

wwPDB NMR Structure Validation Summary Report (i)

Jun 4, 2023 – 11:50 PM EDT

PDB ID : 2LQ5 BMRB ID : 18292

Title : NMR structure of the RNA binding motif 39 (RBM39) from Mus musculus Authors : Serrano, P.; Dutta, S.K.; Geralt, M.; Wuthrich, K.; Joint Center for Structural

Genomics (JCSG); Partnership for T-Cell Biology (TCELL)

Deposited on : 2012-02-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	113	91%	6%	•



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 16 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:4-A:113 (110)	0.75	16		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 5, 8, 9, 12, 13, 15, 16, 17, 18, 19, 20
2	4, 7, 10, 11, 14
3	1, 2, 6



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1607 atoms, of which 729 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called RNA-binding protein 39.

Mol	Chain	Residues		Atoms				Trace	
1	Λ	112	Total	С	Н	N	О	S	0
1	A	113	1607	558	729	150	164	6	U




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: RNA-binding protein 39



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 16. Colouring as in section 4.1 above.

• Molecule 1: RNA-binding protein 39





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: molecular dynamics.

Of the 80 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPALp	refinement	
CYANA	structure solution	
UNIO	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1355
Number of shifts mapped to atoms	1235
Number of unparsed shifts	0
Number of shifts with mapping errors	120
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	E	ond lengths	Bond angles		
IVIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.60 ± 0.01	$0\pm0/876~(~0.0\pm~0.0\%)$	1.02 ± 0.03	$1\pm 1/1194$ ($0.1\pm~0.1\%$)	
All	All	0.60	0/17520 (0.0%)	1.03	14/23880 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	1.1 ± 0.8
All	All	0	22

There are no bond-length outliers.

5 of 8 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	$f hain egin{array}{c c c c c c c c c c c c c c c c c c c $		$\operatorname{Ideal}({}^{o})$	Models				
WIOI	Chain	nes	Type	Atoms		Observed()	ideai()	Worst	Total
1	A	88	TYR	CB-CG-CD2	-5.87	117.48	121.00	16	1
1	A	57	VAL	CA-CB-CG2	5.78	119.56	110.90	4	1
1	A	107	LEU	CB-CG-CD2	5.72	120.72	111.00	9	4
1	A	42	GLY	C-N-CA	5.68	135.90	121.70	20	1
1	A	77	ARG	NE-CZ-NH1	5.52	123.06	120.30	2	1

There are no chirality outliers.

5 of 8 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	112	ARG	Sidechain	6
1	A	88	TYR	Sidechain	4
1	A	94	TYR	Sidechain	4



Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	113	ARG	Sidechain	3
1	A	77	ARG	Sidechain	2

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	855	710	833	2±2
All	All	17100	14200	16660	44

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 29 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:40:HIS:CE1	1:A:73:ALA:HB2	0.68	2.24	6	1
1:A:4:LEU:H	1:A:4:LEU:HD13	0.60	1.56	18	1
1:A:37:CYS:HA	1:A:40:HIS:CD2	0.56	2.35	8	2
1:A:44:ILE:HA	1:A:108:LEU:HD23	0.55	1.77	4	1
1:A:9:PHE:CE2	1:A:59:VAL:HG11	0.54	2.37	16	5

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed Favoured Alle		Allowed	Outliers	Perce	entiles
1	A	109/113 (96%)	94±3 (86±3%)	12±3 (11±3%)	3±1 (2±1%)	9	46
All	All	$2180/2260 \ (96\%)$	1881 (86%)	246 (11%)	53 (2%)	9	46

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of



occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	13	ASN	16
1	A	41	GLY	10
1	A	81	GLY	7
1	A	80	ALA	6
1	A	112	ARG	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles		
1	A	$92/95\ (97\%)$	79±4 (86±4%)	13±4 (14±4%)	7 47		
All	All	1840/1900 (97%)	1586 (86%)	254 (14%)	7 47		

5 of 53 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res		Models (Total)
1	A	45	HIS	13
1	A	79	PHE	13
1	A	50	LYS	12
1	A	112	ARG	12
1	A	82	LYS	11

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.



6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 90% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1355
Number of shifts mapped to atoms	1235
Number of unparsed shifts	0
Number of shifts with mapping errors	120
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

The following errors were found when reading this chemical shift list.

• Chemical shift has been reported more than once. All 3 occurrences are reported below.

Tigt ID	Chain	Dag	Trme	Type Atom		Shift Dat	
LIST ID	Chain	nes	туре	Atom	Value	Uncertainty	Ambiguity
1	A	11	LEU	HD11	0.859	0.000	
1	A	11	LEU	HD12	0.859	0.000	
1	A	11	LEU	HD13	0.859	0.000	•

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 120) occurrences are reported below.

List ID	Chain	Chain Res Typ		Type Atom		Shift Dat	a
LIST ID	Chain	nes	туре	Atom	Value	Uncertainty	Ambiguity
1	A	4	LEU	HB2	1.46	0.000	2
1	A	7	GLN	HB2	1.852	0.000	2
1	A	7	GLN	HG2	2.053	0.000	2
1	A	8	CYS	HB2	2.357	0.000	2
1	A	9	PHE	HB2	2.677	0.000	2



 $Continued\ from\ previous\ page...$

	$\frac{a \text{ from } pro}{a}$			A .		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	10	GLN	HB2	1.776	0.000	2
1	A	10	GLN	HG2	1.525	0.000	2
1	A	11	LEU	HB2	1.731	0.000	2
1	A	12	SER	HB2	3.514	0.000	2
1	A	13	ASN	HB2	3.076	0.000	2
1	A	14	MET	HB2	1.979	0.000	2
1	A	15	PHE	HB2	2.648	0.000	2
1	A	16	ASN	HB2	2.898	0.000	2
1	A	17	PRO	HD2	3.685	0.000	2
1	A	17	PRO	HB2	2.412	0.000	2
1	A	18	GLN	HB2	2.022	0.000	2
1	A	18	GLN	HG2	2.163	0.000	2
1	A	20	GLU	HB2	1.314	0.000	2
1	A	20	GLU	HG2	1.408	0.000	2
1	A	21	GLU	HB2	1.975	0.000	2
1	A	21	GLU	HG2	2.026	0.000	2
1	A	22	GLU	HB2	2.056	0.000	2
1	A	22	GLU	HG2	2.239	0.000	2
1	A	25	TRP	HB2	3.294	0.000	2
1	A	26	ASP	HB2	1.507	0.000	2
1	A	28	GLU	HB2	2.056	0.000	2
1	A	28	GLU	HG2	2.287	0.000	2
1	A	29	ILE	HG12	1.422	0.000	
1	A	30	LYS	HB2	1.683	0.000	2
1	A	30	LYS	HG2	1.231	0.000	2
1	A	30	LYS	HE2	2.636	0.000	2
1	A	30	LYS	HD2	1.458	0.000	2
1	A	31	ASP	HB2	2.758	0.000	2
1	A	32	ASP	HB2	2.554	0.000	2
1	A	34	ILE	HG12	1.797	0.000	•
1	A	35	GLU	HB2	2.113	0.000	2
1	A	35	GLU	HG2	2.317	0.000	2
1	A	36	GLU	HB2	2.254	0.000	2
1	A	36	GLU	HG2	2.349	0.000	2
1	A	37	CYS	HB2	2.805	0.000	2
1	A	38	ASN	HB2	2.834	0.000	2
1	A	39	LYS	HB2	1.42	0.000	2
1	A	39	LYS	HG2	1.19	0.000	2
1	A	39	LYS	HE2	2.782	0.000	2
1	A	39	LYS	HD2	1.574	0.000	2
1	A	40	HIS	HB2	3.459	0.000	2



 $Continued\ from\ previous\ page...$

	$\frac{a \text{ from } pr}{a}$			A .		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	45	HIS	HB2	3.048	0.000	2
1	A	46	ILE	HG12	1.589	0.000	
1	A	47	TYR	HB2	2.113	0.000	2
1	A	49	ASP	HB2	3.573	0.000	2
1	A	50	LYS	HB2	1.831	0.000	2
1	A	50	LYS	HG2	1.327	0.000	2
1	A	50	LYS	HD2	1.486	0.000	2
1	A	50	LYS	HE2	2.664	0.000	2
1	A	51	ASN	HB2	3.006	0.000	2
1	A	52	SER	HB2	4.267	0.000	2
1	A	54	GLN	HB2	2.164	0.000	2
1	A	54	GLN	HG2	2.372	0.000	2
1	A	56	ASN	HB2	2.15	0.000	2
1	A	58	TYR	HB2	3.279	0.000	2
1	A	60	LYS	HB2	1.666	0.000	2
1	A	63	SER	HB2	3.892	0.000	2
1	A	64	ILE	HG12	1.645	0.000	2
1	A	68	ILE	HG12	1.689	0.000	2
1	A	72	ASN	HB2	2.601	0.000	2
1	A	74	LEU	HB2	1.494	0.000	2
1	A	75	HIS	HB2	2.813	0.000	2
1	A	78	TRP	HB2	2.929	0.000	2
1	A	79	PHE	HB2	2.825	0.000	2
1	A	82	LYS	HB2	1.681	0.000	2
1	A	82	LYS	HE2	2.79	0.000	2
1	A	83	MET	HB2	1.307	0.000	2
1	A	84	ILE	HG12	1.215	0.000	2
1	A	88	TYR	HB2	3.282	0.000	2
1	A	92	PRO	HB2	2.264	0.000	2
1	A	94	TYR	HB2	2.595	0.000	2
1	A	95	HIS	HB2	3.024	0.000	2
1	A	108	LEU	HB2	1.266	0.000	2
1	A	100	ASP	HB2	2.689	0.000	2
1	A	102	MET	HB2	1.829	0.000	2
1	A	102	MET	HG2	2.016	0.000	2
1	A	106	GLN	HG2	2.341	0.000	2
1	A	106	GLN	HB2	1.949	0.000	2
1	A	107	LEU	HB2	1.567	0.000	2
1	A	97	LEU	HB2	-0.502	0.000	2
1	A	111	SER	HB2	3.828	0.000	2
1	A	113	ARG	HB2	1.757	0.000	2



 $Continued\ from\ previous\ page...$

	Clasical			A 4		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	61	CYS	HB2	2.852	0.000	2
1	A	112	ARG	HB2	1.737	0.000	2
1	A	112	ARG	HG2	1.474	0.000	2
1	A	14	MET	HG2	2.217	0.000	2
1	A	60	LYS	HE2	2.767	0.000	2
1	A	60	LYS	HG2	1.058	0.000	2
1	A	83	MET	HG2	1.577	0.000	2
1	A	112	ARG	HD2	3.093	0.000	2
1	A	3	PRO	HB2	1.831	0.000	2
1	A	3	PRO	HD2	3.64	0.000	2
1	A	92	PRO	HD2	3.527	0.000	2
1	A	92	PRO	HG2	1.923	0.000	2
1	A	3	PRO	HG2	1.967	0.000	2
1	A	101	SER	HB2	2.515	0.000	2
1	A	2	GLN	HB2	2.026	0.000	2
1	A	2	GLN	HG2	2.341	0.000	2
1	A	77	ARG	HG2	1.596	0.000	2
1	A	77	ARG	HD2	3.09	0.000	2
1	A	62	PRO	HB2	2.152	0.000	2
1	A	62	PRO	HD2	3.171	0.000	2
1	A	62	PRO	HG2	1.843	0.000	2
1	A	110	PRO	HB2	2.024	0.000	2
1	A	110	PRO	HD2	3.555	0.000	2
1	A	99	PRO	HB2	2.29	0.000	2
1	A	99	PRO	HD2	3.175	0.000	2
1	A	110	PRO	HG2	1.695	0.000	2
1	A	90	PRO	HD2	3.99	0.000	2
1	A	90	PRO	HG2	2.157	0.000	2
1	A	90	PRO	HB2	1.87	0.000	2
1	A	91	LEU	HB2	1.588	0.000	2
1	A	98	PHE	HB2	2.37	0.000	2
1	A	44	ILE	HG12	0.947	0.000	2
1	A	96	ASN	HB2	2.753	0.000	2

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	112	2.58 ± 0.19	Should be applied
$^{13}C_{\beta}$	106	2.63 ± 0.13	Should be applied



Continued from previous page...

Nucleus	# values	Correction \pm precision, ppm	Suggested action
¹³ C′	99	2.81 ± 0.20	Should be applied
^{15}N	105	0.62 ± 0.26	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 1329 atoms were assigned a chemical shift out of a possible 1464. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	531/544 (98%)	$219/220 \ (100\%)$	$208/220 \ (95\%)$	104/104 (100%)
Sidechain	703/792 (89%)	475/519 (92%)	216/247 (87%)	12/26 (46%)
Aromatic	95/128 (74%)	49/64 (77%)	44/58 (76%)	2/6 (33%)
Overall	1329/1464 (91%)	743/803 (93%)	468/525 (89%)	118/136 (87%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	26	ASP	HB3	0.17	1.32 - 4.00	-9.3
1	A	59	VAL	НВ	-0.06	0.43 - 3.54	-6.6
1	A	97	LEU	HB2	-0.50	-0.07 - 3.30	-6.3
1	A	55	GLY	HA2	1.80	2.15 - 5.77	-6.0
1	A	101	SER	HB2	2.52	2.61 - 5.13	-5.4

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2121
Intra-residue ($ i-j =0$)	439
Sequential ($ i-j =1$)	607
Medium range ($ i-j >1$ and $ i-j <5$)	428
Long range (i-j ≥5)	647
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	577
Number of restraints per residue	18.8
Number of long range restraints per residue ¹	5.7

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	25.9	0.2
0.2-0.5 (Medium)	49.6	0.5
>0.5 (Large)	111.5	3.54



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dordensinda dom o	Count	% ¹	Vi	olated	3	Consis	tentl	${ m y}$ Violated 4
Restraints type	Count	70	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	439	20.7	74	16.9	3.5	1	0.2	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	361	17.0	65	18.0	3.1	1	0.3	0.0
Sidechain-Sidechain	78	3.7	9	11.5	0.4	0	0.0	0.0
Sequential (i-j =1)	607	28.6	99	16.3	4.7	17	2.8	0.8
Backbone-Backbone	169	8.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	352	16.6	73	20.7	3.4	11	3.1	0.5
Sidechain-Sidechain	86	4.1	26	30.2	1.2	6	7.0	0.3
Medium range ($ i-j >1 \& i-j <5$)	428	20.2	79	18.5	3.7	8	1.9	0.4
Backbone-Backbone	151	7.1	1	0.7	0.0	1	0.7	0.0
Backbone-Sidechain	181	8.5	47	26.0	2.2	5	2.8	0.2
Sidechain-Sidechain	96	4.5	31	32.3	1.5	2	2.1	0.1
Long range ($ i-j \ge 5$)	647	30.5	155	24.0	7.3	3	0.5	0.1
Backbone-Backbone	88	4.1	5	5.7	0.2	0	0.0	0.0
Backbone-Sidechain	301	14.2	80	26.6	3.8	1	0.3	0.0
Sidechain-Sidechain	258	12.2	70	27.1	3.3	2	0.8	0.1
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2121	100.0	407	19.2	19.2	29	1.4	1.4
Backbone-Backbone	408	19.2	6	1.5	0.3	1	0.2	0.0
Backbone-Sidechain	1195	56.3	265	22.2	12.5	18	1.5	0.8
Sidechain-Sidechain	518	24.4	136	26.3	6.4	10	1.9	0.5

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	M (Å)	Max (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (A)	\mathbf{SD}^6 (Å)	Median (Å)
1	38	50	47	58	0	193	0.74	2.98	0.55	0.62
2	35	54	51	71	0	211	0.8	3.27	0.55	0.72
3	31	60	42	73	0	206	0.85	3.44	0.61	0.68
4	35	50	35	68	0	188	0.76	2.67	0.57	0.58
5	34	40	46	58	0	178	0.76	2.47	0.56	0.62
6	32	48	44	64	0	188	0.81	3.09	0.61	0.61
7	32	51	49	83	0	215	0.86	3.07	0.6	0.71
8	33	50	42	61	0	186	0.8	3.42	0.57	0.63
9	33	48	43	71	0	195	0.82	2.78	0.61	0.63
10	31	39	36	53	0	159	0.7	2.99	0.51	0.57
11	31	46	36	77	0	190	0.77	2.64	0.59	0.6




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	34	45	40	73	0	192	0.77	2.57	0.58	0.59
13	37	56	52	68	0	213	0.74	2.94	0.54	0.62
14	31	43	41	60	0	175	0.83	3.54	0.62	0.67
15	32	43	45	56	0	176	0.74	2.64	0.54	0.6
16	32	41	39	52	0	164	0.69	3.13	0.52	0.55
17	35	52	45	66	0	198	0.78	3.22	0.59	0.62
18	30	45	36	57	0	168	0.8	2.82	0.59	0.67
19	33	40	39	44	0	156	0.65	2.71	0.5	0.56
20	31	54	37	68	0	190	0.82	3.02	0.62	0.7

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1714(IR:365, SQ:508, MR:349, LR:492, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
10	13	4	28	0	55	1	5.0
3	7	7	19	0	36	2	10.0
7	6	2	9	0	24	3	15.0
4	7	3	5	0	19	4	20.0
5	3	3	9	0	20	5	25.0
2	3	1	9	0	15	6	30.0
1	6	7	2	0	16	7	35.0
4	3	7	6	0	20	8	40.0
6	4	3	1	0	14	9	45.0
2	3	2	7	0	14	10	50.0
5	2	6	7	0	20	11	55.0
3	1	1	4	0	9	12	60.0
2	14	6	9	0	31	13	65.0
3	3	2	5	0	13	14	70.0
2	2	1	8	0	13	15	75.0
2	1	4	4	0	11	16	80.0
4	1	5	4	0	14	17	85.0
4	1	5	0	0	10	18	90.0
4	2	2	16	0	24	19	95.0
1	17	8	3	0	29	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,90)	1:A:47:TYR:HE2	1:A:50:LYS:H	20	2.16	0.59	2.26
(1,2110)	1:A:47:TYR:HE2	1:A:49:ASP:HA	20	1.86	0.51	1.88
(1,2116)	1:A:47:TYR:HE1	1:A:49:ASP:HB3	20	1.49	0.36	1.58
(1,7)	1:A:47:TYR:HE2	1:A:48:VAL:H	20	1.4	0.26	1.38
(1,809)	1:A:30:LYS:HD3	1:A:31:ASP:HB3	20	1.39	0.56	1.53
(1,661)	1:A:10:GLN:HE22	1:A:57:VAL:HA	20	1.34	0.2	1.38
(1,334)	1:A:27:THR:H	1:A:28:GLU:HB3	20	1.28	0.43	1.44
(1,2026)	1:A:87:ALA:HA	1:A:88:TYR:HB3	20	1.27	0.06	1.27
(1,1116)	1:A:28:GLU:HB3	1:A:29:ILE:HB	20	1.21	0.24	1.25
(1,242)	1:A:30:LYS:HD3	1:A:32:ASP:H	20	1.17	0.41	1.37


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1025)	1:A:112:ARG:HD3	1:A:113:ARG:HG3	14	3.54
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG11	3	3.44
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG12	3	3.44
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG13	3	3.44
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG11	3	3.44
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG12	3	3.44
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG13	3	3.44
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG11	3	3.44
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG12	3	3.44
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG13	3	3.44
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG11	8	3.42



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG12	8	3.42
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG13	8	3.42
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG11	8	3.42
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG12	8	3.42
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG13	8	3.42
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG11	8	3.42
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG12	8	3.42
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG13	8	3.42
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG11	2	3.27
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG12	2	3.27
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG13	2	3.27
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG11	2	3.27
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG12	2	3.27
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG13	2	3.27
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG11	2	3.27
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG12	2	3.27
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG13	2	3.27
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG11	17	3.22
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG12	17	3.22
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG13	17	3.22
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG11	17	3.22
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG12	17	3.22
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG13	17	3.22
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG11	17	3.22
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG12	17	3.22
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG13	17	3.22
(1,616)	1:A:16:ASN:HD22	1:A:18:GLN:HB3	16	3.13
(1,616)	1:A:16:ASN:HD22	1:A:18:GLN:HB3	14	3.11
(1,90)	1:A:47:TYR:HE2	1:A:50:LYS:H	6	3.09
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG11	7	3.07
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG12	7	3.07
(1,1501)	1:A:57:VAL:HG21	1:A:59:VAL:HG13	7	3.07
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG11	7	3.07
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG12	7	3.07
(1,1501)	1:A:57:VAL:HG22	1:A:59:VAL:HG13	7	3.07
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG11	7	3.07
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG12	7	3.07
(1,1501)	1:A:57:VAL:HG23	1:A:59:VAL:HG13	7	3.07
(1,616)	1:A:16:ASN:HD22	1:A:18:GLN:HB3	20	3.02



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

