



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2024 – 11:17 AM EDT

PDB ID : 1LSM
Title : THERMAL STABILITY DETERMINANTS OF CHICKEN EGG-WHITE
LYSOZYME CORE MUTANTS: HYDROPHOBICITY, PACKING VOL-
UME AND CONSERVED BURIED WATER MOLECULES
Authors : Holland, D.R.; Shih, P.
Deposited on : 1994-09-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	129	 76% 19% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEN EGG WHITE LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	1000	613	193	184	10	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	LEU	ILE	conflict	UNP P00698
A	91	THR	SER	conflict	UNP P00698
A	101	SER	ASP	conflict	UNP P00698

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	170	Total	O	0	0
			170	170		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HEN EGG WHITE LYSOZYME



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.08Å 79.08Å 37.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.70	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.160 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1170	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.06	2/1020 (0.2%)	1.36	17/1378 (1.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	ARG	C-N	18.20	1.75	1.34
1	A	7	GLU	CD-OE2	5.05	1.31	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	ARG	NE-CZ-NH1	8.64	124.62	120.30
1	A	52	ASP	CB-CG-OD2	-7.47	111.58	118.30
1	A	48	ASP	CB-CG-OD2	-7.09	111.92	118.30
1	A	66	ASP	CB-CG-OD2	-6.45	112.50	118.30
1	A	48	ASP	CB-CG-OD1	6.26	123.94	118.30
1	A	119	ASP	CB-CG-OD2	-5.97	112.93	118.30
1	A	78	ILE	CB-CA-C	-5.87	99.86	111.60
1	A	128	ARG	NE-CZ-NH1	5.79	123.20	120.30
1	A	87	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	A	112	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	52	ASP	CB-CG-OD1	5.71	123.44	118.30
1	A	66	ASP	CB-CG-OD1	5.62	123.36	118.30
1	A	18	ASP	CB-CG-OD1	5.62	123.35	118.30
1	A	73	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	A	87	ASP	CB-CG-OD1	5.22	123.00	118.30
1	A	18	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	125	ARG	NE-CZ-NH2	-5.18	117.71	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1000	0	961	22	0
2	A	170	0	0	3	1
All	All	1170	0	961	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ARG:C	1:A:129:LEU:N	1.75	1.35
1:A:125:ARG:HH11	1:A:125:ARG:HG2	1.48	0.78
1:A:45:ARG:NH1	2:A:316:HOH:O	2.28	0.66
1:A:125:ARG:HH11	1:A:125:ARG:CG	2.13	0.62
1:A:45:ARG:NH2	1:A:68:ARG:NH1	2.49	0.60
1:A:62:TRP:HB2	1:A:63:TRP:CD1	2.40	0.56
1:A:125:ARG:HG2	1:A:125:ARG:NH1	2.20	0.55
1:A:45:ARG:NE	1:A:68:ARG:NH2	2.57	0.51
1:A:128:ARG:C	1:A:129:LEU:CA	2.73	0.51
1:A:111:TRP:CD1	1:A:115:CYS:HB2	2.46	0.51
1:A:61:ARG:HH11	1:A:61:ARG:HG3	1.76	0.51
1:A:128:ARG:CA	1:A:129:LEU:N	2.69	0.50
1:A:52:ASP:HB3	1:A:57:GLN:HB3	1.94	0.48
1:A:45:ARG:NH2	1:A:68:ARG:CZ	2.78	0.47
1:A:45:ARG:HE	1:A:68:ARG:NH2	2.13	0.46
1:A:15:HIS:HB3	1:A:92:VAL:HG11	1.99	0.45
1:A:121:GLN:HE21	1:A:121:GLN:HB3	1.41	0.44
1:A:63:TRP:CE2	1:A:98:ILE:HG12	2.53	0.44
1:A:125:ARG:CG	1:A:125:ARG:NH1	2.79	0.43
1:A:17:LEU:HA	1:A:17:LEU:HD23	1.74	0.41
1:A:18:ASP:HB3	2:A:315:HOH:O	2.19	0.41
1:A:44:ASN:HB3	2:A:204:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:203:HOH:O	2:A:203:HOH:O[7_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/129 (98%)	124 (98%)	3 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	105/105 (100%)	98 (93%)	7 (7%)	16 4

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	77	ASN
1	A	78	ILE
1	A	91	THR
1	A	101	SER
1	A	121	GLN
1	A	125	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	93	ASN
1	A	121	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	128:ARG	C	129:LEU	N	1.75

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.