



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 19, 2023 – 04:23 PM JST

PDB ID : 6LXJ  
Title : Crystal structure of human Z2B3 Fab in complex with influenza virus neuraminidase from A/Anhui/1/2013 (H7N9)  
Authors : Jiang, H.; Peng, W.; Qi, J.; Chai, Y.; Song, H.; Shi, Y.; Gao, G.F.; Wu, Y.  
Deposited on : 2020-02-11  
Resolution : 2.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	 86% 13%
1	B	388	 80% 19%
1	C	388	 84% 14%
1	D	388	 80% 19%
2	E	237	 % 72% 17% 10%
2	H	237	 2% 74% 21% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	J	237	 2% 67% 19% 13%
2	M	237	 2% 68% 19% 12%
3	F	216	 4% 71% 23% ..
3	K	216	 3% 64% 30% 6%
3	L	216	 73% 25% ..
3	N	216	 4% 61% 34% ..
4	G	5	 40% 40% 20%
5	I	9	 56% 33% 11%
5	P	9	 33% 56% 11%
5	R	9	 56% 22% 22%
5	T	9	 56% 22% 22%
6	O	4	 50% 50%
6	Q	4	 25% 75%
6	S	4	 75% 25%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 25523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3055	1902	541	589	23	0	0	0
1	B	388	3055	1902	541	589	23	0	0	0
1	C	388	3055	1902	541	589	23	0	0	0
1	D	388	3055	1902	541	589	23	0	0	0


- Molecule 2 is a protein called Heavy chain of Z2B3 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	226	1694	1074	275	338	7	0	0	0
2	E	214	1614	1026	262	319	7	0	0	0
2	J	206	1561	993	253	308	7	0	0	0
2	M	208	1563	994	255	307	7	0	0	0

- Molecule 3 is a protein called Light chain of Z2B3 Fab.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1579	982	262	330	5	0	0	0
3	F	207	1540	960	256	319	5	0	0	0
3	K	204	1517	948	251	313	5	0	0	0
3	N	208	1544	962	257	320	5	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	5	61	34	2	25	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	I	9	105	58	2	45	0	0	0
5	P	9	105	58	2	45	0	0	0
5	R	9	105	58	2	45	0	0	0
5	T	9	105	58	2	45	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	O	4	50	28	2	20	0	0	0
6	Q	4	50	28	2	20	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	S	4	50	28	2	20	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	A	1	14	8	1	5	0	0
7	B	1	14	8	1	5	0	0
7	C	1	14	8	1	5	0	0
7	D	1	14	8	1	5	0	0

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
8	A	1	1	1	0	0
8	B	1	1	1	0	0
8	C	1	1	1	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
8	D	1	Total 1	Ca 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neuraminidase




- Molecule 1: Neuraminidase




- Molecule 1: Neuraminidase





- Molecule 1: Neuraminidase









- Molecule 2: Heavy chain of Z2B3 Fab



- Molecule 2: Heavy chain of Z2B3 Fab





- Molecule 2: Heavy chain of Z2B3 Fab



SER  
CYS  
ASP  
LYS

- Molecule 2: Heavy chain of Z2B3 Fab





LYS

• Molecule 3: Light chain of Z2B3 Fab



• Molecule 3: Light chain of Z2B3 Fab



• Molecule 3: Light chain of Z2B3 Fab



P212  
THR  
GLU  
CYS  
SER

• Molecule 3: Light chain of Z2B3 Fab






- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 40% 40% 20%



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 56% 33% 11%




- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P: 33% 56% 11%



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain R: 56% 22% 22%



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-

(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain T:  56% 22% 22%




- Molecule 6: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  50% 50%



- Molecule 6: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Q:  25% 75%



- Molecule 6: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S:  75% 25%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	201.31Å 207.51Å 207.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.91 – 2.90 48.91 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.91-2.90) 99.1 (48.91-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 2.91Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.193 , 0.241 0.193 , 0.242	Depositor DCC
$R_{free}$ test set	4742 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.8	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 27.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25523	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/3137	0.65	0/4271
1	B	0.65	0/3137	0.66	0/4271
1	C	0.66	0/3137	0.69	0/4271
1	D	0.71	0/3137	0.68	0/4271
2	E	0.67	0/1652	0.62	0/2251
2	H	0.65	0/1735	0.62	0/2366
2	J	0.61	0/1598	0.67	0/2175
2	M	0.58	0/1600	0.59	0/2179
3	F	0.52	0/1575	0.61	0/2145
3	K	0.53	0/1551	0.61	0/2111
3	L	0.62	0/1616	0.62	0/2205
3	N	0.52	0/1580	0.64	0/2154
All	All	0.63	0/25455	0.65	0/34670

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3055	0	2877	40	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3055	0	2877	53	0
1	C	3055	0	2877	50	0
1	D	3055	0	2877	53	0
2	E	1614	0	1574	35	0
2	H	1694	0	1657	39	0
2	J	1561	0	1516	38	0
2	M	1563	0	1528	34	0
3	F	1540	0	1488	37	0
3	K	1517	0	1465	52	0
3	L	1579	0	1518	38	0
3	N	1544	0	1488	54	0
4	G	61	0	52	2	0
5	I	105	0	88	1	0
5	P	105	0	88	3	0
5	R	105	0	88	2	0
5	T	105	0	88	2	0
6	O	50	0	43	3	0
6	Q	50	0	43	4	0
6	S	50	0	43	0	0
7	A	14	0	13	0	0
7	B	14	0	13	1	0
7	C	14	0	13	0	0
7	D	14	0	13	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
All	All	25523	0	24327	496	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 496 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:K:27:SER:HB2	3:K:95:ARG:HG3	1.46	0.97
1:C:437:TRP:CD1	6:Q:1:NAG:H82	2.00	0.97
2:J:228:LYS:NZ	3:K:127:GLU:OE1	1.99	0.94
1:C:123:SER:OG	1:C:190:ARG:NH2	2.01	0.93
1:D:121:TYR:CD1	1:D:228:GLU:O	2.23	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	366 (95%)	19 (5%)	1 (0%)	41	71
1	B	386/388 (100%)	369 (96%)	16 (4%)	1 (0%)	41	71
1	C	386/388 (100%)	365 (95%)	21 (5%)	0	100	100
1	D	386/388 (100%)	366 (95%)	20 (5%)	0	100	100
2	E	208/237 (88%)	197 (95%)	11 (5%)	0	100	100
2	H	222/237 (94%)	217 (98%)	5 (2%)	0	100	100
2	J	198/237 (84%)	192 (97%)	6 (3%)	0	100	100
2	M	202/237 (85%)	194 (96%)	8 (4%)	0	100	100
3	F	201/216 (93%)	180 (90%)	20 (10%)	1 (0%)	29	61
3	K	196/216 (91%)	186 (95%)	9 (5%)	1 (0%)	29	61
3	L	211/216 (98%)	203 (96%)	8 (4%)	0	100	100
3	N	204/216 (94%)	190 (93%)	13 (6%)	1 (0%)	29	61
All	All	3186/3364 (95%)	3025 (95%)	156 (5%)	5 (0%)	47	78

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	K	53	VAL
1	A	223	ILE
3	N	53	VAL
1	B	223	ILE
3	F	53	VAL



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	339/339 (100%)	335 (99%)	4 (1%)	71 91
1	B	339/339 (100%)	330 (97%)	9 (3%)	44 77
1	C	339/339 (100%)	328 (97%)	11 (3%)	39 73
1	D	339/339 (100%)	329 (97%)	10 (3%)	42 76
2	E	181/200 (90%)	174 (96%)	7 (4%)	32 66
2	H	191/200 (96%)	187 (98%)	4 (2%)	53 81
2	J	176/200 (88%)	167 (95%)	9 (5%)	24 56
2	M	175/200 (88%)	166 (95%)	9 (5%)	24 56
3	F	176/183 (96%)	170 (97%)	6 (3%)	37 71
3	K	173/183 (94%)	167 (96%)	6 (4%)	36 70
3	L	180/183 (98%)	172 (96%)	8 (4%)	28 61
3	N	176/183 (96%)	168 (96%)	8 (4%)	27 61
All	All	2784/2888 (96%)	2693 (97%)	91 (3%)	38 72

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	191	SER
3	K	67	SER
2	J	4	LEU
2	J	197	LEU
2	M	22	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	K	171	GLN
3	K	174	ASN
2	M	117	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

53 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	G	1	1,4	14,14,15	0.29	0	17,19,21	0.71	0
4	NAG	G	2	4	14,14,15	0.29	0	17,19,21	0.64	0
4	BMA	G	3	4	11,11,12	0.73	0	15,15,17	1.93	3 (20%)
4	MAN	G	4	4	11,11,12	0.26	0	15,15,17	0.64	0
4	MAN	G	5	4	11,11,12	0.26	0	15,15,17	0.65	0
5	NAG	I	1	1,5	14,14,15	0.56	0	17,19,21	0.69	0
5	NAG	I	2	5	14,14,15	0.57	0	17,19,21	0.68	0
5	BMA	I	3	5	11,11,12	0.27	0	15,15,17	0.97	1 (6%)
5	MAN	I	4	5	11,11,12	0.48	0	15,15,17	1.39	2 (13%)
5	MAN	I	5	5	11,11,12	0.51	0	15,15,17	0.79	0
5	MAN	I	6	5	11,11,12	0.43	0	15,15,17	0.97	1 (6%)
5	MAN	I	7	5	11,11,12	0.57	0	15,15,17	0.75	0
5	MAN	I	8	5	11,11,12	0.55	0	15,15,17	0.78	0
5	MAN	I	9	5	11,11,12	0.57	0	15,15,17	1.05	2 (13%)
6	NAG	O	1	1,6	14,14,15	0.29	0	17,19,21	0.74	0
6	NAG	O	2	6	14,14,15	0.30	0	17,19,21	0.64	0
6	BMA	O	3	6	11,11,12	0.33	0	15,15,17	0.69	0
6	MAN	O	4	6	11,11,12	0.26	0	15,15,17	0.64	0
5	NAG	P	1	1,5	14,14,15	0.56	0	17,19,21	0.68	0
5	NAG	P	2	5	14,14,15	0.57	0	17,19,21	0.68	0
5	BMA	P	3	5	11,11,12	0.26	0	15,15,17	0.97	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MAN	P	4	5	11,11,12	0.47	0	15,15,17	1.38	2 (13%)
5	MAN	P	5	5	11,11,12	0.50	0	15,15,17	0.78	0
5	MAN	P	6	5	11,11,12	0.43	0	15,15,17	0.98	1 (6%)
5	MAN	P	7	5	11,11,12	0.58	0	15,15,17	0.76	0
5	MAN	P	8	5	11,11,12	0.55	0	15,15,17	0.78	0
5	MAN	P	9	5	11,11,12	0.57	0	15,15,17	1.06	2 (13%)
6	NAG	Q	1	1,6	14,14,15	0.31	0	17,19,21	0.78	0
6	NAG	Q	2	6	14,14,15	0.29	0	17,19,21	0.76	0
6	BMA	Q	3	6	11,11,12	0.36	0	15,15,17	0.94	1 (6%)
6	MAN	Q	4	6	11,11,12	0.27	0	15,15,17	0.63	0
5	NAG	R	1	1,5	14,14,15	0.55	0	17,19,21	0.69	0
5	NAG	R	2	5	14,14,15	0.57	0	17,19,21	0.68	0
5	BMA	R	3	5	11,11,12	0.26	0	15,15,17	0.97	1 (6%)
5	MAN	R	4	5	11,11,12	0.48	0	15,15,17	1.39	2 (13%)
5	MAN	R	5	5	11,11,12	0.52	0	15,15,17	0.79	0
5	MAN	R	6	5	11,11,12	0.44	0	15,15,17	0.98	1 (6%)
5	MAN	R	7	5	11,11,12	0.57	0	15,15,17	0.75	0
5	MAN	R	8	5	11,11,12	0.54	0	15,15,17	0.78	0
5	MAN	R	9	5	11,11,12	0.57	0	15,15,17	1.06	2 (13%)
6	NAG	S	1	1,6	14,14,15	0.28	0	17,19,21	0.77	0
6	NAG	S	2	6	14,14,15	0.29	0	17,19,21	0.64	0
6	BMA	S	3	6	11,11,12	0.43	0	15,15,17	1.26	2 (13%)
6	MAN	S	4	6	11,11,12	0.23	0	15,15,17	0.76	0
5	NAG	T	1	1,5	14,14,15	0.56	0	17,19,21	0.68	0
5	NAG	T	2	5	14,14,15	0.57	0	17,19,21	0.68	0
5	BMA	T	3	5	11,11,12	0.27	0	15,15,17	0.97	1 (6%)
5	MAN	T	4	5	11,11,12	0.46	0	15,15,17	1.38	2 (13%)
5	MAN	T	5	5	11,11,12	0.50	0	15,15,17	0.79	0
5	MAN	T	6	5	11,11,12	0.43	0	15,15,17	0.97	1 (6%)
5	MAN	T	7	5	11,11,12	0.56	0	15,15,17	0.75	0
5	MAN	T	8	5	11,11,12	0.54	0	15,15,17	0.77	0
5	MAN	T	9	5	11,11,12	0.58	0	15,15,17	1.05	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	1/6/23/26	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
4	MAN	G	5	4	-	1/2/19/22	0/1/1/1
5	NAG	I	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	I	2	5	-	0/6/23/26	0/1/1/1
5	BMA	I	3	5	-	0/2/19/22	0/1/1/1
5	MAN	I	4	5	-	0/2/19/22	0/1/1/1
5	MAN	I	5	5	-	0/2/19/22	0/1/1/1
5	MAN	I	6	5	-	0/2/19/22	0/1/1/1
5	MAN	I	7	5	-	0/2/19/22	0/1/1/1
5	MAN	I	8	5	-	0/2/19/22	0/1/1/1
5	MAN	I	9	5	-	1/2/19/22	0/1/1/1
6	NAG	O	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	O	2	6	-	1/6/23/26	0/1/1/1
6	BMA	O	3	6	-	1/2/19/22	0/1/1/1
6	MAN	O	4	6	-	2/2/19/22	0/1/1/1
5	NAG	P	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	P	2	5	-	0/6/23/26	0/1/1/1
5	BMA	P	3	5	-	0/2/19/22	0/1/1/1
5	MAN	P	4	5	-	0/2/19/22	0/1/1/1
5	MAN	P	5	5	-	0/2/19/22	0/1/1/1
5	MAN	P	6	5	-	0/2/19/22	0/1/1/1
5	MAN	P	7	5	-	0/2/19/22	0/1/1/1
5	MAN	P	8	5	-	0/2/19/22	0/1/1/1
5	MAN	P	9	5	-	1/2/19/22	0/1/1/1
6	NAG	Q	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	Q	2	6	-	2/6/23/26	0/1/1/1
6	BMA	Q	3	6	-	0/2/19/22	0/1/1/1
6	MAN	Q	4	6	-	2/2/19/22	0/1/1/1
5	NAG	R	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	R	2	5	-	0/6/23/26	0/1/1/1
5	BMA	R	3	5	-	0/2/19/22	0/1/1/1
5	MAN	R	4	5	-	0/2/19/22	0/1/1/1
5	MAN	R	5	5	-	0/2/19/22	0/1/1/1
5	MAN	R	6	5	-	0/2/19/22	0/1/1/1
5	MAN	R	7	5	-	0/2/19/22	0/1/1/1
5	MAN	R	8	5	-	0/2/19/22	0/1/1/1
5	MAN	R	9	5	-	1/2/19/22	0/1/1/1
6	NAG	S	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	S	2	6	-	0/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BMA	S	3	6	-	0/2/19/22	0/1/1/1
6	MAN	S	4	6	-	2/2/19/22	0/1/1/1
5	NAG	T	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	T	2	5	-	0/6/23/26	0/1/1/1
5	BMA	T	3	5	-	0/2/19/22	0/1/1/1
5	MAN	T	4	5	-	0/2/19/22	0/1/1/1
5	MAN	T	5	5	-	0/2/19/22	0/1/1/1
5	MAN	T	6	5	-	0/2/19/22	0/1/1/1
5	MAN	T	7	5	-	0/2/19/22	0/1/1/1
5	MAN	T	8	5	-	0/2/19/22	0/1/1/1
5	MAN	T	9	5	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	3	BMA	C1-C2-C3	5.18	116.04	109.67
4	G	3	BMA	C2-C3-C4	3.67	117.24	110.89
5	P	4	MAN	O2-C2-C1	-2.75	103.53	109.15
5	I	4	MAN	O2-C2-C1	-2.74	103.56	109.15
5	T	4	MAN	O2-C2-C1	-2.73	103.56	109.15

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	S	4	MAN	O5-C5-C6-O6
6	Q	1	NAG	C4-C5-C6-O6
6	S	1	NAG	C4-C5-C6-O6
6	S	4	MAN	C4-C5-C6-O6
6	O	1	NAG	C4-C5-C6-O6

There are no ring outliers.

15 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Q	4	MAN	1	0
5	T	3	BMA	1	0
5	P	5	MAN	1	0
4	G	5	MAN	1	0


*Continued on next page...*


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	3	BMA	1	0
5	P	6	MAN	1	0
6	O	4	MAN	1	0
4	G	4	MAN	1	0
5	T	6	MAN	1	0
5	R	6	MAN	1	0
5	P	1	NAG	1	0
5	R	3	BMA	1	0
4	G	3	BMA	1	0
6	O	3	BMA	2	0
6	Q	1	NAG	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





















## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	NAG	C	501	1	14,14,15	0.44	0	17,19,21	0.86	1 (5%)
7	NAG	D	501	1	14,14,15	0.44	0	17,19,21	0.86	1 (5%)
7	NAG	A	501	1	14,14,15	0.44	0	17,19,21	0.86	1 (5%)
7	NAG	B	501	1	14,14,15	0.44	0	17,19,21	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	C	501	1	-	0/6/23/26	0/1/1/1
7	NAG	D	501	1	-	0/6/23/26	0/1/1/1
7	NAG	A	501	1	-	0/6/23/26	0/1/1/1
7	NAG	B	501	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	501	NAG	O5-C5-C6	2.53	111.17	107.20
7	D	501	NAG	O5-C5-C6	2.52	111.15	107.20
7	B	501	NAG	O5-C5-C6	2.51	111.14	107.20
7	A	501	NAG	O5-C5-C6	2.51	111.14	107.20

There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	501	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9	
1	A	388/388 (100%)	-0.30	0	100 100	17, 27, 41, 62	0
1	B	388/388 (100%)	-0.28	0	100 100	19, 31, 53, 62	0
1	C	388/388 (100%)	-0.29	0	100 100	17, 29, 46, 62	0
1	D	388/388 (100%)	-0.31	0	100 100	16, 27, 43, 55	0
2	E	214/237 (90%)	-0.06	2 (0%)	84 84	27, 39, 64, 78	0
2	H	226/237 (95%)	-0.17	4 (1%)	68 67	22, 33, 62, 84	0
2	J	206/237 (86%)	-0.13	4 (1%)	66 65	26, 39, 75, 96	0
2	M	208/237 (87%)	-0.08	4 (1%)	66 65	24, 37, 70, 97	0
3	F	207/216 (95%)	0.18	8 (3%)	39 35	35, 57, 96, 114	0
3	K	204/216 (94%)	0.20	6 (2%)	51 47	35, 61, 90, 113	0
3	L	213/216 (98%)	-0.15	1 (0%)	91 91	23, 43, 66, 77	0
3	N	208/216 (96%)	0.27	9 (4%)	35 31	31, 59, 92, 120	0
All	All	3238/3364 (96%)	-0.14	38 (1%)	79 79	16, 35, 77, 120	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	J	213	TYR	5.6
3	F	82	ALA	4.0
2	M	178	LEU	3.5
2	H	26	GLY	3.4
2	M	230	VAL	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MAN	R	9	11/12	0.71	0.27	70,78,90,98	0
5	MAN	P	9	11/12	0.72	0.25	56,63,72,73	0
4	MAN	G	5	11/12	0.75	0.28	71,89,101,103	0
4	MAN	G	4	11/12	0.80	0.27	80,92,97,99	0
5	MAN	I	9	11/12	0.84	0.22	54,56,61,67	0
6	MAN	Q	4	11/12	0.84	0.19	51,69,73,74	0
5	MAN	T	9	11/12	0.85	0.21	45,54,61,65	0
5	MAN	T	8	11/12	0.85	0.17	37,47,56,56	0
6	BMA	S	3	11/12	0.85	0.12	54,58,63,66	0
6	BMA	Q	3	11/12	0.86	0.11	50,57,62,62	0
6	BMA	O	3	11/12	0.87	0.10	54,64,66,67	0
4	BMA	G	3	11/12	0.87	0.16	67,77,86,88	0
5	MAN	P	8	11/12	0.88	0.24	42,52,56,57	0
5	MAN	I	8	11/12	0.88	0.19	46,52,57,62	0
6	MAN	O	4	11/12	0.88	0.14	64,70,76,77	0
6	MAN	S	4	11/12	0.88	0.17	61,66,75,78	0
5	BMA	R	3	11/12	0.90	0.18	29,38,45,47	0
4	NAG	G	2	14/15	0.90	0.14	46,56,64,70	0
5	MAN	P	5	11/12	0.90	0.25	36,42,47,48	0
6	NAG	Q	1	14/15	0.90	0.16	25,32,35,37	0
6	NAG	S	2	14/15	0.91	0.15	35,42,52,61	0
5	BMA	P	3	11/12	0.91	0.15	35,39,44,45	0
5	MAN	R	7	11/12	0.91	0.14	38,44,54,66	0
5	MAN	R	5	11/12	0.92	0.30	37,40,46,47	0
6	NAG	O	2	14/15	0.92	0.14	39,56,62,66	0
5	MAN	P	6	11/12	0.92	0.18	37,41,45,48	0
5	MAN	P	4	11/12	0.92	0.18	28,31,38,38	0
5	MAN	R	4	11/12	0.92	0.19	26,32,39,41	0
5	MAN	I	5	11/12	0.93	0.15	42,44,50,52	0
5	MAN	I	7	11/12	0.93	0.15	39,49,52,59	0
5	MAN	R	8	11/12	0.93	0.20	44,45,51,51	0
5	MAN	P	7	11/12	0.94	0.13	42,47,55,57	0
5	MAN	R	6	11/12	0.94	0.17	32,35,37,39	0
5	MAN	I	6	11/12	0.94	0.13	38,43,47,48	0
5	MAN	I	4	11/12	0.94	0.13	27,33,41,43	0
5	NAG	P	1	14/15	0.94	0.13	27,31,41,48	0
6	NAG	Q	2	14/15	0.94	0.12	33,49,52,54	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	T	1	14/15	0.94	0.17	23,27,35,39	0
5	MAN	T	5	11/12	0.94	0.19	34,37,42,45	0
5	MAN	T	6	11/12	0.94	0.17	29,35,40,40	0
5	MAN	T	7	11/12	0.94	0.15	44,50,55,62	0
5	NAG	P	2	14/15	0.94	0.17	28,33,39,49	0
5	BMA	I	3	11/12	0.95	0.16	32,40,46,49	0
5	NAG	I	2	14/15	0.95	0.15	29,30,44,46	0
5	NAG	T	2	14/15	0.95	0.18	24,30,36,50	0
5	NAG	R	1	14/15	0.95	0.15	28,32,43,50	0
6	NAG	O	1	14/15	0.95	0.13	29,37,42,43	0
5	NAG	I	1	14/15	0.96	0.11	25,32,35,38	0
5	NAG	R	2	14/15	0.96	0.19	30,33,38,49	0
5	BMA	T	3	11/12	0.96	0.16	32,35,43,47	0
5	MAN	T	4	11/12	0.96	0.12	25,32,34,35	0
4	NAG	G	1	14/15	0.96	0.12	27,32,41,47	0
6	NAG	S	1	14/15	0.97	0.12	23,26,34,37	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain P:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain R:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain T:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain O:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain Q:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	CA	D	502	1/1	0.60	0.19	90,90,90,90	0
8	CA	B	502	1/1	0.71	0.17	79,79,79,79	0
8	CA	C	502	1/1	0.79	0.11	72,72,72,72	0
7	NAG	B	501	14/15	0.80	0.32	47,58,62,65	0
7	NAG	C	501	14/15	0.81	0.42	56,68,75,76	0
7	NAG	D	501	14/15	0.84	0.28	46,58,65,66	0
7	NAG	A	501	14/15	0.86	0.30	49,58,63,64	0
8	CA	A	502	1/1	0.91	0.10	65,65,65,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around CA D 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around CA B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CA C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around NAG B 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around NAG C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAG D 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around NAG A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.