



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 07:36 am BST

PDB ID : 5M0J  
Title : Crystal structure of the cytoplasmic complex with She2p, She3p, and the ASH1 mRNA E3-localization element  
Authors : Edelman, F.T.; Janowski, R.; Niessing, D.  
Deposited on : 2016-10-05  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	 2% 64% 5% 29%
1	B	328	 2% 69% 5% 29%
1	C	328	 0% 70% 5% 27%
1	D	328	 2% 65% 5% 29%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	328	 95%
1	H	328	 93%
1	I	328	 94%
1	J	328	 94%
2	E	28	 71% 25%
2	F	28	 82% 18%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SWI5-dependent HO expression protein 2, SWI5-dependent HO expression protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	232	1914	1231	307	374	2	0	3	0
1	C	240	1952	1257	313	380	2	0	1	0
1	B	234	1920	1238	310	370	2	0	2	0
1	A	223	1811	1172	289	348	2	0	0	0
1	J	21	170	109	34	27		0	0	0
1	G	16	127	80	25	22		0	0	0
1	H	22	171	108	34	29		0	0	0
1	I	20	153	101	27	25		0	0	0

There are 128 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	4	GLY	-	expression tag	UNP B3LQW9
D	5	PRO	-	expression tag	UNP B3LQW9
D	14	SER	CYS	engineered mutation	UNP B3LQW9
D	68	SER	CYS	engineered mutation	UNP B3LQW9
D	106	SER	CYS	engineered mutation	UNP B3LQW9
D	180	SER	CYS	engineered mutation	UNP B3LQW9
D	247	GLY	-	linker	UNP B3LQW9
D	248	GLY	-	linker	UNP B3LQW9
D	249	SER	-	linker	UNP B3LQW9
D	250	GLY	-	linker	UNP B3LQW9
D	251	GLY	-	linker	UNP B3LQW9
D	252	GLY	-	linker	UNP B3LQW9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	253	GLY	-	linker	UNP B3LQW9
D	254	SER	-	linker	UNP B3LQW9
D	255	GLY	-	linker	UNP B3LQW9
D	256	GLY	-	linker	UNP B3LQW9
C	4	GLY	-	expression tag	UNP B3LQW9
C	5	PRO	-	expression tag	UNP B3LQW9
C	14	SER	CYS	engineered mutation	UNP B3LQW9
C	68	SER	CYS	engineered mutation	UNP B3LQW9
C	106	SER	CYS	engineered mutation	UNP B3LQW9
C	180	SER	CYS	engineered mutation	UNP B3LQW9
C	247	GLY	-	linker	UNP B3LQW9
C	248	GLY	-	linker	UNP B3LQW9
C	249	SER	-	linker	UNP B3LQW9
C	250	GLY	-	linker	UNP B3LQW9
C	251	GLY	-	linker	UNP B3LQW9
C	252	GLY	-	linker	UNP B3LQW9
C	253	GLY	-	linker	UNP B3LQW9
C	254	SER	-	linker	UNP B3LQW9
C	255	GLY	-	linker	UNP B3LQW9
C	256	GLY	-	linker	UNP B3LQW9
B	4	GLY	-	expression tag	UNP B3LQW9
B	5	PRO	-	expression tag	UNP B3LQW9
B	14	SER	CYS	engineered mutation	UNP B3LQW9
B	68	SER	CYS	engineered mutation	UNP B3LQW9
B	106	SER	CYS	engineered mutation	UNP B3LQW9
B	180	SER	CYS	engineered mutation	UNP B3LQW9
B	247	GLY	-	linker	UNP B3LQW9
B	248	GLY	-	linker	UNP B3LQW9
B	249	SER	-	linker	UNP B3LQW9
B	250	GLY	-	linker	UNP B3LQW9
B	251	GLY	-	linker	UNP B3LQW9
B	252	GLY	-	linker	UNP B3LQW9
B	253	GLY	-	linker	UNP B3LQW9
B	254	SER	-	linker	UNP B3LQW9
B	255	GLY	-	linker	UNP B3LQW9
B	256	GLY	-	linker	UNP B3LQW9
A	4	GLY	-	expression tag	UNP B3LQW9
A	5	PRO	-	expression tag	UNP B3LQW9
A	14	SER	CYS	engineered mutation	UNP B3LQW9
A	68	SER	CYS	engineered mutation	UNP B3LQW9
A	106	SER	CYS	engineered mutation	UNP B3LQW9
A	180	SER	CYS	engineered mutation	UNP B3LQW9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	GLY	-	linker	UNP B3LQW9
A	248	GLY	-	linker	UNP B3LQW9
A	249	SER	-	linker	UNP B3LQW9
A	250	GLY	-	linker	UNP B3LQW9
A	251	GLY	-	linker	UNP B3LQW9
A	252	GLY	-	linker	UNP B3LQW9
A	253	GLY	-	linker	UNP B3LQW9
A	254	SER	-	linker	UNP B3LQW9
A	255	GLY	-	linker	UNP B3LQW9
A	256	GLY	-	linker	UNP B3LQW9
J	78	GLY	-	expression tag	UNP B3LQW9
J	79	PRO	-	expression tag	UNP B3LQW9
J	88	SER	CYS	engineered mutation	UNP B3LQW9
J	142	SER	CYS	engineered mutation	UNP B3LQW9
J	180	SER	CYS	engineered mutation	UNP B3LQW9
J	254	SER	CYS	engineered mutation	UNP B3LQW9
J	321	GLY	-	linker	UNP B3LQW9
J	322	GLY	-	linker	UNP B3LQW9
J	323	SER	-	linker	UNP B3LQW9
J	324	GLY	-	linker	UNP B3LQW9
J	325	GLY	-	linker	UNP B3LQW9
J	326	GLY	-	linker	UNP B3LQW9
J	327	GLY	-	linker	UNP B3LQW9
J	328	SER	-	linker	UNP B3LQW9
J	329	GLY	-	linker	UNP B3LQW9
J	330	GLY	-	linker	UNP B3LQW9
G	78	GLY	-	expression tag	UNP B3LQW9
G	79	PRO	-	expression tag	UNP B3LQW9
G	88	SER	CYS	engineered mutation	UNP B3LQW9
G	142	SER	CYS	engineered mutation	UNP B3LQW9
G	180	SER	CYS	engineered mutation	UNP B3LQW9
G	254	SER	CYS	engineered mutation	UNP B3LQW9
G	321	GLY	-	linker	UNP B3LQW9
G	322	GLY	-	linker	UNP B3LQW9
G	323	SER	-	linker	UNP B3LQW9
G	324	GLY	-	linker	UNP B3LQW9
G	325	GLY	-	linker	UNP B3LQW9
G	326	GLY	-	linker	UNP B3LQW9
G	327	GLY	-	linker	UNP B3LQW9
G	328	SER	-	linker	UNP B3LQW9
G	329	GLY	-	linker	UNP B3LQW9
G	330	GLY	-	linker	UNP B3LQW9

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	78	GLY	-	expression tag	UNP B3LQW9
H	79	PRO	-	expression tag	UNP B3LQW9
H	88	SER	CYS	engineered mutation	UNP B3LQW9
H	142	SER	CYS	engineered mutation	UNP B3LQW9
H	180	SER	CYS	engineered mutation	UNP B3LQW9
H	254	SER	CYS	engineered mutation	UNP B3LQW9
H	321	GLY	-	linker	UNP B3LQW9
H	322	GLY	-	linker	UNP B3LQW9
H	323	SER	-	linker	UNP B3LQW9
H	324	GLY	-	linker	UNP B3LQW9
H	325	GLY	-	linker	UNP B3LQW9
H	326	GLY	-	linker	UNP B3LQW9
H	327	GLY	-	linker	UNP B3LQW9
H	328	SER	-	linker	UNP B3LQW9
H	329	GLY	-	linker	UNP B3LQW9
H	330	GLY	-	linker	UNP B3LQW9
I	78	GLY	-	expression tag	UNP B3LQW9
I	79	PRO	-	expression tag	UNP B3LQW9
I	88	SER	CYS	engineered mutation	UNP B3LQW9
I	142	SER	CYS	engineered mutation	UNP B3LQW9
I	180	SER	CYS	engineered mutation	UNP B3LQW9
I	254	SER	CYS	engineered mutation	UNP B3LQW9
I	321	GLY	-	linker	UNP B3LQW9
I	322	GLY	-	linker	UNP B3LQW9
I	323	SER	-	linker	UNP B3LQW9
I	324	GLY	-	linker	UNP B3LQW9
I	325	GLY	-	linker	UNP B3LQW9
I	326	GLY	-	linker	UNP B3LQW9
I	327	GLY	-	linker	UNP B3LQW9
I	328	SER	-	linker	UNP B3LQW9
I	329	GLY	-	linker	UNP B3LQW9
I	330	GLY	-	linker	UNP B3LQW9

- Molecule 2 is a RNA chain called ASH1 E3 (28 nt-loop).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	28	Total	C	N	O	P	0	0	0
			596	269	112	188	27			
2	F	28	Total	C	N	O	P	0	0	0
			596	269	112	188	27			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	39	Total O 39 39	0	0
4	C	37	Total O 37 37	0	0
4	B	45	Total O 45 45	0	0
4	A	33	Total O 33 33	0	0
4	J	3	Total O 3 3	0	0
4	G	1	Total O 1 1	0	0
4	H	5	Total O 5 5	0	0
4	I	5	Total O 5 5	0	0
4	E	19	Total O 19 19	0	0
4	F	20	Total O 20 20	0	0











## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	219.03Å 58.98Å 144.68Å 90.00° 126.91° 90.00°	Depositor
Resolution (Å)	50.01 – 2.80 47.29 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.01-2.80) 100.0 (47.29-2.80)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.194 , 0.246 0.198 , 0.246	Depositor DCC
$R_{free}$ test set	1828 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.2	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9621	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1844	0.74	2/2494 (0.1%)
1	B	0.58	0/1956	0.76	0/2645
1	C	0.57	0/1989	0.72	0/2693
1	D	0.58	0/1950	0.78	1/2640 (0.0%)
1	G	0.68	0/129	0.88	0/172
1	H	0.72	0/174	0.90	0/233
1	I	0.67	0/156	0.83	0/210
1	J	0.69	0/172	0.99	0/229
2	E	0.56	0/668	0.91	4/1039 (0.4%)
2	F	0.51	0/668	0.81	0/1039
All	All	0.57	0/9706	0.78	7/13394 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	26	ARG	NE-CZ-NH1	-7.93	116.33	120.30
1	A	26	ARG	NE-CZ-NH2	7.13	123.86	120.30
2	E	7	U	N1-C1'-C2'	6.87	122.93	114.00
2	E	26	A	N9-C1'-C2'	6.38	122.29	114.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	26	A	O4'-C1'-N9	6.22	113.18	108.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	184	ASP	Peptide
1	D	82	LYS	Peptide
1	D	85	LEU	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1811	0	1811	4	0
1	B	1920	0	1916	2	0
1	C	1952	0	1945	3	0
1	D	1914	0	1892	8	0
1	G	127	0	126	0	0
1	H	171	0	179	0	0
1	I	153	0	165	1	0
1	J	170	0	183	2	0
2	E	596	0	304	1	0
2	F	596	0	304	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	33	0	0	0	0
4	B	45	0	0	1	0
4	C	37	0	0	0	0
4	D	39	0	0	1	0
4	E	19	0	0	0	0
4	F	20	0	0	0	0
4	G	1	0	0	0	0
4	H	5	0	0	0	0
4	I	5	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	J	3	0	0	0	0
All	All	9621	0	8825	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 16 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:197:GLN:HG3	1:J:367:VAL:HG13	1.78	0.65
1:C:191:THR:HG21	1:C:196:LEU:HB2	1.83	0.60
1:D:191:THR:HG21	1:D:196:LEU:HB2	1.85	0.58
1:D:47:THR:HG21	1:A:47:THR:HG21	1.89	0.53
1:D:197:GLN:HG3	1:J:367:VAL:CG1	2.44	0.47

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/328 (66%)	213 (98%)	4 (2%)	0	100	100
1	B	232/328 (71%)	226 (97%)	5 (2%)	1 (0%)	34	66
1	C	239/328 (73%)	235 (98%)	3 (1%)	1 (0%)	34	66
1	D	231/328 (70%)	223 (96%)	6 (3%)	2 (1%)	17	46
1	G	12/328 (4%)	12 (100%)	0	0	100	100
1	H	18/328 (6%)	16 (89%)	2 (11%)	0	100	100
1	I	16/328 (5%)	16 (100%)	0	0	100	100
1	J	17/328 (5%)	14 (82%)	2 (12%)	1 (6%)	1	4
All	All	982/2624 (37%)	955 (97%)	22 (2%)	5 (0%)	29	61



All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	83	ASN
1	C	84	GLU
1	J	350	LEU
1	B	84	GLU
1	D	7	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/292 (70%)	199 (97%)	6 (3%)	42	76
1	B	217/292 (74%)	212 (98%)	5 (2%)	50	82
1	C	221/292 (76%)	217 (98%)	4 (2%)	59	86
1	D	217/292 (74%)	210 (97%)	7 (3%)	39	73
1	G	14/292 (5%)	12 (86%)	2 (14%)	3	10
1	H	19/292 (6%)	17 (90%)	2 (10%)	7	20
1	I	17/292 (6%)	16 (94%)	1 (6%)	19	49
1	J	19/292 (6%)	18 (95%)	1 (5%)	22	54
All	All	929/2336 (40%)	901 (97%)	28 (3%)	41	75

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	80	GLU
1	B	182	ASP
1	H	363	VAL
1	B	84	GLU
1	B	142	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	66	ASN
1	B	66	ASN
1	A	41	HIS

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	27/28 (96%)	4 (14%)	0
2	F	27/28 (96%)	5 (18%)	0
All	All	54/56 (96%)	9 (16%)	0

5 of 9 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	7	U
2	E	20	A
2	E	27	C
2	E	28	G
2	F	7	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	223/328 (67%)	-0.07	8 (3%) 42 32	23, 50, 98, 118	1 (0%)
1	B	234/328 (71%)	-0.28	6 (2%) 56 46	20, 40, 94, 121	1 (0%)
1	C	240/328 (73%)	-0.22	4 (1%) 70 63	23, 39, 90, 134	0
1	D	232/328 (70%)	-0.22	8 (3%) 45 35	20, 41, 95, 141	0
1	G	16/328 (4%)	0.38	2 (12%) 3 2	33, 80, 98, 106	0
1	H	22/328 (6%)	0.37	3 (13%) 3 1	45, 68, 109, 126	0
1	I	20/328 (6%)	0.61	1 (5%) 28 19	35, 66, 94, 103	0
1	J	21/328 (6%)	0.33	2 (9%) 8 4	33, 62, 108, 114	0
2	E	28/28 (100%)	0.02	0 100 100	31, 45, 69, 77	0
2	F	28/28 (100%)	0.11	2 (7%) 16 9	28, 42, 78, 82	0
All	All	1064/2680 (39%)	-0.14	36 (3%) 45 35	20, 43, 98, 141	2 (0%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	83	ASN	5.6
1	A	11	PRO	4.9
1	C	83	ASN	4.8
1	C	81	GLY	4.0
1	I	350	LEU	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	D	401	1/1	0.95	0.53	62,62,62,62	0
3	MG	A	401	1/1	0.95	0.52	70,70,70,70	0
3	MG	B	401	1/1	0.98	0.45	50,50,50,50	0
3	MG	C	401	1/1	0.98	0.46	54,54,54,54	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.