

wwPDB X-ray Structure Validation Summary Report (i)

Nov 22, 2023 – 03:43 PM JST

PDB ID : 6M2P

Title: Crystal structure of a NIR-emitting DNA-stabilized Ag16 nanocluster (A10-

deletion mutant)

Authors: Kondo, J.; Cerretani, C.; Vosch, T.

Deposited on : 2020-02-28

Resolution : 1.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $\begin{array}{ccc} & Mol Probity & : & 4.02b\text{-}467 \\ Xtriage \left(Phenix\right) & : & 1.13 \end{array}$

EDS: 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

 $CCP4 : 7.0.044 ext{ (Gargrove)}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1168 (1.14-1.10)
Clashscore	141614	1205 (1.14-1.10)
RSRZ outliers	127900	1146 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	9	89%	11%
1	В	9	89%	11%
1	С	9	89%	11%
1	D	9	89%	11%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 1442 atoms, of which 474 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a DNA chain called DNA (5'-D(*CP*AP*CP*CP*TP*AP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Λ	0	Total C	Н	N	О	Р	0	$0 \qquad 2$	0	
1	Λ	9	316 96	115	34	61	10	U	2		
1	В	0	Total C	Н	N	О	Р	0	3	0	
1	Б	9	334 101	122	34	66	11	U	3	0	
1	С	9	Total C	Н	N	О	Р	0	9	0	
1		9	316 96	115	34	61	10	U	2	U	
1	D	9	Total C	Н	N	О	Р	0	2	0	
1	ש	9	334 101	122	34	66	11	U	3		

• Molecule 2 is SILVER ION (three-letter code: AG) (formula: Ag) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	8	Total Ag 8 8	0	0
2	В	8	Total Ag 8 8	0	0
2	С	8	Total Ag 8 8	0	0
2	D	8	Total Ag 8 8	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	D	1	Total Ca 2 2	0	1

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	В	1	Total Cl 1 1	0	0
4	C	1	Total Cl 1 1	0	0
4	D	1	Total Cl 1 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	29	Total O 29 29	0	0
5	В	35	Total O 35 35	0	0
5	С	14	Total O 14 14	0	0
5	D	25	Total O 25 25	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA (5'-D(*CP*AP*CP*CP*TP*AP*GP*CP*G)-3')





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	24.21Å 33.71Å 58.55Å	Donositor
a, b, c, α , β , γ	90.00° 98.11° 90.00°	Depositor
Resolution (Å)	29.14 - 1.13	Depositor
Resolution (A)	29.14 - 1.13	EDS
% Data completeness	96.8 (29.14-1.13)	Depositor
(in resolution range)	97.0 (29.14-1.13)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.74 (at 1.12Å)	Xtriage
Refinement program	PHENIX 1.17.1	Depositor
D D.	0.085 , 0.096	Depositor
R, R_{free}	0.082 , 0.091	DCC
R_{free} test set	2019 reflections (5.84%)	wwPDB-VP
Wilson B-factor (Å ²)	4.3	Xtriage
Anisotropy	0.387	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.44,62.3	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	1442	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 29.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4102e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AG, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ # Z > 5		RMSZ	# Z > 5
1	A	0.96	0/243	1.27	1/372~(0.3%)
1	В	1.02	0/264	1.25	1/404 (0.2%)
1	С	0.85	0/243	1.25	1/372 (0.3%)
1	D	0.94	0/264	1.19	1/404 (0.2%)
All	All	0.95	0/1014	1.24	4/1552 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers	
1	A	0	1	

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	A	7	DG	O4'-C1'-N9	8.72	114.11	108.00
1	С	7	DG	O4'-C1'-N9	8.68	114.08	108.00
1	D	7	DG	O4'-C1'-N9	8.00	113.60	108.00
1	В	7	DG	O4'-C1'-N9	7.49	113.24	108.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mo	1	Chain	Res	Type	Group
1		A	7	DG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	201	115	106	0	0
1	В	212	122	109	0	0
1	С	201	115	106	0	0
1	D	212	122	109	0	0
2	A	8	0	0	0	0
2	В	8	0	0	0	0
2	С	8	0	0	0	0
2	D	8	0	0	0	0
3	A	1	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
5	A	29	0	0	0	0
5	В	35	0	0	0	0
5	С	14	0	0	0	0
5	D	25	0	0	0	0
All	All	968	474	430	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 39 ligands modelled in this entry, 39 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		ZZ>2	$OWAB(A^2)$	Q < 0.9
1	A	9/9~(100%)	0.17	0	100	100	7, 8, 9, 10	0
1	В	9/9~(100%)	0.16	0	100	100	8, 9, 11, 11	0
1	С	9/9~(100%)	0.31	0	100	100	9, 11, 13, 16	0
1	D	9/9~(100%)	0.26	0	100	100	9, 10, 13, 14	0
All	All	$36/36 \ (100\%)$	0.22	0	100	100	7, 9, 13, 16	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q < 0.9
3	CA	D	109[A]	1/1	0.99	0.06	11,11,11,11	1
3	CA	D	109[B]	1/1	0.99	0.06	12,12,12,12	1
2	AG	A	103	1/1	1.00	0.06	6,6,6,6	0
2	AG	A	104	1/1	1.00	0.05	6,6,6,6	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q < 0.9
2	AG	A	105	1/1	1.00	0.05	6,6,6,6	0
2	AG	A	106	1/1	1.00	0.06	6,6,6,6	0
2	AG	A	107	1/1	1.00	0.06	6,6,6,6	0
2	AG	A	108	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	101	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	102	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	103	1/1	1.00	0.06	6,6,6,6	0
2	AG	В	104	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	105	1/1	1.00	0.06	6,6,6,6	0
2	AG	В	106	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	107	1/1	1.00	0.05	6,6,6,6	0
2	AG	В	108	1/1	1.00	0.05	6,6,6,6	0
2	AG	С	101	1/1	1.00	0.05	8,8,8,8	0
2	AG	С	102	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	103	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	104	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	105	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	106	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	107	1/1	1.00	0.05	7,7,7,7	0
2	AG	С	108	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	101	1/1	1.00	0.05	8,8,8,8	0
2	AG	D	102	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	103	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	104	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	105	1/1	1.00	0.05	8,8,8,8	0
2	AG	D	106	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	107	1/1	1.00	0.05	7,7,7,7	0
2	AG	D	108	1/1	1.00	0.05	7,7,7,7	0
3	CA	A	109	1/1	1.00	0.05	8,8,8,8	0
2	AG	A	101	1/1	1.00	0.05	6,6,6,6	0
2	AG	A	102	1/1	1.00	0.06	6,6,6,6	0
4	CL	A	110	1/1	1.00	0.06	6,6,6,6	0
4	CL	В	109	1/1	1.00	0.06	6,6,6,6	0
4	CL	С	109	1/1	1.00	0.06	7,7,7,7	0
4	CL	D	110	1/1	1.00	0.05	7,7,7,7	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around AG A 103:













Electron density around AG A 104:









Electron density around AG A 105:













Electron density around AG A 106:









Electron density around AG A 107:












Electron density around AG A 108:












Electron density around AG B 101:












Electron density around AG B 102:












Electron density around AG B 103:













Electron density around AG B 104:









Electron density around AG B 105:












Electron density around AG B 106:












Electron density around AG B 107:












Electron density around AG B 108:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around AG C 101:









Electron density around AG C 102:












Electron density around AG C 103:












Electron density around AG C 104:












Electron density around AG C 105:












Electron density around AG C 106:












Electron density around AG C 107:












Electron density around AG C 108:












Electron density around AG D 101:












Electron density around AG D 102:













Electron density around AG D 103:









Electron density around AG D 104:












Electron density around AG D 105:












Electron density around AG D 106:












Electron density around AG D 107:












Electron density around AG D 108:












Electron density around AG A 101:













Electron density around AG A 102:










Electron density around CL A 110:









Electron density around CL B 109:













Electron density around CL C 109:









6.5 Other polymers (i)

There are no such residues in this entry.

