



# wwPDB X-ray Structure Validation Summary Report

Sep 22, 2022 – 02:12 PM EDT

PDB ID : 7M66  
Title : Targeting Enterococcus faecalis HMG-CoA reductase with a novel non-statin inhibitor  
Authors : Bose, S.; Steussy, C.N.  
Deposited on : 2021-03-25  
Resolution : 2.25 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	
1	B	423	
1	C	423	
1	D	423	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 23361 atoms, of which 11422 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydroxymethylglutaryl-CoA reductase, degradative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	360	5560	1741	2790	473	543	13	0	4	0
1	B	360	5651	1766	2846	482	545	12	0	12	0
1	C	360	5580	1745	2801	477	545	12	0	8	0
1	D	357	5635	1761	2836	478	546	14	0	17	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	C	1	5	4	1	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
3	A	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		
3	B	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		
3	B	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		
3	D	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		
3	D	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	17	4	10	3	0	0
4	A	1	17	4	10	3	0	0
4	A	1	17	4	10	3	0	0
4	C	1	17	4	10	3	0	0
4	C	1	17	4	10	3	0	0
4	D	1	17	4	10	3	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
5	A	3	3	3	0	0
5	B	2	2	2	0	0
5	C	2	2	2	0	0
5	D	3	3	3	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
6	B	1	Total	C	H	O	0	0
			10	2	6	2		
6	B	1	Total	C	H	O	0	0
			10	2	6	2		
6	B	1	Total	C	H	O	0	0
			10	2	6	2		
6	C	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	153	Total	O	0	0
			153	153		
7	B	156	Total	O	0	0
			156	156		
7	C	158	Total	O	0	0
			158	158		
7	D	179	Total	O	0	2
			181	181		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydroxymethylglutaryl-CoA reductase, degradative



- Molecule 1: Hydroxymethylglutaryl-CoA reductase, degradative



- Molecule 1: Hydroxymethylglutaryl-CoA reductase, degradative



THR  
GLY  
LYS  
GLU  
VAL  
GLU  
ALA  
VAL  
ALA  
GLN  
GLN  
LEU  
LYS  
ARG  
GLN  
LYS  
THR  
MET  
ASN  
GLN  
ASP  
ARG  
ALA  
LEU  
ALA  
ILE  
LEU  
ASN  
ASP  
LEU  
ARG  
LYS  
GLN

- Molecule 1: Hydroxymethylglutaryl-CoA reductase, degradative

Chain D: 6% 82% 16%

MET  
LEU  
LEU  
GLU  
ARG  
PRO  
GLN  
GLN  
LYS  
LYS  
ASN  
SER  
R13  
M112  
R113  
L211  
P227  
V228  
S229  
R230  
L231  
S232  
K233  
G234  
S235  
R238  
E239  
I240  
K243  
Y250  
K263  
M266  
N267  
G268  
I269  
E270  
A271  
V272  
V273  
R281  
K282  
V283  
S284  
H288  
D306  
G307  
E308  
L340

A341  
V342  
T343  
E347  
S369  
GLU  
GLY  
ILE  
GLN  
LYS  
GLY  
HIS  
MET  
ALA  
LEU  
GLN  
ALA  
ARG  
SER  
LEU  
ALA  
ALA  
MET  
THR  
VAL  
GLY  
ALA  
THR  
GLY  
LYS  
GLU  
VAL  
GLU  
ALA  
VAL  
ALA  
GLN  
GLN  
LEU  
LYS  
ARG  
GLN  
LYS  
THR  
MET  
ASN  
GLN  
ASP  
ARG  
ALA  
LEU  
ALA  
ILE  
LEU  
ASN  
ASP  
LEU  
ARG  
LYS

GLN



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	168.66Å 168.66Å 120.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.13 – 2.25 45.13 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.13-2.25) 99.8 (45.13-2.25)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.39 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.175 , 0.209 0.173 , 0.206	Depositor DCC
$R_{free}$ test set	4593 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.7	Xtrriage
Anisotropy	0.005	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 42.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	23361	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.5905e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MES, PEG, SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2808	0.48	0/3801
1	B	0.27	0/2875	0.51	0/3892
1	C	0.26	0/2851	0.49	0/3858
1	D	0.26	0/2886	0.50	0/3907
All	All	0.26	0/11420	0.50	0/15458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2770	2790	2786	8	0
1	B	2805	2846	2799	6	0
1	C	2779	2801	2765	3	0
1	D	2799	2836	2782	6	0
2	A	5	0	0	0	0
2	C	5	0	0	0	0
3	A	12	13	13	0	0
3	B	24	26	25	0	0
3	D	24	26	26	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	21	30	30	0	0
4	C	14	20	20	0	0
4	D	7	10	10	0	0
5	A	3	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	3	0	0	0	0
6	B	12	18	18	0	0
6	C	4	6	6	0	0
7	A	153	0	0	2	0
7	B	156	0	0	1	0
7	C	158	0	0	1	0
7	D	181	0	0	0	0
All	All	11939	11422	11280	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:ARG:O	1:B:211[A]:LEU:HD22	2.06	0.55
1:D:112[B]:MET:SD	1:D:211[B]:LEU:HD11	2.46	0.55
1:D:113:ARG:O	1:D:211[A]:LEU:HD22	2.07	0.55
1:B:183:ALA:HA	1:B:211[B]:LEU:HD22	1.91	0.54
1:C:342:VAL:HG22	7:C:658:HOH:O	2.10	0.52

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/423 (86%)	354 (98%)	8 (2%)	0	100	100
1	B	370/423 (88%)	360 (97%)	10 (3%)	0	100	100
1	C	366/423 (86%)	358 (98%)	8 (2%)	0	100	100
1	D	372/423 (88%)	363 (98%)	9 (2%)	0	100	100
All	All	1470/1692 (87%)	1435 (98%)	35 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/344 (86%)	294 (99%)	3 (1%)	76	84
1	B	304/344 (88%)	298 (98%)	6 (2%)	55	64
1	C	301/344 (88%)	294 (98%)	7 (2%)	50	59
1	D	306/344 (89%)	303 (99%)	3 (1%)	76	84
All	All	1208/1376 (88%)	1189 (98%)	19 (2%)	65	73

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	281	ARG
1	D	235	SER
1	D	281	ARG
1	D	229	SER
1	B	281	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 10 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	504	-	6,6,6	0.66	0	5,5,5	1.08	0
4	PEG	C	503	-	6,6,6	0.66	0	5,5,5	1.08	0
2	SO4	A	501	-	4,4,4	0.14	0	6,6,6	0.06	0
6	EDO	C	502	-	3,3,3	0.48	0	2,2,2	0.30	0
6	EDO	B	504	-	3,3,3	0.47	0	2,2,2	0.31	0
3	MES	A	502	-	12,12,12	2.22	1 (8%)	14,16,16	1.60	4 (28%)
2	SO4	C	501	-	4,4,4	0.14	0	6,6,6	0.06	0
4	PEG	A	505	-	6,6,6	0.61	0	5,5,5	1.12	0
4	PEG	A	503	-	6,6,6	0.63	0	5,5,5	1.03	0
4	PEG	D	503	-	6,6,6	0.60	0	5,5,5	1.08	0
6	EDO	B	505	-	3,3,3	0.47	0	2,2,2	0.33	0
3	MES	B	502	-	12,12,12	2.17	1 (8%)	14,16,16	1.54	3 (21%)
3	MES	B	501	-	12,12,12	2.22	1 (8%)	14,16,16	1.56	3 (21%)
3	MES	D	502	-	12,12,12	2.18	1 (8%)	14,16,16	1.80	5 (35%)
4	PEG	C	504	-	6,6,6	0.64	0	5,5,5	1.07	0
3	MES	D	501	-	12,12,12	2.07	1 (8%)	14,16,16	2.01	6 (42%)
6	EDO	B	503	-	3,3,3	0.46	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	504	-	-	3/4/4/4	-
4	PEG	C	503	-	-	1/4/4/4	-
6	EDO	C	502	-	-	1/1/1/1	-
6	EDO	B	504	-	-	1/1/1/1	-
3	MES	A	502	-	-	2/6/14/14	0/1/1/1
4	PEG	A	505	-	-	2/4/4/4	-
4	PEG	A	503	-	-	1/4/4/4	-
4	PEG	D	503	-	-	1/4/4/4	-
6	EDO	B	505	-	-	1/1/1/1	-
3	MES	B	502	-	-	5/6/14/14	0/1/1/1
3	MES	B	501	-	-	0/6/14/14	0/1/1/1
3	MES	D	502	-	-	1/6/14/14	0/1/1/1
4	PEG	C	504	-	-	3/4/4/4	-
3	MES	D	501	-	-	1/6/14/14	0/1/1/1
6	EDO	B	503	-	-	0/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	MES	C8-S	-7.43	1.66	1.77
3	A	502	MES	C8-S	-7.42	1.67	1.77
3	D	502	MES	C8-S	-7.26	1.67	1.77
3	B	502	MES	C8-S	-7.24	1.67	1.77
3	D	501	MES	C8-S	-6.87	1.67	1.77

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	501	MES	O2S-S-C8	4.16	111.92	106.92
3	D	502	MES	O3S-S-C8	3.88	112.04	105.77
3	B	501	MES	C5-N4-C3	3.55	116.81	108.83
3	B	502	MES	O3S-S-C8	3.31	111.13	105.77
3	D	502	MES	C2-C3-N4	-2.73	105.97	110.10

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	MES	C7-C8-S-O1S
3	B	502	MES	C7-C8-S-O2S

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	502	MES	C7-C8-S-O3S
4	D	503	PEG	O2-C3-C4-O4
4	A	503	PEG	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/423 (85%)	0.68	62 (17%) <b>1</b> <b>1</b>	27, 42, 85, 97	10 (2%)
1	B	360/423 (85%)	0.18	30 (8%) <b>11</b> <b>12</b>	29, 41, 75, 107	9 (2%)
1	C	360/423 (85%)	0.64	63 (17%) <b>1</b> <b>1</b>	26, 41, 87, 100	14 (3%)
1	D	357/423 (84%)	0.12	25 (7%) <b>16</b> <b>17</b>	28, 39, 72, 88	7 (1%)
All	All	1437/1692 (84%)	0.41	180 (12%) <b>3</b> <b>3</b>	26, 41, 82, 107	40 (2%)

The worst 5 of 180 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	PHE	7.2
1	C	166	GLU	6.6
1	A	166	GLU	5.8
1	A	11	ASN	5.6
1	C	133	VAL	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	C	502	4/4	0.63	0.18	59,71,79,82	0
6	EDO	B	505	4/4	0.65	0.30	54,65,70,76	10
4	PEG	C	503	7/7	0.70	0.20	53,65,84,84	0
6	EDO	B	503	4/4	0.75	0.23	63,76,81,82	0
3	MES	A	502	12/12	0.78	0.27	47,56,68,70	25
4	PEG	A	505	7/7	0.78	0.25	35,54,63,65	0
4	PEG	A	503	7/7	0.79	0.15	49,63,75,76	0
4	PEG	A	504	7/7	0.80	0.29	56,69,84,85	0
6	EDO	B	504	4/4	0.82	0.27	54,65,67,76	0
3	MES	B	502	12/12	0.84	0.28	53,70,87,89	25
4	PEG	C	504	7/7	0.85	0.45	59,73,80,86	0
2	SO4	C	501	5/5	0.87	0.23	52,55,64,66	5
4	PEG	D	503	7/7	0.88	0.21	39,48,57,65	17
2	SO4	A	501	5/5	0.92	0.16	68,73,80,90	0
3	MES	D	502	12/12	0.94	0.14	48,70,93,108	0
5	CA	D	505	1/1	0.94	0.07	35,35,35,35	0
3	MES	D	501	12/12	0.94	0.11	36,45,53,57	0
5	CA	C	506	1/1	0.95	0.03	64,64,64,64	0
5	CA	D	504	1/1	0.97	0.05	51,51,51,51	0
5	CA	A	507	1/1	0.97	0.12	42,42,42,42	0
5	CA	A	508	1/1	0.97	0.03	68,68,68,68	0
5	CA	B	506	1/1	0.97	0.05	51,51,51,51	0
5	CA	B	507	1/1	0.97	0.11	74,74,74,74	0
3	MES	B	501	12/12	0.97	0.18	39,54,63,64	25
5	CA	A	506	1/1	0.98	0.21	34,34,34,34	0
5	CA	D	506	1/1	0.98	0.21	71,71,71,71	0
5	CA	C	505	1/1	0.98	0.09	43,43,43,43	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.