



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 12:25 pm BST

PDB ID : 3M9C
Title : Crystal structure of the membrane domain of respiratory complex I from Escherichia coli
Authors : Efremov, R.G.; Baradaran, R.; Sazanov, L.A.
Deposited on : 2010-03-22
Resolution : 3.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-quinone oxidoreductase subunit NuoL.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
1	L	474	Total	C	N	0	0	0
			1422	948	474			

- Molecule 2 is a protein called NADH-quinone oxidoreductase subunit NuoM.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	M	391	Total	C	N	0	0	0
			1173	782	391			

- Molecule 3 is a protein called NADH-quinone oxidoreductase subunit NuoN.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	N	378	Total	C	N	0	0	0
			1134	756	378			


- Molecule 4 is a protein called NADH-quinone oxidoreductase subunits NuoA,J and K.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	R	281	Total	C	N	0	0	0
			843	562	281			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH-quinone oxidoreductase subunit NuoL




- Molecule 2: NADH-quinone oxidoreductase subunit NuoM



- Molecule 3: NADH-quinone oxidoreductase subunit NuoN



- Molecule 4: NADH-quinone oxidoreductase subunits NuoA,J and K



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	114.06Å 182.17Å 185.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.90 29.96 – 3.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-3.90) 97.5 (29.96-3.90)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 3.86Å)	Xtrriage
Refinement program		Depositor
R, R_{free}	(Not available) , (Not available) 0.457 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	108.5	Xtrriage
Anisotropy	0.720	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.18$	Xtrriage
Estimated twinning fraction	0.129 for -h,l,k	Xtrriage
F_o, F_c correlation	0.65	EDS
Total number of atoms	4572	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	1212	UNK	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1422	0	52	10	0
2	M	1173	0	37	13	0
3	N	1134	0	36	10	0
4	R	843	0	24	3	0
All	All	4572	0	149	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:N:1001:UNK:CA	3:N:1059:UNK:C	1.85	1.50
2:M:920:UNK:CA	2:M:1200:UNK:N	2.03	1.21
3:N:310:UNK:CA	3:N:350:UNK:CA	2.31	1.08
3:N:1001:UNK:C	3:N:1059:UNK:C	2.40	0.98
2:M:920:UNK:C	2:M:1200:UNK:H2	1.78	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	L	20
3	N	16
2	M	16
4	R	10

The worst 5 of 62 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	L	428:UNK	C	495:UNK	N	56.90
1	M	1024:UNK	C	1050:UNK	N	55.04
1	N	366:UNK	C	395:UNK	N	50.88
1	L	1417:UNK	C	1501:UNK	N	47.71
1	N	665:UNK	C	700:UNK	N	47.65

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.