



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 11:23 AM EDT

PDB ID : 4MAR  
Title : Crystal structure of purine nucleoside phosphorylase from *Meiothermus ruber* DSM 1279 complexed with sulfate.  
Authors : Malashkevich, V.N.; Bhosle, R.; Toro, R.; Hillerich, B.; Gizzi, A.; Garforth, S.; Kar, A.; Chan, M.K.; Laffuer, J.; Patel, H.; Matikainen, B.; Chamala, S.; Lim, S.; Celikgil, A.; Villegas, G.; Evans, B.; Love, J.; Fiser, A.; Khafizov, K.; Seidel, R.; Bonanno, J.B.; Almo, S.C.; New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2013-08-16  
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div>
1	B	254	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div>
1	C	254	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div>

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.35.1

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purine nucleoside phosphorylase DeoD-type.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	231	1808	1156	308	336	3	5	0	4	0
1	B	231	1800	1147	308	337	3	5	0	2	0
1	C	231	1811	1155	307	341	3	5	0	5	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MSE	-	expression tag	UNP D3PPS9
A	-20	HIS	-	expression tag	UNP D3PPS9
A	-19	HIS	-	expression tag	UNP D3PPS9
A	-18	HIS	-	expression tag	UNP D3PPS9
A	-17	HIS	-	expression tag	UNP D3PPS9
A	-16	HIS	-	expression tag	UNP D3PPS9
A	-15	HIS	-	expression tag	UNP D3PPS9
A	-14	SER	-	expression tag	UNP D3PPS9
A	-13	SER	-	expression tag	UNP D3PPS9
A	-12	GLY	-	expression tag	UNP D3PPS9
A	-11	VAL	-	expression tag	UNP D3PPS9
A	-10	ASP	-	expression tag	UNP D3PPS9
A	-9	LEU	-	expression tag	UNP D3PPS9
A	-8	GLY	-	expression tag	UNP D3PPS9
A	-7	THR	-	expression tag	UNP D3PPS9
A	-6	GLU	-	expression tag	UNP D3PPS9
A	-5	ASN	-	expression tag	UNP D3PPS9
A	-4	LEU	-	expression tag	UNP D3PPS9
A	-3	TYR	-	expression tag	UNP D3PPS9
A	-2	PHE	-	expression tag	UNP D3PPS9
A	-1	GLN	-	expression tag	UNP D3PPS9
A	0	SER	-	expression tag	UNP D3PPS9
B	-21	MSE	-	expression tag	UNP D3PPS9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	HIS	-	expression tag	UNP D3PPS9
B	-19	HIS	-	expression tag	UNP D3PPS9
B	-18	HIS	-	expression tag	UNP D3PPS9
B	-17	HIS	-	expression tag	UNP D3PPS9
B	-16	HIS	-	expression tag	UNP D3PPS9
B	-15	HIS	-	expression tag	UNP D3PPS9
B	-14	SER	-	expression tag	UNP D3PPS9
B	-13	SER	-	expression tag	UNP D3PPS9
B	-12	GLY	-	expression tag	UNP D3PPS9
B	-11	VAL	-	expression tag	UNP D3PPS9
B	-10	ASP	-	expression tag	UNP D3PPS9
B	-9	LEU	-	expression tag	UNP D3PPS9
B	-8	GLY	-	expression tag	UNP D3PPS9
B	-7	THR	-	expression tag	UNP D3PPS9
B	-6	GLU	-	expression tag	UNP D3PPS9
B	-5	ASN	-	expression tag	UNP D3PPS9
B	-4	LEU	-	expression tag	UNP D3PPS9
B	-3	TYR	-	expression tag	UNP D3PPS9
B	-2	PHE	-	expression tag	UNP D3PPS9
B	-1	GLN	-	expression tag	UNP D3PPS9
B	0	SER	-	expression tag	UNP D3PPS9
C	-21	MSE	-	expression tag	UNP D3PPS9
C	-20	HIS	-	expression tag	UNP D3PPS9
C	-19	HIS	-	expression tag	UNP D3PPS9
C	-18	HIS	-	expression tag	UNP D3PPS9
C	-17	HIS	-	expression tag	UNP D3PPS9
C	-16	HIS	-	expression tag	UNP D3PPS9
C	-15	HIS	-	expression tag	UNP D3PPS9
C	-14	SER	-	expression tag	UNP D3PPS9
C	-13	SER	-	expression tag	UNP D3PPS9
C	-12	GLY	-	expression tag	UNP D3PPS9
C	-11	VAL	-	expression tag	UNP D3PPS9
C	-10	ASP	-	expression tag	UNP D3PPS9
C	-9	LEU	-	expression tag	UNP D3PPS9
C	-8	GLY	-	expression tag	UNP D3PPS9
C	-7	THR	-	expression tag	UNP D3PPS9
C	-6	GLU	-	expression tag	UNP D3PPS9
C	-5	ASN	-	expression tag	UNP D3PPS9
C	-4	LEU	-	expression tag	UNP D3PPS9
C	-3	TYR	-	expression tag	UNP D3PPS9
C	-2	PHE	-	expression tag	UNP D3PPS9
C	-1	GLN	-	expression tag	UNP D3PPS9

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP D3PPS9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	105	Total O 105 105	0	0
4	B	71	Total O 71 71	0	0
4	C	107	Total O 107 107	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Purine nucleoside phosphorylase DeoD-type



- Molecule 1: Purine nucleoside phosphorylase DeoD-type



- Molecule 1: Purine nucleoside phosphorylase DeoD-type



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.40Å 189.10Å 152.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.56 – 2.16 32.56 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.0 (32.56-2.16) 99.0 (32.56-2.16)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.28 (at 2.16Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.188 , 0.226 0.188 , 0.225	Depositor DCC
$R_{free}$ test set	2242 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtrriage
Anisotropy	1.237	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5758	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1853	0.59	0/2512
1	B	0.36	0/1839	0.56	0/2492
1	C	0.41	0/1854	0.60	0/2513
All	All	0.39	0/5546	0.58	0/7517

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1821	14	0
1	B	1800	0	1798	17	0
1	C	1811	0	1808	16	0
2	A	1	0	0	0	0
3	A	15	0	0	1	0
3	B	20	0	0	0	0
3	C	20	0	0	0	0
4	A	105	0	0	3	0
4	B	71	0	0	0	0
4	C	107	0	0	1	0
All	All	5758	0	5427	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:ARG:HH11	1:C:136:ARG:HG2	1.49	0.77
1:C:136:ARG:HG2	1:C:136:ARG:NH1	2.01	0.73
1:A:105:ILE:HG22	1:A:134:LEU:HD12	1.73	0.69
1:A:205:HIS:HD2	1:A:207:ILE:H	1.43	0.66
1:C:105:ILE:HG22	1:C:134:LEU:HD12	1.79	0.64
1:C:38:LEU:HD21	1:C:41:GLN:HG3	1.81	0.62
1:C:136:ARG:NH2	4:C:499:HOH:O	2.32	0.60
1:C:204:ASP:OD1	1:C:211:LYS:HD3	2.01	0.60
1:A:162:ASP:HA	4:A:658:HOH:O	2.02	0.59
1:B:42:VAL:HG12	1:B:43:ARG:HG3	1.85	0.58
4:A:686:HOH:O	1:C:162:ASP:HB3	2.03	0.58
1:A:105:ILE:CG2	1:A:134:LEU:HD12	2.34	0.58
1:B:100:LEU:HD11	1:B:210:GLU:HB3	1.85	0.57
1:C:88:VAL:HB	1:C:224:MSE:HE3	1.87	0.57
1:A:24:ARG:NH2	4:A:679:HOH:O	2.39	0.55
1:B:15:ALA:HB1	1:B:232:ILE:HG12	1.88	0.55
1:B:2:THR:HB	1:B:79:PHE:CD2	2.43	0.53
1:B:105:ILE:HG22	1:B:134:LEU:HD12	1.92	0.52
1:C:100:LEU:HD11	1:C:210:GLU:HB3	1.91	0.52
1:B:205:HIS:HD2	1:B:208:THR:H	1.56	0.52
1:B:163:GLN:HB3	1:B:164:PRO:HD2	1.93	0.50
1:B:101:ARG:HB3	1:B:220:THR:HG21	1.93	0.50
1:A:100:LEU:HD11	1:A:210:GLU:HB3	1.93	0.49
1:C:105:ILE:CG2	1:C:134:LEU:HD12	2.43	0.49
1:B:205:HIS:CD2	1:B:208:THR:H	2.31	0.48
1:C:38:LEU:HD21	1:C:41:GLN:CG	2.43	0.47
1:C:18:LEU:HD11	1:C:85:LEU:HD23	1.98	0.46
1:A:173:PHE:CZ	1:B:119:ARG:HD3	2.51	0.46
1:C:100:LEU:O	1:C:101:ARG:HB2	2.16	0.46
1:B:229:LEU:O	1:B:232:ILE:HG22	2.16	0.45
1:A:24:ARG:NH2	3:A:504:SO4:O1	2.39	0.44
1:A:216:GLU:O	1:A:220:THR:HG22	2.18	0.44
1:B:144:ARG:HH21	1:B:226:GLU:HG2	1.83	0.43
1:B:98:LEU:HD22	1:B:149:HIS:CD2	2.53	0.43
1:B:100:LEU:O	1:B:101:ARG:HB2	2.19	0.42
1:A:119:ARG:HD3	1:B:173:PHE:CZ	2.55	0.42
1:B:205:HIS:CD2	1:B:207:ILE:HB	2.55	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:ARG:HH11	1:C:136:ARG:CG	2.22	0.42
1:A:105:ILE:HG22	1:A:134:LEU:CD1	2.45	0.42
1:C:21:ASP:HA	1:C:22:PRO:HD2	1.95	0.41
1:A:173:PHE:CZ	1:B:119:ARG:NE	2.88	0.41
1:A:204:ASP:OD1	1:A:211:LYS:HG3	2.20	0.41
1:C:205:HIS:CD2	1:C:208:THR:H	2.39	0.41
1:A:217:ARG:O	1:A:221:PHE:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/254 (92%)	229 (98%)	3 (1%)	1 (0%)	34	29
1	B	231/254 (91%)	224 (97%)	7 (3%)	0	100	100
1	C	234/254 (92%)	228 (97%)	5 (2%)	1 (0%)	34	29
All	All	698/762 (92%)	681 (98%)	15 (2%)	2 (0%)	41	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASP
1	C	162	ASP

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/202 (95%)	189 (98%)	3 (2%)	62	67
1	B	190/202 (94%)	187 (98%)	3 (2%)	62	67
1	C	192/202 (95%)	188 (98%)	4 (2%)	53	57
All	All	574/606 (95%)	564 (98%)	10 (2%)	59	65

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	162	ASP
1	A	179	GLU
1	B	2	THR
1	B	55	LYS
1	B	179	GLU
1	C	2	THR
1	C	136	ARG
1	C	162	ASP
1	C	179	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	161	HIS
1	A	165	ASN
1	A	168	GLN
1	A	205	HIS
1	B	124	ASN
1	B	161	HIS
1	B	205	HIS
1	C	205	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	B	302	-	4,4,4	0.36	0	6,6,6	0.15	0
3	SO4	A	504	-	4,4,4	0.29	0	6,6,6	0.23	0
3	SO4	A	503	-	4,4,4	0.17	0	6,6,6	0.17	0
3	SO4	B	303	-	4,4,4	0.30	0	6,6,6	0.13	0
3	SO4	C	301	-	4,4,4	0.37	0	6,6,6	0.33	0
3	SO4	C	302	-	4,4,4	0.35	0	6,6,6	0.12	0
3	SO4	C	304	-	4,4,4	0.19	0	6,6,6	0.21	0
3	SO4	C	303	-	4,4,4	0.28	0	6,6,6	0.13	0
3	SO4	B	304	-	4,4,4	0.36	0	6,6,6	0.14	0
3	SO4	B	301	-	4,4,4	0.26	0	6,6,6	0.19	0
3	SO4	A	502	-	4,4,4	0.37	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	504	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	226/254 (88%)	0.08	12 (5%) 26 35	16, 27, 57, 79	0
1	B	226/254 (88%)	0.33	19 (8%) 11 15	18, 34, 66, 85	0
1	C	226/254 (88%)	0.23	13 (5%) 23 31	16, 27, 57, 83	0
All	All	678/762 (88%)	0.21	44 (6%) 18 25	16, 29, 61, 85	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	THR	6.9
1	A	211	LYS	5.5
1	C	211	LYS	5.2
1	C	212	THR	5.0
1	B	212	THR	3.8
1	C	9[A]	PRO	3.8
1	B	211	LYS	3.4
1	C	213	THR	3.4
1	B	9	PRO	3.3
1	A	212	THR	3.3
1	A	164	PRO	3.2
1	C	164	PRO	3.1
1	A	214	PRO	3.1
1	C	2	THR	3.1
1	B	215	GLN	3.1
1	A	210	GLU	3.0
1	B	101	ARG	3.0
1	C	214	PRO	2.9
1	C	210	GLU	2.9
1	C	209	GLY	2.8
1	B	214	PRO	2.6
1	A	165	ASN	2.6
1	C	208	THR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	164	PRO	2.6
1	B	208	THR	2.5
1	B	163	GLN	2.5
1	B	210	GLU	2.5
1	B	216	GLU	2.4
1	B	213	THR	2.4
1	C	96	GLU	2.4
1	C	101	ARG	2.3
1	A	96	GLU	2.3
1	A	99	LYS	2.3
1	A	215	GLN	2.2
1	B	8	PRO	2.2
1	A	213	THR	2.1
1	B	79	PHE	2.1
1	B	162[A]	ASP	2.1
1	A	163	GLN	2.1
1	B	18	LEU	2.0
1	C	69	ALA	2.0
1	A	9	PRO	2.0
1	B	69	ALA	2.0
1	B	128	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	501	1/1	0.82	0.23	67,67,67,67	0
3	SO4	A	502	5/5	0.90	0.21	69,69,73,74	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	B	304	5/5	0.91	0.16	68,74,75,78	0
3	SO4	C	303	5/5	0.92	0.20	70,72,76,80	0
3	SO4	C	302	5/5	0.93	0.21	56,61,62,63	0
3	SO4	C	301	5/5	0.93	0.15	42,46,51,52	0
3	SO4	A	504	5/5	0.96	0.15	45,50,51,55	0
3	SO4	B	302	5/5	0.96	0.11	56,57,61,63	0
3	SO4	B	303	5/5	0.97	0.14	64,65,68,69	0
3	SO4	A	503	5/5	0.99	0.08	30,30,32,32	0
3	SO4	B	301	5/5	0.99	0.07	47,49,51,52	0
3	SO4	C	304	5/5	0.99	0.07	33,34,35,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.