



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 10:17 pm GMT

PDB ID : 5MA4
Title : GFP-binding DARPin fusion gc_K7
Authors : Hansen, S.; Stueber, J.; Ernst, P.; Koch, A.; Bojar, D.; Batyuk, A.; Plueckthun, A.
Deposited on : 2016-11-03
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	243	 86% 9% 5%
2	A	298	 9% 85% 11% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9217 atoms, of which 4207 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	B	230	3933	1255	1954	334	383	7	0	34	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP P42212
B	-3	PRO	-	expression tag	UNP P42212
B	-2	GLY	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	0	MET	-	expression tag	UNP P42212
B	1	VAL	-	expression tag	UNP P42212
B	64	LEU	PHE	conflict	UNP P42212
B	?	-	SER	chromophore	UNP P42212
B	?	-	TYR	chromophore	UNP P42212
B	66	CRO	GLY	chromophore	UNP P42212
B	231	LEU	HIS	conflict	UNP P42212
B	239	GLN	-	expression tag	UNP P42212
B	240	ALA	-	expression tag	UNP P42212

- Molecule 2 is a protein called K7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	A	294	4559	1459	2253	392	454	1	0	25	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	318	Total 318	O 318	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	407	Total 407	O 407	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Green fluorescent protein

Chain B: 



- Molecule 2: K7

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.88Å 92.34Å 56.53Å 90.00° 114.60° 90.00°	Depositor
Resolution (Å)	47.29 – 1.40 47.29 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.29-1.40) 99.9 (47.29-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 1.40Å)	Xtrriage
Refinement program	PHENIX (1.11_2567: ???)	Depositor
R, R_{free}	0.142 , 0.170 0.142 , 0.171	Depositor DCC
R_{free} test set	4853 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	15.5	Xtrriage
Anisotropy	0.112	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.021 for l,-k,h	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	9217	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.66	0/2138	0.77	0/2889
2	A	0.63	0/2448	0.76	4/3335 (0.1%)
All	All	0.64	0/4586	0.77	4/6224 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	155[A]	ASP	CB-CG-OD1	5.93	123.64	118.30
2	A	155[B]	ASP	CB-CG-OD1	5.93	123.64	118.30
2	A	160	ASP	CB-CG-OD1	5.39	123.16	118.30
2	A	297	ASP	CB-CG-OD1	5.31	123.08	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1979	1954	1814	25	0
2	A	2306	2253	2138	60	0
3	A	407	0	0	46	2
3	B	318	0	0	19	1
All	All	5010	4207	3952	86	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:155[B]:ASP:OD2	3:A:401:HOH:O	1.55	1.24
2:A:97[A]:GLU:O	3:A:402:HOH:O	1.66	1.11
2:A:31:ARG:O	3:A:403:HOH:O	1.71	1.08
1:B:168[B]:ARG:NH2	3:B:301:HOH:O	1.87	1.06
2:A:138[B]:ASP:OD2	3:A:404:HOH:O	1.75	1.02
2:A:97[B]:GLU:O	3:A:402:HOH:O	1.76	1.02
2:A:155[B]:ASP:OD1	3:A:406:HOH:O	1.83	0.96
2:A:246:ASP:OD1	3:A:407:HOH:O	1.86	0.94
1:B:113:LYS:NZ	3:B:303:HOH:O	2.01	0.93
2:A:100:LEU:N	3:A:402:HOH:O	2.06	0.88
1:B:144[A]:ASN:OD1	3:B:302:HOH:O	1.92	0.88
1:B:142[B]:GLU:OE2	1:B:144[B]:ASN:ND2	2.08	0.86
2:A:109:ARG:NH1	3:A:410:HOH:O	2.09	0.86
1:B:157:GLN:NE2	3:B:306:HOH:O	2.08	0.84
2:A:247:ASP:O	3:A:408:HOH:O	1.97	0.82
1:B:73:ARG:NH1	3:B:304:HOH:O	2.01	0.81
2:A:101:LYS:N	3:A:402:HOH:O	1.90	0.79
2:A:208:ALA:O	3:A:409:HOH:O	2.04	0.76
2:A:251[A]:PHE:N	3:A:412:HOH:O	2.19	0.75
2:A:249:ALA:N	3:A:407:HOH:O	2.18	0.74
2:A:249:ALA:HB3	3:A:405:HOH:O	1.88	0.72
1:B:5:GLU:OE2	3:B:305:HOH:O	2.07	0.70
2:A:144[A]:LYS:NZ	3:A:421:HOH:O	2.28	0.67
1:B:162[B]:LYS:NZ	3:B:312:HOH:O	2.27	0.67
1:B:225[B]:THR:CG2	2:A:122[B]:TRP:CH2	2.78	0.66
1:B:147[B]:SER:OG	3:B:307:HOH:O	2.13	0.66
2:A:251[B]:PHE:N	3:A:412:HOH:O	2.28	0.66
1:B:90[A]:GLU:OE1	3:B:309:HOH:O	2.13	0.66
1:B:133:ASP:OD2	3:B:308:HOH:O	2.13	0.66
2:A:250:GLY:N	3:A:407:HOH:O	2.06	0.65
2:A:251[B]:PHE:HA	3:A:416:HOH:O	1.98	0.64
2:A:296:GLU:OE2	3:A:411:HOH:O	2.15	0.63
3:B:304:HOH:O	2:A:122[B]:TRP:O	2.05	0.63
2:A:249:ALA:C	3:A:412:HOH:O	2.37	0.62
2:A:34:MET:N	3:A:403:HOH:O	2.36	0.59
1:B:225[B]:THR:HG21	2:A:122[B]:TRP:CH2	2.38	0.58
2:A:183[B]:GLU:HG2	3:A:641:HOH:O	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225[B]:THR:HG23	2:A:122[B]:TRP:CH2	2.40	0.57
2:A:144[A]:LYS:HD3	3:A:421:HOH:O	2.03	0.57
1:B:142[B]:GLU:CD	1:B:144[B]:ASN:HD21	2.09	0.56
2:A:101:LYS:NZ	3:A:415:HOH:O	2.23	0.55
1:B:142[B]:GLU:CD	1:B:144[B]:ASN:ND2	2.59	0.55
3:B:302:HOH:O	2:A:144[A]:LYS:NZ	2.40	0.54
2:A:246:ASP:OD1	3:A:405:HOH:O	2.18	0.53
2:A:57:ARG:NH2	3:A:427:HOH:O	2.42	0.53
3:B:307:HOH:O	2:A:183[B]:GLU:HG3	2.08	0.53
2:A:246:ASP:HA	3:A:405:HOH:O	2.10	0.52
2:A:35:ALA:N	3:A:403:HOH:O	2.13	0.50
2:A:128:ILE:HA	2:A:131[B]:VAL:HG22	1.94	0.50
2:A:14:LEU:HD22	2:A:15:GLY:N	2.27	0.50
1:B:73:ARG:NE	3:B:314:HOH:O	2.35	0.49
2:A:61:GLU:O	2:A:65:VAL:HG13	2.12	0.49
2:A:249:ALA:CB	3:A:412:HOH:O	2.61	0.48
1:B:186[B]:THR:HG22	3:B:393:HOH:O	2.12	0.48
2:A:14:LEU:HD13	2:A:15:GLY:H	1.79	0.47
2:A:100:LEU:CA	3:A:402:HOH:O	2.58	0.47
2:A:246:ASP:CG	3:A:405:HOH:O	2.53	0.46
2:A:267:GLU:HG3	3:A:716:HOH:O	2.16	0.46
2:A:57:ARG:NH1	3:A:439:HOH:O	2.49	0.45
2:A:251[B]:PHE:CD1	3:A:412:HOH:O	2.68	0.45
2:A:157:GLY:N	3:A:430:HOH:O	2.43	0.45
2:A:17:LYS:O	2:A:21:ALA:N	2.43	0.45
2:A:42:ALA:O	2:A:50:PRO:HD3	2.17	0.44
3:B:358:HOH:O	2:A:122[B]:TRP:CD1	2.57	0.44
2:A:183[B]:GLU:CD	2:A:183[B]:GLU:H	2.14	0.43
1:B:155:ASP:OD2	1:B:158:LYS:HD2	2.19	0.43
2:A:251[B]:PHE:CD2	3:A:412:HOH:O	2.72	0.43
1:B:73:ARG:NH2	3:B:304:HOH:O	2.53	0.42
2:A:122[B]:TRP:HD1	3:A:555:HOH:O	2.03	0.42
2:A:248:GLN:C	3:A:407:HOH:O	2.53	0.41
2:A:198:GLU:HG3	3:A:545:HOH:O	2.20	0.41
2:A:249:ALA:HB1	3:A:412:HOH:O	2.20	0.41
2:A:57:ARG:HG2	2:A:57:ARG:HH11	1.86	0.41
2:A:250:GLY:N	3:A:412:HOH:O	2.51	0.41
1:B:153:MET:HB2	1:B:162[B]:LYS:HG3	2.03	0.41
2:A:134:LYS:NZ	3:A:451:HOH:O	2.55	0.40
1:B:6:GLU:HG2	3:B:579:HOH:O	2.20	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:675:HOH:O	3:A:692:HOH:O[2_545]	1.77	0.43
3:B:391:HOH:O	3:A:415:HOH:O[2_546]	1.80	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	258/243 (106%)	256 (99%)	2 (1%)	0	100	100
2	A	317/298 (106%)	306 (96%)	11 (4%)	0	100	100
All	All	575/541 (106%)	562 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	232/210 (110%)	231 (100%)	1 (0%)	91	78
2	A	242/220 (110%)	231 (96%)	11 (4%)	27	4
All	All	474/430 (110%)	462 (98%)	12 (2%)	49	14

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	0	MET
2	A	14	LEU
2	A	57	ARG
2	A	65	VAL
2	A	101	LYS
2	A	122[A]	TRP
2	A	122[B]	TRP
2	A	154	ILE
2	A	156[A]	ASN
2	A	156[B]	ASN
2	A	248	GLN
2	A	281	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	B	66	1	23,23,24	1.77	6 (26%)	30,32,34	1.44	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CRO	C1-N2	4.99	1.39	1.32
1	B	66	CRO	CG2-CB2	3.78	1.54	1.46
1	B	66	CRO	CA1-C1	-2.63	1.47	1.51
1	B	66	CRO	C1-N3	2.60	1.41	1.37
1	B	66	CRO	CA2-N2	2.57	1.44	1.38
1	B	66	CRO	CB2-CA2	-2.14	1.33	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	CRO	C2-N3-C1	3.80	109.89	107.97
1	B	66	CRO	O3-C3-CA3	-3.54	115.70	126.39
1	B	66	CRO	CG2-CB2-CA2	-2.33	127.09	129.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	229/243 (94%)	-0.33	3 (1%) 77 75	10, 20, 36, 53	0
2	A	294/298 (98%)	0.07	28 (9%) 8 7	11, 21, 67, 99	0
All	All	523/541 (96%)	-0.10	31 (5%) 22 20	10, 20, 56, 99	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	14	LEU	8.7
2	A	69	TYR	5.9
2	A	33	LEU	5.7
2	A	13	ASP	4.5
1	B	231[A]	LEU	4.0
2	A	250	GLY	3.6
2	A	17	LYS	3.5
2	A	57	ARG	3.4
2	A	156[A]	ASN	3.4
2	A	247	ASP	3.3
2	A	15	GLY	3.2
2	A	248	GLN	3.2
2	A	68	LYS	3.1
2	A	21	ALA	3.0
2	A	281	LYS	3.0
2	A	35	ALA	2.9
2	A	16	LYS	2.9
2	A	37	GLY	2.8
2	A	31	ARG	2.7
2	A	36	ASN	2.6
2	A	34	MET	2.6
2	A	18	LEU	2.4
2	A	43	ALA	2.4
2	A	306	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	38	ALA	2.4
2	A	32	ILE	2.3
2	A	30	VAL	2.2
2	A	20	GLU	2.2
2	A	154	ILE	2.1
1	B	0	MET	2.1
1	B	230	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	B	66	22/23	0.99	0.07	10,12,18,18	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.