

wwPDB X-ray Structure Validation Summary Report (i)

Oct 1, 2023 – 11:22 PM EDT

PDB ID	:	6MDZ
Title	:	Human Argonaute2-miR-122 bound to a target RNA with two central mis-
		matches (bu2)
Authors	:	Sheu-Gruttadauria, J.; MacRae, I.J.
Deposited on	:	2018-09-05
Resolution	:	3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 3.40 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 14382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein argonaute-2.

Mol	Chain	Residues		Α	toms		ZeroOcc	AltConf	Trace	
1	А	804	Total 6446	C 4106	N 1158	0 1141	S 41	0	0	0
1	В	796	Total 6388	-	N 1148	0 1128	S 39	0	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	387	ASP	SER	engineered mutation	UNP Q9UKV8
А	669	ALA	ASP	engineered mutation	UNP Q9UKV8
A	824	ALA	SER	engineered mutation	UNP Q9UKV8
А	828	ASP	SER	engineered mutation	UNP Q9UKV8
А	831	ASP	SER	engineered mutation	UNP Q9UKV8
А	834	ALA	SER	engineered mutation	UNP Q9UKV8
В	387	ASP	SER	engineered mutation	UNP Q9UKV8
В	669	ALA	ASP	engineered mutation	UNP Q9UKV8
В	824	ALA	SER	engineered mutation	UNP Q9UKV8
В	828	ASP	SER	engineered mutation	UNP Q9UKV8
В	831	ASP	SER	engineered mutation	UNP Q9UKV8
В	834	ALA	SER	engineered mutation	UNP Q9UKV8

There are 12 discrepancies between the modelled and reference sequences:

• Molecule 2 is a RNA chain called RNA (5'-R(P*UP*GP*GP*AP*GP*UP*GP*UP*GP*AP *CP*AP*AP*UP*GP*UP*GP*UP*U)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
0	C	20	Total	С	Ν	0	Р	0	0	0
			433	192	77	144	20	0		
0	Л	14	Total	С	Ν	0	Р	0	0	0
	2 D		308	136	58	100	14	0	U	U

• Molecule 3 is a RNA chain called RNA (5'-R(P*AP*AP*CP*AP*CP*AP*CP*AP*UP*UP *UP*CP*CP*AP*CP*AP*CP*AP*CP*AP*AP*A)-3').



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	23	Total	С	Ν	0	Р	0	0	0
0	Ľ		480	217	85	155	23	0		
2	Б	17	Total	С	Ν	0	Р	0	0	1
0	3 F	17	299	131	50	102	16	0	0	1

• Molecule 4 is PHENOL (three-letter code: IPH) (formula: C_6H_6O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 6 1 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 6 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 6 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 6 1 \end{array}$	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	107.61Å 138.29Å 152.61Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.29 - 3.40	Depositor
% Data completeness	99.3 (38.29-3.40)	Depositor
(in resolution range)		-
R _{merge}	0.17	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.00 (at 3.40 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.9_{1692}	Depositor
R, R_{free}	0.220 , 0.264	Depositor
Wilson B-factor $(Å^2)$	69.3	Xtriage
Anisotropy	0.233	Xtriage
L-test for twinning ²	$ L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14382	wwPDB-VP
Average B, all atoms $(Å^2)$	63.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 24.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6319e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



	The second		Der	T 1.	B	ond leng	gths	Bond angles			
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	IPH	В	902	-	7,7,7	0.36	0	8,8,8	0.24	0	
4	IPH	А	902	-	7,7,7	0.36	0	8,8,8	0.24	0	
4	IPH	В	901	-	7,7,7	0.37	0	8,8,8	0.23	0	
4	IPH	А	901	-	7,7,7	0.33	0	8,8,8	0.27	0	

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IPH	В	902	-	-	-	0/1/1/1
4	IPH	А	902	-	-	-	0/1/1/1
4	IPH	В	901	-	-	-	0/1/1/1
4	IPH	А	901	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

