

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 04:40 AM EDT

PDB ID : 2MJC BMRB ID : 19716

Title: Zn-binding domain of eukaryotic translation initiation factor 3, subunit G

Authors: Al-Abdul-Wahid, M.; Menade, M.; Xie, J.; Kozlov, G.; Gehring, K.

Deposited on : 2014-01-03

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 79%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive
Metric	$(\# ext{Entries})$	$(\# ext{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length		Quality of chain				
		0.4						
1	A	34	35%	12%	24%	29%		



2 Ensemble composition and analysis (i)

This entry contains 15 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core	Medoid model				
1 A:11-A:26 (16)		0.38	1		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 14, 15
2	10, 13



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 384 atoms, of which 191 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Eukaryotic translation initiation factor 3 subunit G.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	24	Total	С	Н	N	О	S	0
1	A	A 24 38	383	120	191	36	33	3	U

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	PRO	-	expression tag	UNP O75821
A	3	LEU	-	expression tag	UNP O75821
A	4	GLY	-	expression tag	UNP O75821
A	5	SER	-	expression tag	UNP O75821

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
9	Λ	1	Total Zn	
2	A	1	1 1	



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Eukaryotic translation initiation factor 3 subunit G



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: Eukaryotic translation initiation factor 3 subunit G





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 15 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	structure solution	
XPLOR-NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	345
Number of shifts mapped to atoms	254
Number of unparsed shifts	0
Number of shifts with mapping errors	91
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	79%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	
1	A	127 120		121	2±2	
All	All	1920	1800	1815	29	

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 18 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models		
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total	
1:A:14:ARG:NE	1:A:14:ARG:O	0.52	2.42	5	1	
1:A:21:TRP:O	1:A:23:THR:N	0.51	2.43	13	2	
1:A:14:ARG:NE	1:A:14:ARG:C	0.47	2.67	5	1	
1:A:13:CYS:SG	1:A:14:ARG:N	0.47	2.87	3	2	
1:A:11:VAL:HG23	1:A:21:TRP:CD1	0.47	2.45	3	1	



6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed Outliers Perc		Favoured Allowed Outliers Percenti		entiles
1	A	16/34 (47%)	10±1 (60±7%)	5±1 (32±7%)	1±1 (8±5%)	2	15	
All	All	240/510 (47%)	145 (60%)	77 (32%)	18 (8%)	2	15	

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	21	TRP	12
1	A	13	CYS	5
1	A	26	PRO	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric		Outliers	Perce	ntiles
1	A	15/30 (50%)	15±0 (97±3%)	0±0 (3±3%)	48	90
All	All	225/450~(50%)	219 (97%)	6 (3%)	48	90

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	23	THR	2
1	A	20	HIS	2
1	A	14	ARG	1
1	A	17	LYS	1



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 79% for the well-defined parts and 78% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	345
Number of shifts mapped to atoms	254
Number of unparsed shifts	0
Number of shifts with mapping errors	91
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	56

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 91) occurrences are reported below.

I int ID Chain		D	Т	A 4		Shift Data	
List ID	ist ID \mid Chain \mid Re	Res	Type	Type Atom		Uncertainty	Ambiguity
1	A	2	PRO	CA	35.3515	0.0000	1
1	A	2	PRO	HA	4.4659	0.0000	1
1	A	2	PRO	СВ	32.3343	0.0000	1
1	A	2	PRO	HB2	2.3148	0.0000	2
1	A	2	PRO	CG	27.1066	0.0000	1
1	A	2	PRO	HG2	2.0099	0.0000	2
1	A	2	PRO	CD	49.7183	0.0000	1
1	A	2	PRO	HD2	3.5683	0.0000	2
1	A	3	LEU	N	122.6532	0.0000	1
1	A	3	LEU	Н	8.5603	0.0000	1
1	A	3	LEU	CA	27.2875	0.0000	1
1	A	3	LEU	HA	4.343	0.0000	1
1	A	3	LEU	СВ	43.0704	0.0000	1
1	A	3	LEU	HB2	1.6537	0.0000	2

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from previous page					Shift Data			
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity	
1	A	3	LEU	CG	27.099	0.0000	1	
1	A	3	LEU	HG	1.6503	0.0000	1	
1	A	3	LEU	CD1	24.876	0.0000	2	
1	A	3	LEU	HD11	0.8942	0.0000	2	
1	A	3	LEU	HD12	0.8942	0.0000	2	
1	A	3	LEU	HD13	0.8942	0.0000	2	
1	A	3	LEU	CD2	51.4641	0.0000	2	
1	A	3	LEU	HD21	0.911	0.0000	2	
1	A	3	LEU	HD22	0.911	0.0000	2	
1	A	3	LEU	HD23	0.911	0.0000	2	
1	A	4	GLY	N	109.8654	0.0000	1	
1	A	4	GLY	Н	8.4436	0.0000	1	
1	A	4	GLY	CA	45.9357	0.0000	1	
1	A	4	GLY	HA2	3.9513	0.0000	2	
1	A	5	SER	N	115.8753	0.0000	1	
1	A	5	SER	Н	8.2676	0.0000	1	
1	A	5	SER	CA	30.4234	0.0000	1	
1	A	5	SER	HA	4.4489	0.0000	1	
1	A	5	SER	СВ	35.9952	0.0000	1	
1	A	5	SER	HB2	3.8594	0.0000	2	
1	A	6	LYS	N	123.354	0.0000	1	
1	A	6	LYS	Н	8.4447	0.0000	1	
1	A	6	LYS	CA	28.4489	0.0000	1	
1	A	6	LYS	HA	4.3464	0.0000	1	
1	A	6	LYS	СВ	33.4432	0.0000	1	
1	A	6	LYS	HB2	1.7139	0.0000	2	
1	A	6	LYS	HB3	1.3657	0.0000	2	
1	A	6	LYS	CG	24.7879	0.0000	1	
1	A	6	LYS	HG2	1.4075	0.0000	2	
1	A	6	LYS	HG3	1.4099	0.0000	2	
1	A	6	LYS	HD2	1.717	0.0000	2	
1	A	6	LYS	HE2	2.9744	0.0000	2	
1	A	7	GLY	N	109.8433	0.0000	1	
1	A	7	GLY	Н	8.4302	0.0000	1	
1	A	7	GLY	CA	46.0171	0.0000	1	
1	A	7	GLY	HA2	4.8107	0.0000	2	
1	A	8	GLN	N	120.0368	0.0000	1	
1	A	8	GLN	Н	8.1927	0.0000	1	
1	A	8	GLN	CA	56.3108	0.0000	1	
1	A	8	GLN	HA	4.3063	0.0000	1	
1	A	8	GLN	СВ	30.4397	0.0000	1	

Continued on next page...



Continued from previous page...

	Continued from previous page				Shift Data			
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity	
1	A	8	GLN	HB2	2.3152	0.0000	2	
1	A	8	GLN	HB3	1.9513	0.0000	2	
1	A	8	GLN	HG2	2.3152	0.0000	2	
1	A	8	GLN	HG3	1.9513	0.0000	2	
1	A	33	PRO	CA	35.2803	0.0000	1	
1	A	33	PRO	HA	4.3975	0.0000	1	
1	A	33	PRO	СВ	32.1147	0.0000	1	
1	A	33	PRO	HB2	2.219	0.0000	2	
1	A	33	PRO	CG	27.1892	0.0000	1	
1	A	33	PRO	HG2	1.9763	0.0000	2	
1	A	33	PRO	CD	49.7856	0.0000	1	
1	A	33	PRO	HD2	3.5583	0.0000	2	
1	A	34	MET	N	120.7562	0.0000	1	
1	A	34	MET	Н	8.45	0.0000	1	
1	A	34	MET	CA	27.566	0.0000	1	
1	A	34	MET	HA	4.4415	0.0000	1	
1	A	34	MET	СВ	32.8394	0.0000	1	
1	A	34	MET	HB2	2.1	0.0000	2	
1	A	34	MET	HB3	1.9965	0.0000	2	
1	A	34	MET	CG	32.1922	0.0000	1	
1	A	34	MET	HG2	2.6019	0.0000	2	
1	A	34	MET	HG3	2.5447	0.0000	2	
1	A	34	MET	CE	44.9347	0.0000	1	
1	A	34	MET	HE1	2.0744	0.0000	1	
1	A	34	MET	HE2	2.0744	0.0000	1	
1	A	34	MET	HE3	2.0744	0.0000	1	
1	A	35	GLN	N	126.2075	0.0000	1	
1	A	35	GLN	Н	7.929	0.0000	1	
1	A	35	GLN	CA	29.3391	0.0000	1	
1	A	35	GLN	HA	4.1533	0.0000	1	
1	A	35	GLN	СВ	30.4888	0.0000	1	
1	A	35	GLN	HB2	2.0977	0.0000	2	
1	A	35	GLN	HB3	1.9189	0.0000	2	
1	A	35	GLN	CG	34.2691	0.0000	1	
1	A	35	GLN	HG2	2.2711	0.0000	2	
1	A	35	GLN	HG3	1.7531	0.0000	2	

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.



Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	34	0.00 ± 0.00	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	29	1.04 ± 0.56	None needed (imprecise)
¹³ C′	0	_	None (insufficient data)
^{15}N	31	-0.57 ± 1.06	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 79%, i.e. 168 atoms were assigned a chemical shift out of a possible 212. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	63/79 (80%)	32/32 (100%)	16/32~(50%)	15/15 (100%)
Sidechain	89/115 (77%)	59/74 (80%)	30/34 (88%)	0/7 (0%)
Aromatic	16/18 (89%)	8/10 (80%)	7/7 (100%)	1/1 (100%)
Overall	168/212 (79%)	99/116 (85%)	53/73 (73%)	16/23 (70%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	27	TYR	CE2	34.12	111.68 - 124.17	-67.1
1	A	21	TRP	CZ2	30.73	107.20 - 121.33	-59.1
1	A	27	TYR	CD2	49.20	125.28 - 140.14	-56.2
1	A	21	TRP	CH2	40.90	116.19 – 131.43	-54.4
1	A	21	TRP	CZ3	38.03	113.48 - 129.28	-52.8
1	A	20	HIS	CE1	25.61	126.08 - 149.12	-48.6
1	A	21	TRP	CE3	36.83	111.58 - 129.41	-46.9
1	A	21	TRP	CD1	42.03	117.34 - 135.80	-45.8
1	A	22	THR	CG2	50.86	16.06 - 27.03	26.7
1	A	23	THR	CG2	50.74	16.06 - 27.03	26.6
1	A	30	THR	CG2	49.78	16.06 - 27.03	25.7
1	A	28	LYS	CG	52.03	19.35 - 30.45	24.4
1	A	20	HIS	CD2	44.37	103.95 - 136.66	-23.2
1	A	10	ILE	CG2	45.27	10.93 - 24.12	21.0
1	A	15	ILE	CD1	46.01	5.18 - 21.60	19.9
1	A	11	VAL	CG1	48.23	14.71 - 28.29	19.7
1	A	33	PRO	CA	35.28	55.85 - 70.84	-18.7

Continued on next page...



Continued from previous page...


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	2	PRO	CA	35.35	55.85 - 70.84	-18.7
1	A	5	SER	СВ	36.00	56.28 - 71.32	-18.5
1	A	26	PRO	CA	35.88	55.85 - 70.84	-18.3
1	A	15	ILE	CG2	41.53	10.93 - 24.12	18.2
1	A	12	SER	СВ	36.52	56.28 - 71.32	-18.1
1	A	11	VAL	CG2	48.44	13.71 - 28.88	17.9
1	A	23	THR	СВ	40.88	61.12 - 78.27	-16.8
1	A	10	ILE	CD1	40.62	5.18 - 21.60	16.6
1	A	3	LEU	CD2	51.46	15.73 - 32.47	16.4
1	A	30	THR	СВ	41.92	61.12 - 78.27	-16.2
1	A	31	LEU	CD2	51.20	15.73 - 32.47	16.2
1	A	22	THR	СВ	42.00	61.12 - 78.27	-16.1
1	A	34	MET	CE	44.93	8.39 - 25.85	15.9
1	A	19	ASP	CA	24.84	44.71 - 64.67	-15.0
1	A	12	SER	CA	29.11	48.46 - 68.96	-14.4
1	A	29	ASP	CA	26.99	44.71 - 64.67	-13.9
1	A	5	SER	CA	30.42	48.46 - 68.96	-13.8
1	A	31	LEU	CA	27.09	45.17 - 66.21	-13.6
1	A	24	ARG	CA	26.03	45.44 - 68.13	-13.6
1	A	3	LEU	CA	27.29	45.17 - 66.21	-13.5
1	A	20	HIS	CA	25.99	45.04 - 67.94	-13.3
1	A	9	LYS	CA	28.38	46.18 - 67.77	-13.2
1	A	6	LYS	CA	28.45	46.18 - 67.77	-13.2
1	A	34	MET	CA	27.57	45.26 - 67.07	-13.1
1	A	35	GLN	CA	29.34	46.17 - 66.97	-13.1
1	A	17	LYS	CA	30.68	46.18 - 67.77	-12.2
1	A	28	LYS	CA	30.70	46.18 - 67.77	-12.2
1	A	27	TYR	CA	28.90	45.75 - 70.63	-11.8
1	A	21	TRP	CA	28.74	45.21 - 70.26	-11.6
1	A	14	ARG	CA	30.67	45.44 - 68.13	-11.5
1	A	30	THR	CA	34.44	49.41 - 75.05	-10.8
1	A	10	ILE	CA	33.21	48.30 - 75.08	-10.6
1	A	23	THR	CA	35.62	49.41 - 75.05	-10.4
1	A	11	VAL	CA	33.84	48.38 - 76.73	-10.1
1	A	22	THR	CA	37.99	49.41 - 75.05	-9.4
1	A	15	ILE	CA	36.41	48.30 - 75.08	-9.4
1	A	16	CYS	CA	30.67	40.80 - 75.33	-7.9
1	A	25	CYS	CA	30.89	40.80 - 75.33	-7.9
1	A	13	CYS	CA	31.86	40.80 - 75.33	-7.6



7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	548
Intra-residue ($ i-j =0$)	250
Sequential ($ i-j =1$)	131
Medium range ($ i-j >1$ and $ i-j <5$)	72
Long range (i-j ≥5)	89
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	6
Total dihedral-angle restraints	22
Number of unmapped restraints	0
Number of restraints per residue	16.8
Number of long range restraints per residue ¹	2.7

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	4.7	0.19
0.2-0.5 (Medium)	4.5	0.5
>0.5 (Large)	0.9	2.07



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	$\mathbf{Max} \ (^{\circ})$
1.0-10.0 (Small)	0.5	2.0
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dordensinda dom o	Count	% ¹	Vic	lated ³	3	Consis	tentl	$\overline{ m y~Violated^4}$
Restraints type	Count	%0°	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	250	45.6	6	2.4	1.1	1	0.4	0.2
Backbone-Backbone	20	3.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	145	26.5	6	4.1	1.1	1	0.7	0.2
Sidechain-Sidechain	85	15.5	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	131	23.9	10	7.6	1.8	3	2.3	0.5
Backbone-Backbone	64	11.7	5	7.8	0.9	2	3.1	0.4
Backbone-Sidechain	56	10.2	4	7.1	0.7	1	1.8	0.2
Sidechain-Sidechain	11	2.0	1	9.1	0.2	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	72	13.1	6	8.3	1.1	1	1.4	0.2
Backbone-Backbone	25	4.6	3	12.0	0.5	1	4.0	0.2
Backbone-Sidechain	26	4.7	2	7.7	0.4	0	0.0	0.0
Sidechain-Sidechain	21	3.8	1	4.8	0.2	0	0.0	0.0
Long range ($ i-j \ge 5$)	89	16.2	2	2.2	0.4	0	0.0	0.0
Backbone-Backbone	10	1.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	42	7.7	1	2.4	0.2	0	0.0	0.0
Sidechain-Sidechain	37	6.8	1	2.7	0.2	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	6	1.1	0	0.0	0.0	0	0.0	0.0
Total	548	100.0	24	4.4	4.4	5	0.9	0.9
Backbone-Backbone	119	21.7	8	6.7	1.5	3	2.5	0.5
Backbone-Sidechain	269	49.1	13	4.8	2.4	2	0.7	0.4
Sidechain-Sidechain	160	29.2	3	1.9	0.5	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	3	Magn (Å)	Morr (Å)	\mathbf{SD}^6 (Å)	Modian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	$SD^*(A)$	Median (Å)
1	3	6	1	1	0	11	0.34	1.85	0.49	0.13
2	1	6	2	1	0	10	0.39	2.07	0.57	0.18
3	2	5	1	0	0	8	0.23	0.47	0.11	0.23
4	1	4	1	2	0	8	0.45	1.98	0.59	0.25
5	2	5	1	1	0	9	0.35	1.48	0.41	0.17
6	2	5	2	1	0	10	0.31	1.08	0.27	0.23
7	3	8	2	2	0	15	0.25	1.22	0.28	0.13
8	2	4	1	1	0	8	0.41	1.4	0.39	0.27
9	2	5	1	1	0	9	0.38	1.68	0.47	0.27
10	3	5	2	1	0	11	0.36	1.71	0.44	0.25
11	3	5	1	1	0	10	0.31	1.16	0.3	0.2

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model ID	IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Mean (A)	Max (A)	SD (A)	Median (A)
12	1	7	1	2	0	11	0.28	1.09	0.28	0.22
13	3	4	1	1	0	9	0.41	1.83	0.51	0.26
14	3	5	1	1	0	10	0.27	0.62	0.15	0.22
15	2	7	2	2	0	13	0.26	1.14	0.27	0.13

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints




for a given fraction of the ensemble. In total, 518 (IR:244, SQ:121, MR:66, LR:87, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio		Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	3	5	0	0	8	1	6.7
2	0	0	0	0	2	2	13.3
1	1	0	0	0	2	3	20.0
1	0	0	1	0	2	4	26.7
0	0	0	0	0	0	5	33.3
0	0	0	0	0	0	6	40.0
1	0	0	0	0	1	7	46.7
0	0	0	0	0	0	8	53.3
0	1	0	0	0	1	9	60.0
0	1	0	0	0	1	10	66.7
0	1	0	0	0	1	11	73.3
0	0	0	0	0	0	12	80.0
0	0	0	0	0	0	13	86.7
0	0	0	1	0	1	14	93.3
1	3	1	0	0	5	15	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,334)	1:A:29:ASP:H	1:A:29:ASP:HB2	15	0.46	0.01	0.46
(1,434)	1:A:13:CYS:H	1:A:11:VAL:HA	15	0.27	0.01	0.28
(1,352)	1:A:13:CYS:H	1:A:12:SER:HB2	15	0.25	0.01	0.25
(1,352)	1:A:13:CYS:H	1:A:12:SER:HB3	15	0.25	0.01	0.25
(1,426)	1:A:30:THR:H	1:A:29:ASP:HA	15	0.25	0.03	0.25
(1,395)	1:A:21:TRP:H	1:A:20:HIS:H	15	0.18	0.07	0.14
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	14	1.45	0.4	1.44
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	14	1.45	0.4	1.44
(1,497)	1:A:16:CYS:H	1:A:17:LYS:HB3	11	0.13	0.03	0.12
(1,344)	1:A:12:SER:H	1:A:11:VAL:HB	10	0.12	0.01	0.12
(1,428)	1:A:30:THR:H	1:A:29:ASP:HB3	9	0.13	0.02	0.12
(1,284)	1:A:10:ILE:H	1:A:10:ILE:HB	7	0.15	0.05	0.13


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	2	2.07
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	2	2.07
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	4	1.98
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	4	1.98
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	1	1.85
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	1	1.85
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	13	1.83
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	13	1.83
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	10	1.71
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	10	1.71
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	9	1.68

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	9	1.68
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	5	1.48
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	5	1.48
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	8	1.4
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	8	1.4
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	7	1.22
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	7	1.22
(1,528)	1:A:16:CYS:HB2	1:A:27:TYR:HE1	11	1.16



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	C	$\%^{1}$	Vic	lated ⁵	3	\mid Consistently Violated $^4\mid$		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^{1}$
PHI	10	45.5	0	0.0	0.0	0	0.0	0.0
CHI1	12	54.5	2	16.7	9.1	0	0.0	0.0
Total	22	100.0	2	9.1	9.1	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Num	ber of	violations	Mean (°)	Max (°)	SD (°)	Median (°)
Model 1D	PHI	CHI1	Total	Mean ()	Max ()	SD ()	Median ()
1	0	0	0	0.0	0.0	0.0	0.0
2	0	2	2	1.2	1.2	0.0	1.2
3	0	0	0	0.0	0.0	0.0	0.0
4	0	1	1	1.8	1.8	0.0	1.8
5	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0.0	0.0	0.0	0.0
8	0	2	2	1.4	1.7	0.3	1.4
9	0	1	1	1.1	1.1	0.0	1.1
10	0	0	0	0.0	0.0	0.0	0.0
11	0	0	0	0.0	0.0	0.0	0.0
12	0	0	0	0.0	0.0	0.0	0.0
13	0	1	1	2.0	2.0	0.0	2.0
14	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0.0	0.0	0.0	0.0

10.2.1 Bar graph: Dihedral violation statistics for each model (i)





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

Dihedral-angle violation statistics for the ensemble (i) 10.3




Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Num	ber of	violated restraints	Fraction of the ensemble				
PHI	CHI1	Total	Count ¹	%			
0	0	0	1	6.7			
0	0	0	2	13.3			
0	1	1	3	20.0			
0	1	1	4	26.7			
0	0	0	5	33.3			
0	0	0	6	40.0			
0	0	0	7	46.7			
0	0	0	8	53.3			
0	0	0	9	60.0			
0	0	0	10	66.7			
0	0	0	11	73.3			
0	0	0	12	80.0			
0	0	0	13	86.7			
0	0	0	14	93.3			
0	0	0	15	100.0			

¹ Number of models with violations



10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,11)	1:A:19:ASP:N	1:A:19:ASP:CA	1:A:19:ASP:CB	1:A:19:ASP:CG	4	1.68	0.29	1.75
(1,10)	1:A:14:ARG:N	1:A:14:ARG:CA	1:A:14:ARG:CB	1:A:14:ARG:CG	3	1.13	0.05	1.1


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,11)	1:A:19:ASP:N	1:A:19:ASP:CA	1:A:19:ASP:CB	1:A:19:ASP:CG	13	2.0
(1,11)	1:A:19:ASP:N	1:A:19:ASP:CA	1:A:19:ASP:CB	1:A:19:ASP:CG	4	1.8
(1,11)	1:A:19:ASP:N	1:A:19:ASP:CA	1:A:19:ASP:CB	1:A:19:ASP:CG	8	1.7
(1,11)	1:A:19:ASP:N	1:A:19:ASP:CA	1:A:19:ASP:CB	1:A:19:ASP:CG	2	1.2
(1,10)	1:A:14:ARG:N	1:A:14:ARG:CA	1:A:14:ARG:CB	1:A:14:ARG:CG	2	1.2
(1,10)	1:A:14:ARG:N	1:A:14:ARG:CA	1:A:14:ARG:CB	1:A:14:ARG:CG	8	1.1
(1,10)	1:A:14:ARG:N	1:A:14:ARG:CA	1:A:14:ARG:CB	1:A:14:ARG:CG	9	1.1

