

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 11:17 AM EDT

PDB ID : 6MNX

Title : Structural basis of impaired hydrolysis in KRAS Q61H mutant

Authors: Bera, A.K.; Westover, K.D.

Deposited on : 2018-10-03

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16455 atoms, of which 8043 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GTPase KRas.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	167	Total	С	Н	N	О	S	0	4	0
1	Λ	107	2701	849	1345	236	263	8	0	T	0
1	В	168	Total	С	Н	N	О	S	0	1	0
1	Ъ	100	2668	842	1323	231	264	8	U	1	
1	С	166	Total	С	Н	N	О	S	0	1	0
1		100	2631	831	1304	228	260	8			
1	D	166	Total	С	Н	N	О	S	0	3	0
1	D	100	2665	840	1326	231	260	8	U	3	
1	E	169	Total	С	Н	N	Ο	S	0	2	0
1	Ľ	109	2701	851	1343	233	266	8	U	2	U
1	F	164	Total	С	Н	N	О	S	0	2	0
1	I.	104	2623	825	1311	227	253	7			U

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	HIS	GLN	engineered mutation	UNP P01116
В	61	HIS	GLN	engineered mutation	UNP P01116
С	61	HIS	GLN	engineered mutation	UNP P01116
D	61	HIS	GLN	engineered mutation	UNP P01116
Е	61	HIS	GLN	engineered mutation	UNP P01116
F	61	HIS	GLN	engineered mutation	UNP P01116

• Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
9	2 A	1	Total	С	Н	N	О	Р	0	0	
2			47	10	15	5	14	3	U	U	
2	В	1	Total	С	Н	N	О	Р	0	0	
2		Б	D	1	47	10	15	5	14	3	U
9	2 C	С	1	Total	С	Н	N	О	Р	0	0
		1	47	10	15	5	14	3	U	U	
2	D	1	Total	С	Н	N	О	Р	0	0	
	D	1	47	10	15	5	14	3		U	
2	E	1	Total	С	Н	N	О	Р	0	0	
		1	47	10	15	5	14	3	U	0	
2	F	E 1	Total	С	Н	N	О	Р	0	0	
		48	10	16	5	14	3	U	U		

 \bullet Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	Ε	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0



• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	43	Total O 43 43	0	0
4	В	28	Total O 28 28	0	0
4	С	34	Total O 34 34	0	0
4	D	34	Total O 34 34	0	0
4	E	30	Total O 30 30	0	0
4	F	8	Total O 8 8	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	56.77Å 127.67Å 145.03Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.91 - 2.20	Depositor
% Data completeness	92.1 (47.91-2.20)	Depositor
(in resolution range)	,	
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.62 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.14rc3_3199, PHENIX 1.14rc3_3199	Depositor
R, R_{free}	0.204 , 0.254	Depositor
Wilson B-factor (\mathring{A}^2)	30.6	Xtriage
Anisotropy	0.009	Xtriage
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	16455	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 14.45% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GTP	Е	201	3	26,34,34	1.07	3 (11%)	32,54,54	0.58	0
2	GTP	D	201	3	26,34,34	1.19	3 (11%)	32,54,54	0.63	0
2	GTP	С	201	3	26,34,34	0.96	2 (7%)	32,54,54	0.63	0
2	GTP	A	201	3	26,34,34	1.34	5 (19%)	32,54,54	0.58	0
2	GTP	В	201	3	26,34,34	1.09	3 (11%)	32,54,54	0.57	0
2	GTP	F	201	3	26,34,34	0.87	2 (7%)	32,54,54	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	Е	201	3	-	1/18/38/38	0/3/3/3
2	GTP	D	201	3	-	1/18/38/38	0/3/3/3
2	GTP	С	201	3	-	1/18/38/38	0/3/3/3
2	GTP	A	201	3	-	1/18/38/38	0/3/3/3
2	GTP	В	201	3	-	1/18/38/38	0/3/3/3
2	GTP	F	201	3	-	2/18/38/38	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	A	201	GTP	C5-C6	-3.30	1.40	1.47
2	D	201	GTP	C8-N7	-3.14	1.29	1.35
2	A	201	GTP	C5-C4	-3.06	1.35	1.43
2	В	201	GTP	C5-C6	-2.85	1.41	1.47
2	D	201	GTP	C5-C6	-2.62	1.42	1.47
2	F	201	GTP	C5-C6	-2.56	1.42	1.47
2	D	201	GTP	C5-C4	-2.50	1.36	1.43
2	В	201	GTP	C5-C4	-2.29	1.37	1.43
2	С	201	GTP	C8-N7	-2.26	1.31	1.35
2	В	201	GTP	C8-N7	-2.23	1.31	1.35
2	A	201	GTP	PA-O1A	-2.19	1.43	1.50
2	С	201	GTP	C5-C4	-2.14	1.37	1.43
2	A	201	GTP	PA-O5'	2.13	1.67	1.59
2	Е	201	GTP	C5-C6	-2.12	1.43	1.47

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	F	201	GTP	C5-C4	-2.12	1.37	1.43
2	Е	201	GTP	C5-C4	-2.11	1.37	1.43
2	A	201	GTP	C8-N7	-2.07	1.31	1.35
2	Е	201	GTP	PA-O1A	-2.02	1.43	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	201	GTP	C5'-O5'-PA-O3A
2	A	201	GTP	PA-O3A-PB-O2B
2	В	201	GTP	PA-O3A-PB-O2B
2	С	201	GTP	PA-O3A-PB-O2B
2	D	201	GTP	PA-O3A-PB-O2B
2	Е	201	GTP	PA-O3A-PB-O2B
2	F	201	GTP	PA-O3A-PB-O2B


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

