

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 06:11 AM EDT

PDB ID	:	6MO1
Title	:	Structure of dengue virus protease with an allosteric Inhibitor that blocks
		replication
Authors	:	Lin, YL.; Hua, Y.; Nie, S.; Wu, J.; Wu, F.; Huo, T.; Yao, Y.; Song, Y.
Deposited on		
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 3.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	\mathbf{Res}	Chirality	Geometry	Clashes	Electron density
2	I16	А	1201	-	Х	-	-



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2578 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	٨	164	Total	С	Ν	0	S	0	0	0
	I A		1256	796	217	240	3			
1	Р	164	Total	С	Ν	0	S	0	0	0
	D		1256	796	217	240	3	0	0	0

• Molecule 1 is a protein called FLAVIVIRUS_NS2B/Peptidase S7.

Chain	Residue	Modelled	Actual	Comment	Reference
А	43	GLY	-	expression tag	UNP B3RFT3
А	44	SER	-	expression tag	UNP B3RFT3
А	45	HIS	-	expression tag	UNP B3RFT3
А	46	MET	-	expression tag	UNP B3RFT3
A	47	LEU	-	expression tag	UNP B3RFT3
А	48	MET	-	expression tag	UNP B3RFT3
А	992	GLY	-	linker	UNP B3RFT3
А	993	GLY	-	linker	UNP B3RFT3
А	994	GLY	-	linker	UNP B3RFT3
А	995	GLY	-	linker	UNP B3RFT3
А	996	SER	-	linker	UNP B3RFT3
А	997	GLY	-	linker	UNP B3RFT3
А	998	GLY	-	linker	UNP B3RFT3
А	999	GLY	-	linker	UNP B3RFT3
A	1000	GLY	-	linker	UNP B3RFT3
А	1167	ASN	GLN	conflict	UNP $Q91H74$
В	43	GLY	-	expression tag	UNP B3RFT3
В	44	SER	-	expression tag	UNP B3RFT3
В	45	HIS	-	expression tag	UNP B3RFT3
В	46	MET	-	expression tag	UNP B3RFT3
В	47	LEU	-	expression tag	UNP B3RFT3
В	48	MET	-	expression tag	UNP B3RFT3
В	992	GLY	-	linker	UNP B3RFT3
В	993	GLY	-	linker	UNP B3RFT3
В	994	GLY	-	linker	UNP B3RFT3

There are 32 discrepancies between the modelled and reference sequences:


Continued on next page...



Continu	eu jioni pre	vious puye			
Chain	Residue	Modelled	Actual	Comment	Reference
В	995	GLY	-	linker	UNP B3RFT3
В	996	SER	-	linker	UNP B3RFT3
В	997	GLY	-	linker	UNP B3RFT3
В	998	GLY	-	linker	UNP B3RFT3
В	999	GLY	-	linker	UNP B3RFT3
В	1000	GLY	-	linker	UNP B3RFT3
В	1167	ASN	GLN	conflict	UNP Q91H74

Continued from previous page...

• Molecule 2 is 5-[4-(aminomethyl)phenyl]-6-[4-(furan-3-yl)phenyl]-N-[(piperidin-4-yl)methyl] pyrazin-2-amine (three-letter code: I16) (formula: C₂₇H₂₉N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	А	1	Total C 33 2				0	0
2	В	1	Total C 33 2	C N 27 5		O 1	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants	109.33Å 109.33Å 76.25Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.38 - 3.00	Depositor
% Data completeness	99.9 (47.38-3.00)	Depositor
(in resolution range)	· · · · ·	-
R _{merge}	0.45	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.87 (at 3.01 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.238 , 0.312	Depositor
Wilson B-factor $(Å^2)$	56.5	Xtriage
Anisotropy	0.165	Xtriage
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtriage
Total number of atoms	2578	wwPDB-VP
Average B, all atoms $(Å^2)$	63.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



лмод	TAM52 is the root-mean-square of an 2 scores of the bolid lengths (of angles).											
Mol	Type	Chain	Res	Link	B	ond leng	gths	Bond angles				
MOI	Moi Type Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2			
2	I16	А	1201	-	34,37,37	3.91	24 (70%)	45,50,50	2.37	20 (44%)		
2	I16	В	1201	-	34,37,37	3.90	23 (67%)	45,50,50	2.48	18 (40%)		

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	I16	А	1201	-	-	5/19/27/27	0/5/5/5
2	I16	В	1201	-	-	6/19/27/27	0/5/5/5

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	В	1201	I16	C30-C29	13.87	1.53	1.37
2	А	1201	I16	C30-C29	10.41	1.49	1.37
2	А	1201	I16	C10-N11	8.71	1.48	1.34
2	А	1201	I16	C23-C10	7.32	1.57	1.49
2	В	1201	I16	C27-C28	7.08	1.51	1.38

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
2	В	1201	I16	C8-C7-C6	-6.41	111.91	121.13
2	В	1201	I16	C5-C4-C3	-4.89	114.30	121.03
2	В	1201	I16	C5-C6-C9	-4.80	113.00	120.61
2	А	1201	I16	C14-N13-C12	-4.71	114.76	123.36
2	А	1201	I16	C23-C10-N11	4.67	121.29	115.09

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	1201	I16	N13-C14-C15-C16
2	В	1201	I16	N13-C14-C15-C16
2	А	1201	I16	C9-C10-C23-C24

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	В	1201	I16	C9-C10-C23-C24
2	А	1201	I16	C9-C10-C23-C28

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

