



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 08:42 AM EDT

PDB ID : 6MOJ
Title : Dimeric DARPin A_angle_R5 complex with EpoR
Authors : Jude, K.M.; Mohan, K.; Garcia, K.C.
Deposited on : 2018-10-04
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TAR	A	301	-	X	-	-
3	TAR	A	302	-	X	-	-
3	TAR	B	301	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dimeric DARPin ACR5 (A_angle_R5).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	223	1599	1000	278	314	7	0	0	0


- Molecule 2 is a protein called Erythropoietin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	219	1697	1076	299	315	7	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	PHE	-	expression tag	UNP P19235
B	4	ALA	-	expression tag	UNP P19235
B	5	GLY	-	expression tag	UNP P19235
B	6	SER	-	expression tag	UNP P19235
B	7	ALA	-	expression tag	UNP P19235
B	52	GLN	ASN	conflict	UNP P19235
B	164	GLN	ASN	conflict	UNP P19235
B	226	LYS	-	expression tag	UNP P19235
B	227	GLU	-	expression tag	UNP P19235
B	228	LYS	-	expression tag	UNP P19235
B	229	ALA	-	expression tag	UNP P19235
B	230	ALA	-	expression tag	UNP P19235
B	231	ALA	-	expression tag	UNP P19235

- Molecule 3 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		
3	A	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	9	Total	O	0	0
			9	9		
5	B	13	Total	O	0	0
			13	13		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	130.42Å 130.42Å 293.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.11 – 2.43	Depositor
% Data completeness (in resolution range)	52.2 (46.11-2.43)	Depositor
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.42Å)	Xtriage
Refinement program	PHENIX (1.14_3211: ???)	Depositor
R, R_{free}	0.210 , 0.250	Depositor
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.007	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3366	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TAR	B	301	-	9,9,9	1.01	0	12,12,12	1.36	2 (16%)
3	TAR	A	302	-	9,9,9	1.04	0	12,12,12	1.41	3 (25%)
3	TAR	A	301	-	9,9,9	1.00	0	12,12,12	1.37	2 (16%)
4	GOL	B	304	-	5,5,5	0.84	0	5,5,5	0.97	0
4	GOL	B	303	-	5,5,5	0.96	0	5,5,5	0.98	0
4	GOL	B	302	-	5,5,5	0.93	0	5,5,5	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TAR	B	301	-	-	12/12/12/12	-
3	TAR	A	302	-	-	12/12/12/12	-
3	TAR	A	301	-	-	12/12/12/12	-
4	GOL	B	304	-	-	2/4/4/4	-
4	GOL	B	303	-	-	0/4/4/4	-
4	GOL	B	302	-	-	1/4/4/4	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	302	TAR	O11-C1-C2	2.84	120.94	113.27
3	B	301	TAR	O41-C4-C3	2.62	120.36	113.27
3	A	301	TAR	O41-C4-C3	2.62	120.35	113.27
3	B	301	TAR	O11-C1-C2	2.54	120.13	113.27
3	A	301	TAR	O11-C1-C2	2.44	119.88	113.27
3	A	302	TAR	O41-C4-C3	2.34	119.59	113.27
3	A	302	TAR	O11-C1-O1	-2.01	119.53	124.09

There are no chirality outliers.

All (39) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	TAR	O3-C3-C4-O4
3	A	301	TAR	O3-C3-C4-O41
3	B	301	TAR	O11-C1-C2-C3
3	B	301	TAR	C2-C3-C4-O4
3	B	301	TAR	C2-C3-C4-O41
3	B	301	TAR	O3-C3-C4-O4
4	B	304	GOL	C1-C2-C3-O3
3	A	302	TAR	O3-C3-C4-O4
3	A	302	TAR	O3-C3-C4-O41
3	B	301	TAR	O3-C3-C4-O41
3	A	301	TAR	O1-C1-C2-C3
3	A	301	TAR	O11-C1-C2-C3
3	A	301	TAR	C2-C3-C4-O4
3	A	301	TAR	C2-C3-C4-O41
3	A	302	TAR	O11-C1-C2-C3
3	A	302	TAR	C2-C3-C4-O4
3	A	302	TAR	C2-C3-C4-O41
3	B	301	TAR	O1-C1-C2-C3
3	A	302	TAR	C1-C2-C3-O3
3	A	302	TAR	O2-C2-C3-C4
3	A	302	TAR	O1-C1-C2-O2
3	B	301	TAR	O1-C1-C2-O2
3	A	302	TAR	O1-C1-C2-C3
3	A	302	TAR	O2-C2-C3-O3
3	A	301	TAR	O1-C1-C2-O2
3	A	301	TAR	O11-C1-C2-O2
3	A	302	TAR	O11-C1-C2-O2
3	B	301	TAR	O11-C1-C2-O2
3	A	302	TAR	C1-C2-C3-C4
3	B	301	TAR	C1-C2-C3-O3
3	B	301	TAR	O2-C2-C3-C4
3	B	301	TAR	C1-C2-C3-C4
4	B	302	GOL	O2-C2-C3-O3
3	A	301	TAR	C1-C2-C3-O3
3	A	301	TAR	O2-C2-C3-C4
3	B	301	TAR	O2-C2-C3-O3
4	B	304	GOL	O2-C2-C3-O3
3	A	301	TAR	O2-C2-C3-O3
3	A	301	TAR	C1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.