



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 03:20 PM EDT

PDB ID : 4MP4
Title : Crystal structure of a glutathione transferase family member from *Acinetobacter baumannii*, Target EFI-501785, apo structure
Authors : Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.F.; Morisco, L.L.; Wasserman, S.R.; Sojitra, S.; Stead, M.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Gerlt, J.A.; Armstrong, R.N.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2013-09-12
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 82%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">82% 10% 9%</p>
1	B	217	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 74%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">74% 14% 11%</p>
1	C	217	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">75% 14% 11%</p>

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	Total 1630	C 1061	N 268	O 300	S 1	0	1	0
1	B	194	Total 1590	C 1033	N 263	O 293	S 1	0	0	0
1	C	193	Total 1570	C 1020	N 261	O 288	S 1	0	0	0

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP B2HYC1
A	1	VAL	-	expression tag	UNP B2HYC1
A	210	ALA	-	expression tag	UNP B2HYC1
A	211	GLU	-	expression tag	UNP B2HYC1
A	212	ASN	-	expression tag	UNP B2HYC1
A	213	LEU	-	expression tag	UNP B2HYC1
A	214	TYR	-	expression tag	UNP B2HYC1
A	215	PHE	-	expression tag	UNP B2HYC1
A	216	GLN	-	expression tag	UNP B2HYC1
B	0	MET	-	expression tag	UNP B2HYC1
B	1	VAL	-	expression tag	UNP B2HYC1
B	210	ALA	-	expression tag	UNP B2HYC1
B	211	GLU	-	expression tag	UNP B2HYC1
B	212	ASN	-	expression tag	UNP B2HYC1
B	213	LEU	-	expression tag	UNP B2HYC1
B	214	TYR	-	expression tag	UNP B2HYC1
B	215	PHE	-	expression tag	UNP B2HYC1
B	216	GLN	-	expression tag	UNP B2HYC1
C	0	MET	-	expression tag	UNP B2HYC1
C	1	VAL	-	expression tag	UNP B2HYC1
C	210	ALA	-	expression tag	UNP B2HYC1
C	211	GLU	-	expression tag	UNP B2HYC1
C	212	ASN	-	expression tag	UNP B2HYC1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	213	LEU	-	expression tag	UNP B2HYC1
C	214	TYR	-	expression tag	UNP B2HYC1
C	215	PHE	-	expression tag	UNP B2HYC1
C	216	GLN	-	expression tag	UNP B2HYC1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	81	82	82	0	1
3	B	29	29	29	0	0
3	C	51	51	51	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutathione S-transferase



- Molecule 1: Glutathione S-transferase



- Molecule 1: Glutathione S-transferase



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.20Å 103.36Å 51.06Å 90.00° 91.09° 90.00°	Depositor
Resolution (Å)	36.32 – 2.50 36.32 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.3 (36.32-2.50) 96.3 (36.32-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.191 , 0.255 0.196 , 0.256	Depositor DCC
R_{free} test set	1608 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.000 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.005 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.007 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4957	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1669	0.53	0/2262
1	B	0.38	0/1628	0.53	0/2205
1	C	0.41	0/1606	0.54	0/2176
All	All	0.40	0/4903	0.53	0/6643

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1630	0	1624	13	0
1	B	1590	0	1582	17	0
1	C	1570	0	1580	16	0
2	A	5	0	0	1	0
3	A	82	0	0	2	0
3	B	29	0	0	0	0
3	C	51	0	0	0	0
All	All	4957	0	4786	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ASP:OD2	1:A:142:LYS:NZ	2.17	0.74
1:C:8:LYS:HE2	1:C:37:ALA:HB2	1.79	0.65
2:A:301:SO4:O1	1:C:208:HIS:NE2	2.31	0.64
1:B:13:SER:HA	1:B:38:ALA:HB1	1.83	0.60
1:B:36:ILE:HD13	1:B:206:LEU:HD21	1.84	0.59
1:A:5:GLU:N	3:A:425:HOH:O	2.36	0.58
1:C:86:ASP:N	1:C:86:ASP:OD1	2.38	0.56
1:C:122:VAL:HA	1:C:126:GLN:O	2.05	0.56
1:B:131:PRO:O	1:B:132:LEU:HB2	2.05	0.56
1:C:99:ARG:NH1	1:C:102:GLN:OE1	2.39	0.55
1:B:14:GLN:OE1	1:B:14:GLN:N	2.42	0.53
1:C:92:LYS:O	1:C:97:ARG:NH2	2.43	0.52
1:C:13:SER:HA	1:C:38:ALA:HB1	1.92	0.52
1:B:213:LEU:HD13	1:B:214:TYR:N	2.24	0.51
1:B:29:GLU:HG3	1:B:162:TRP:CH2	2.45	0.51
1:B:151:LEU:HB3	1:B:186:ARG:NH2	2.26	0.50
1:A:57:THR:HG21	1:A:62:VAL:CG2	2.42	0.50
3:A:480:HOH:O	1:C:188:LYS:HE3	2.11	0.49
1:A:8:LYS:NZ	1:A:35:GLU:OE1	2.45	0.49
1:B:132:LEU:HB3	1:B:137:LYS:NZ	2.26	0.49
1:A:29:GLU:HG3	1:A:162:TRP:CH2	2.49	0.48
1:A:8:LYS:HD3	1:A:10:TYR:CZ	2.48	0.48
1:B:14:GLN:HG2	1:B:16:LEU:HG	1.95	0.48
1:C:151:LEU:HD13	1:C:186:ARG:NH2	2.30	0.46
1:C:29:GLU:HG3	1:C:162:TRP:CH2	2.51	0.46
1:A:110:ASP:CG	1:A:111:LEU:H	2.19	0.46
1:A:13:SER:HA	1:A:38:ALA:HB1	1.98	0.45
1:B:110:ASP:OD1	1:B:111:LEU:N	2.49	0.44
1:A:102:GLN:OE1	1:B:70:LEU:HD12	2.18	0.44
1:B:39:ILE:HD12	1:B:39:ILE:HA	1.84	0.43
1:C:203:TRP:O	1:C:207:ARG:HG3	2.18	0.43
1:B:130:THR:HA	1:B:131:PRO:HD2	1.84	0.43
1:C:183:VAL:HB	1:C:188:LYS:HZ3	1.84	0.42
1:B:149:LYS:HD3	1:B:149:LYS:HA	1.91	0.42
1:C:183:VAL:HB	1:C:188:LYS:NZ	2.34	0.42
1:A:97:ARG:NH1	1:B:82:GLU:OE1	2.45	0.42
1:A:57:THR:HG21	1:A:62:VAL:HG22	2.00	0.42
1:A:205:ALA:HB2	1:C:208:HIS:HB2	2.02	0.41
1:B:207:ARG:CZ	1:B:207:ARG:HB3	2.50	0.41
1:C:152:ALA:HA	1:C:153:SER:HA	1.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:LEU:HB3	1:A:186:ARG:NH2	2.36	0.41
1:C:130:THR:HA	1:C:131:PRO:HD3	1.91	0.41
1:B:11:THR:HG22	1:B:20:ALA:HB1	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/217 (90%)	189 (97%)	5 (3%)	1 (0%)	29	48
1	B	190/217 (88%)	175 (92%)	11 (6%)	4 (2%)	7	11
1	C	189/217 (87%)	180 (95%)	8 (4%)	1 (0%)	29	48
All	All	574/651 (88%)	544 (95%)	24 (4%)	6 (1%)	15	28

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	132	LEU
1	A	72	GLU
1	B	131	PRO
1	C	72	GLU
1	B	72	GLU
1	B	125	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/188 (92%)	170 (99%)	2 (1%)	71	88
1	B	168/188 (89%)	160 (95%)	8 (5%)	25	48
1	C	167/188 (89%)	162 (97%)	5 (3%)	41	68
All	All	507/564 (90%)	492 (97%)	15 (3%)	41	68

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ASP
1	A	148	GLU
1	B	31	GLN
1	B	121	ASP
1	B	184	SER
1	B	192	LEU
1	B	197	ARG
1	B	209	LYS
1	B	212	ASN
1	B	213	LEU
1	C	31	GLN
1	C	109	SER
1	C	110	ASP
1	C	129	SER
1	C	195	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	B	31	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	301	-	4,4,4	0.15	0	6,6,6	0.37	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/217 (91%)	-0.24	3 (1%) 73 75	12, 26, 52, 77	0
1	B	194/217 (89%)	0.21	15 (7%) 13 13	15, 41, 80, 97	0
1	C	193/217 (88%)	-0.05	5 (2%) 56 59	14, 29, 64, 84	0
All	All	585/651 (89%)	-0.03	23 (3%) 39 42	12, 32, 68, 97	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	41	LEU	6.3
1	C	40	ASP	4.3
1	B	128	LYS	3.5
1	B	134	GLU	3.2
1	B	138	LYS	3.1
1	B	135	GLU	3.1
1	A	153	SER	2.9
1	C	134	GLU	2.9
1	B	131	PRO	2.8
1	B	6	TYR	2.8
1	B	137	LYS	2.8
1	B	5	GLU	2.4
1	B	215	PHE	2.3
1	A	134	GLU	2.3
1	B	113	ALA	2.2
1	B	153	SER	2.2
1	C	6	TYR	2.2
1	B	66	ASN	2.2
1	C	58	ALA	2.1
1	A	155	ALA	2.1
1	B	141	GLU	2.1
1	B	133	SER	2.1
1	B	126	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	301	5/5	0.92	0.24	44,51,59,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.