

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 05:43 PM EDT

PDB ID : 2MQJ BMRB ID : 25036

Title : Solution structure of ubiquitin-like protein from Caldiarchaeum subterraneum

Authors : Ye, K. Deposited on : 2014-06-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 70%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m NMR~archive} \ (\#{ m Entries})$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Qua	ality of chain		
1	A	88	43%	36%	8%	13%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model		
1	A:10-A:16, A:20-A:82 (70)	0.53	19		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20
2	3, 11



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1177 atoms, of which 612 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Ubiquitin-like protein.

Mol	Chain	Residues	Atoms			Trace			
1	Λ	77	Total	С	Н	N	О	S	0
1	A	11	1177	358	612	98	106	3	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	HIS	-	expression tag	UNP E6N8B8




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Ubiquitin-like protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

• Molecule 1: Ubiquitin-like protein





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: distance geometry.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Sparky	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	796
Number of shifts mapped to atoms	727
Number of unparsed shifts	0
Number of shifts with mapping errors	69
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	70%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	524	569	569	17±4
All	All	10480	11380	11380	336

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

5 of 165 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:13:ILE:HD11	1:A:40:VAL:HG11	1.01	1.30	11	1
1:A:13:ILE:HD13	1:A:40:VAL:HG11	0.95	1.38	7	4
1:A:11:ILE:HD12	1:A:13:ILE:HD11	0.94	1.40	1	2
1:A:60:LEU:HD22	1:A:66:LEU:HD21	0.86	1.48	3	2
1:A:25:LEU:HD13	1:A:40:VAL:HG13	0.85	1.47	2	2

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	70/88 (80%)	63±1 (90±2%)	5±1 (7±2%)	2±1 (3±1%)	7 40
All	All	1400/1760 (80%)	1265 (90%)	93 (7%)	42 (3%)	7 40

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	64	GLU	20
1	A	61	LYS	15
1	A	20	GLY	7

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	nalysed Rotameric		Percentiles		
1	A	58/69 (84%)	42±3 (72±5%)	16±3 (28±5%)	2 20		
All	All	1160/1380 (84%)	840 (72%)	320 (28%)	2 20		

5 of 37 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	32	THR	19
1	A	39	LYS	18
1	A	43	MET	18
1	A	56	LYS	16
1	A	54	THR	14

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 70% for the well-defined parts and 71% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	796
Number of shifts mapped to atoms	727
Number of unparsed shifts	0
Number of shifts with mapping errors	69
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 69) occurrences are reported below.

T:-4 ID	Cl :	D	Т	A 4		Shift Data	l
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	8	HIS	HA	4.797	0.02	1
1	A	8	HIS	HB2	3.006	0.02	2
1	A	8	HIS	HB3	3.006	0.02	2
1	A	8	HIS	CA	55.345	5.00	1
1	A	8	HIS	СВ	31.142	5.00	1
1	A	86	GLY	Н	8.494	0.02	1
1	A	86	GLY	HA2	4.011	0.02	2
1	A	86	GLY	HA3	4.011	0.02	2
1	A	86	GLY	CA	45.238	5.00	1
1	A	86	GLY	N	108.268	0.20	1
1	A	87	CYS	Н	8.482	0.02	1
1	A	87	CYS	HA	4.559	0.02	1
1	A	87	CYS	HB2	2.965	0.02	2
1	A	87	CYS	HB3	2.965	0.02	2

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from previo				A .	Shift Data				
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity		
1	A	87	CYS	CA	58.491	5.00	1		
1	A	87	CYS	СВ	28.229	5.00	1		
1	A	87	CYS	N	118.687	0.20	1		
1	A	88	GLY	Н	8.594	0.02	1		
1	A	88	GLY	HA2	3.933	0.02	2		
1	A	88	GLY	HA3	3.933	0.02	2		
1	A	88	GLY	CA	45.092	5.00	1		
1	A	88	GLY	N	111.528	0.20	1		
1	A	89	GLU	Н	7.974	0.02	1		
1	A	89	GLU	CA	54.258	5.00	1		
1	A	89	GLU	СВ	30.604	5.00	1		
1	A	89	GLU	N	119.482	0.20	1		
1	A	90	PRO	HA	4.439	0.02	1		
1	A	90	PRO	HB2	2.282	0.02	2		
1	A	90	PRO	HB3	2.282	0.02	2		
1	A	90	PRO	HG3	1.828	0.02	2		
1	A	90	PRO	HD2	3.794	0.02	2		
1	A	90	PRO	HD3	3.677	0.02	2		
1	A	90	PRO	CA	62.733	5.00	1		
1	A	90	PRO	СВ	31.958	5.00	1		
1	A	90	PRO	CG	27.189	5.00	1		
1	A	90	PRO	CD	50.192	5.00	1		
1	A	91	ILE	Н	8.298	0.02	1		
1	A	91	ILE	HA	4.072	0.02	1		
1	A	91	ILE	НВ	1.792	0.02	1		
1	A	91	ILE	CA	61.101	5.00	1		
1	A	91	ILE	СВ	38.38	5.00	1		
1	A	91	ILE	CG2	16.997	5.00	1		
1	A	91	ILE	CD1	12.049	5.00	1		
1	A	91	ILE	N	121.704	0.20	1		
1	A	92	ARG	Н	8.492	0.02	1		
1	A	92	ARG	HA	4.368	0.02	1		
1	A	92	ARG	HB2	1.78	0.02	2		
1	A	92	ARG	HB3	1.78	0.02	2		
1	A	92	ARG	CA	55.676	5.00	1		
1	A	92	ARG	СВ	30.855	5.00	1		
1	A	92	ARG	CG	26.894	5.00	1		
1	A	92	ARG	CD	43.126	5.00	1		
1	A	92	ARG	N	126.325	0.20	1		
1	A	93	ARG	Н	8.496	0.02	1		
1	A	93	ARG	HA	4.33	0.02	1		

Continued on next page...



Continued from previous page...

List ID	Chain	Peg	Trme	Atom		ı	
LIST ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	93	ARG	CA	55.738	5.00	1
1	A	93	ARG	СВ	30.972	5.00	1
1	A	93	ARG	CG	26.839	5.00	1
1	A	93	ARG	CD	43.141	5.00	1
1	A	93	ARG	N	123.839	0.20	1
1	A	94	ALA	Н	8.479	0.02	1
1	A	94	ALA	HA	4.304	0.02	1
1	A	94	ALA	CA	52.15	5.00	1
1	A	94	ALA	СВ	19.168	5.00	1
1	A	94	ALA	N	126.719	0.20	1
1	A	95	ALA	Н	8.008	0.02	1
1	A	95	ALA	CA	53.841	5.00	1
1	A	95	ALA	CB	19.842	5.00	1
1	A	95	ALA	N	129.249	0.20	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	85	0.08 ± 0.32	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	76	0.02 ± 0.11	None needed (< 0.5 ppm)
¹³ C′	0		None (insufficient data)
^{15}N	80	0.91 ± 0.51	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 70%, i.e. 669 atoms were assigned a chemical shift out of a possible 950. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	$265/345 \ (77\%)$	133/140 (95%)	68/140 (49%)	64/65 (98%)
Sidechain	404/586~(69%)	270/384 (70%)	134/181 (74%)	0/21 (0%)
Aromatic	0/19 (0%)	0/9 (0%)	0/10 (0%)	0/0 (%)
Overall	669/950 (70%)	403/533 (76%)	202/331 (61%)	64/86 (74%)




7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	736
Intra-residue ($ i-j =0$)	139
Sequential ($ i-j =1$)	288
Medium range ($ i-j >1$ and $ i-j <5$)	125
Long range (i-j ≥5)	154
Inter-chain	0
Hydrogen bond restraints	30
Disulfide bond restraints	0
Total dihedral-angle restraints	112
Number of unmapped restraints	0
Number of restraints per residue	9.6
Number of long range restraints per residue ¹	2.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	4.3	0.2
0.2-0.5 (Medium)	0.9	0.46
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	$\operatorname{Max}(^{\circ})$
1.0-10.0 (Small)	14.6	5.7
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dordensinda dom o	Count	$rac{1}{1}$	Vic	${f Violated}^3$			Consistently Violated ⁴		
Restraints type	Count	901	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$	
Intra-residue (i-j =0)	139	18.9	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	51	6.9	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	88	12.0	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Sequential (i-j =1)	288	39.1	5	1.7	0.7	1	0.3	0.1	
Backbone-Backbone	158	21.5	2	1.3	0.3	1	0.6	0.1	
Backbone-Sidechain	130	17.7	3	2.3	0.4	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Medium range ($ i-j >1 \& i-j <5$)	125	17.0	6	4.8	0.8	1	0.8	0.1	
Backbone-Backbone	93	12.6	4	4.3	0.5	0	0.0	0.0	
Backbone-Sidechain	32	4.3	2	6.2	0.3	1	3.1	0.1	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Long range ($ i-j \ge 5$)	154	20.9	2	1.3	0.3	0	0.0	0.0	
Backbone-Backbone	116	15.8	1	0.9	0.1	0	0.0	0.0	
Backbone-Sidechain	38	5.2	1	2.6	0.1	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0	
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	
Hydrogen bond	30	4.1	3	10.0	0.4	0	0.0	0.0	
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0	
Total	736	100.0	16	2.2	2.2	2	0.3	0.3	
Backbone-Backbone	448	60.9	10	2.2	1.4	1	0.2	0.1	
Backbone-Sidechain	288	39.1	6	2.1	0.8	1	0.3	0.1	
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0	

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	3	Magn (Å)	Morr (Å)	\mathbf{SD}^6 (Å)	Modian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	$SD^*(A)$	Median (Å)
1	0	2	5	1	0	8	0.15	0.3	0.06	0.12
2	0	3	1	0	0	4	0.14	0.17	0.02	0.14
3	0	2	2	1	0	5	0.13	0.18	0.03	0.12
4	0	2	1	0	0	3	0.16	0.18	0.01	0.16
5	0	1	5	0	0	6	0.17	0.33	0.08	0.14
6	0	2	2	0	0	4	0.22	0.43	0.12	0.18
7	0	2	4	0	0	6	0.14	0.16	0.02	0.14
8	0	3	3	0	0	6	0.19	0.23	0.03	0.18
9	0	2	4	0	0	6	0.17	0.35	0.08	0.14
10	0	3	2	0	0	5	0.17	0.24	0.04	0.16
11	0	1	3	0	0	4	0.2	0.29	0.05	0.18

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	0	2	3	0	0	5	0.21	0.46	0.13	0.15
13	0	2	2	1	0	5	0.13	0.19	0.03	0.11
14	0	2	2	0	0	4	0.19	0.3	0.07	0.17
15	0	2	5	0	0	7	0.17	0.39	0.09	0.14
16	0	2	2	0	0	4	0.19	0.22	0.03	0.19
17	0	3	3	0	0	6	0.21	0.37	0.08	0.18
18	0	2	3	1	0	6	0.16	0.25	0.05	0.14
19	0	3	3	0	0	6	0.19	0.32	0.07	0.17
20	0	3	3	0	0	6	0.14	0.19	0.03	0.12

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 693(IR:139, SQ:283, MR:119, LR:152, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	1	1	1	0	3	1	5.0
0	0	0	1	0	1	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	1	1	0	0	2	5	25.0
0	0	1	0	0	1	6	30.0
0	0	0	0	0	0	7	35.0
0	1	1	0	0	2	8	40.0
0	0	0	0	0	0	9	45.0
0	1	0	0	0	1	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	1	0	0	1	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	1	1	0	0	2	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,164)	1:A:32:THR:H	1:A:29:PRO:HB3	20	0.19	0.01	0.18
(1,209)	1:A:36:VAL:H	1:A:35:ALA:HA	20	0.15	0.0	0.15
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	15	0.3	0.09	0.3
(1,337)	1:A:45:LYS:H	1:A:44:LYS:HG2	10	0.14	0.03	0.14
(1,337)	1:A:45:LYS:H	1:A:44:LYS:HG3	10	0.14	0.03	0.14
(1,234)	1:A:37:ARG:H	1:A:33:VAL:H	8	0.12	0.01	0.12
(1,335)	1:A:45:LYS:H	1:A:44:LYS:HB3	8	0.11	0.0	0.11
(1,198)	1:A:35:ALA:H	1:A:32:THR:HA	6	0.11	0.01	0.11
(1,334)	1:A:45:LYS:H	1:A:44:LYS:HB2	5	0.23	0.01	0.24
(1,80)	1:A:17:VAL:H	1:A:21:SER:H	5	0.13	0.02	0.13
(1,521)	1:A:63:THR:H	1:A:32:THR:HA	2	0.11	0.0	0.11


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	12	0.46
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	6	0.43
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	15	0.39
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	17	0.37
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	9	0.35
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	5	0.33
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	19	0.32
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	1	0.3
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	14	0.3
(1,253)	1:A:39:LYS:H	1:A:36:VAL:HG21	11	0.29



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	Count	$\%^{1}$	Vie	olated	3	Consis	tent	${ m ly~Violated^4}$		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	y Violated ⁴ $\frac{\%^1}{4.5}$ $\frac{3.6}{8.0}$		
PSI	58	51.8	20	34.5	17.9	5	8.6	4.5		
PHI	54	48.2	16	29.6	14.3	4	7.4	3.6		
Total	112	100.0	36	32.1	32.1	9	8.0	8.0		

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	f violations	Moon (°)	Mov (°)	SD (°)	Modian (°)
Wiodei 1D	PSI	PHI	Total	$ \mathbf{Mean} (^{\circ}) $	$\mathbf{Max} (^{\circ})$	$\mathbf{SD} \ (^{\circ})$	\mid Median (°) \mid
1	7	8	15	2.39	3.9	0.92	2.3
2	7	7	14	2.47	4.5	0.99	2.4
3	8	5	13	2.78	4.3	0.77	2.8
4	8	8	16	2.26	4.6	1.11	1.9
5	11	8	19	2.28	4.1	0.9	2.2
6	9	6	15	2.57	5.7	1.18	2.0
7	7	6	13	2.54	3.9	0.95	2.3
8	8	8	16	2.39	4.3	1.0	2.15
9	7	6	13	2.68	4.2	0.92	2.8
10	9	5	14	2.86	5.2	1.26	2.45
11	7	5	12	2.97	5.1	1.19	3.2
12	10	6	16	2.56	4.7	1.07	2.45
13	12	8	20	2.32	5.0	1.09	2.05
14	6	5	11	3.07	5.7	1.06	2.8
15	8	7	15	2.6	4.4	0.93	2.9
16	8	8	16	2.29	4.5	0.97	2.2
17	8	5	13	3.05	5.6	1.35	3.1
18	9	7	16	2.77	4.9	1.26	2.6
19	8	5	13	2.7	4.8	0.99	2.5
20	7	5	12	2.94	4.7	1.25	2.7



10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count ¹	%
5	2	7	1	5.0
3	5	8	2	10.0
3	1	4	3	15.0
0	1	1	4	20.0
1	1	2	5	25.0
0	1	1	6	30.0
0	0	0	7	35.0
1	0	1	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0

Continued on next page...




Continued from previous page...

Nun	nber o	of violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count ¹	%
1	0	1	12	60.0
0	0	0	13	65.0
0	0	0	14	70.0
0	0	0	15	75.0
0	0	0	16	80.0
0	0	0	17	85.0
0	1	1	18	90.0
1	0	1	19	95.0
5	4	9	20	100.0

¹ Number of models with violations

10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	20	4.61	0.65	4.55
(1,30)	1:A:31:ALA:C	1:A:32:THR:N	1:A:32:THR:CA	1:A:32:THR:C	20	4.28	0.39	4.2
(1,31)	1:A:32:THR:N	1:A:32:THR:CA	1:A:32:THR:C	1:A:33:VAL:N	20	3.55	0.41	3.45
(1,32)	1:A:32:THR:C	1:A:33:VAL:N	1:A:33:VAL:CA	1:A:33:VAL:C	20	3.39	0.47	3.4
(1,28)	1:A:29:PRO:C	1:A:30:ASN:N	1:A:30:ASN:CA	1:A:30:ASN:C	20	2.99	0.28	2.95
(1,23)	1:A:26:GLU:C	1:A:27:VAL:N	1:A:27:VAL:CA	1:A:27:VAL:C	20	2.72	0.5	2.65
(1,22)	1:A:26:GLU:N	1:A:26:GLU:CA	1:A:26:GLU:C	1:A:27:VAL:N	20	2.7	0.5	2.6
(1,84)	1:A:63:THR:N	1:A:63:THR:CA	1:A:63:THR:C	1:A:64:GLU:N	20	2.44	0.41	2.4
(1,27)	1:A:29:PRO:N	1:A:29:PRO:CA	1:A:29:PRO:C	1:A:30:ASN:N	20	2.36	0.3	2.25
(1,83)	1:A:62:ASP:N	1:A:62:ASP:CA	1:A:62:ASP:C	1:A:63:THR:N	19	1.68	0.37	1.7


 $^{^1}$ Number of violated models, $^2\mathrm{Standard}$ deviation, All angle values are in degree (°)



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	6	5.7
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	14	5.7
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	17	5.6
(1,30)	1:A:31:ALA:C	1:A:32:THR:N	1:A:32:THR:CA	1:A:32:THR:C	17	5.2
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	10	5.2
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	11	5.1
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	13	5.0
(1,30)	1:A:31:ALA:C	1:A:32:THR:N	1:A:32:THR:CA	1:A:32:THR:C	18	4.9
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	18	4.9
(1,26)	1:A:28:ALA:N	1:A:28:ALA:CA	1:A:28:ALA:C	1:A:29:PRO:N	19	4.8

