



## wwPDB EM Validation Summary Report ⓘ

Nov 19, 2022 – 02:12 pm GMT

PDB ID : 5MQF  
EMDB ID : EMD-3545  
Title : Cryo-EM structure of a human spliceosome activated for step 2 of splicing (C\* complex)  
Authors : Bertram, K.; Hartmuth, K.; Kastner, B.  
Deposited on : 2016-12-20  
Resolution : 5.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 5.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.


























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2335	72% 12% 16%
2	B	972	77% 15% 7%
3	C	536	40% 55%
4	D	515	45% 16% 39%
5	E	579	43% 14% 43%
6	F	357	65% 21% 14%
7	G	504	24% 74%






Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	H	504	
7	I	504	
7	J	504	
8	K	225	
9	L	802	
10	M	855	
11	N	243	
12	O	848	
13	P	420	
14	Q	144	
15	R	229	
16	S	2752	
17	T	908	
18	U	1485	
19	V	166	
20	W	255	
21	X	225	
22	a	118	
22	h	118	
23	b	86	
23	i	86	
24	c	92	
24	j	92	
25	d	76	
25	k	76	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
26	e	126	 67% 33%
26	l	126	 66% 34%
27	f	240	 30% 70%
27	m	240	 30% 70%
28	g	119	 68% 31%
28	n	119	 69% 31%
29	o	301	 74% 26%
30	p	411	 31% 94% 6%
31	q	1218	 51% 49%
32	Y	324	 6% 90%
32	Z	324	 96%
33	2	188	 47% 20% 7% 26%
34	5	116	 42% 45% 11%
35	6	106	 36% 38% 10% 16%

## 2 Entry composition [i](#)

There are 35 unique types of molecules in this entry. The entry contains 55236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-processing-splicing factor 8.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
1	A	1965	9955	6025	1965	1965	13	0

- Molecule 2 is a protein called 116 kDa U5 small nuclear ribonucleoprotein component.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	B	902	4566	2762	902	902	0	0

- Molecule 3 is a protein called SNW domain-containing protein 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
3	C	240	1231	751	240	240	0	0

- Molecule 4 is a protein called Pleiotropic regulator 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	D	315	1576	946	315	315	8	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	454	ALA	VAL	conflict	UNP O43660
D	455	VAL	GLN	conflict	UNP O43660
D	456	GLN	PRO	conflict	UNP O43660
D	457	PRO	GLY	conflict	UNP O43660
D	458	GLY	SER	conflict	UNP O43660
D	459	SER	LEU	conflict	UNP O43660
D	460	LEU	ASP	conflict	UNP O43660
D	461	ASP	SER	conflict	UNP O43660
D	492	ASP	-	insertion	UNP O43660

- Molecule 5 is a protein called Pre-mRNA-processing factor 17.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
5	E	329	1646	988	329	329	0	0

- Molecule 6 is a protein called U5 small nuclear ribonucleoprotein 40 kDa protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
6	F	307	1531	917	307	307	0	0

- Molecule 7 is a protein called Pre-mRNA-processing factor 19.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
7	G	132	679	415	132	132	0	0
7	H	135	696	426	135	135	0	0
7	I	134	691	423	134	134	0	0
7	J	135	696	426	135	135	0	0

- Molecule 8 is a protein called Pre-mRNA-splicing factor SPF27.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
8	K	191	960	578	191	191	0	0

- Molecule 9 is a protein called Cell division cycle 5-like protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
9	L	336	1680	1008	336	336	0	0

- Molecule 10 is a protein called Pre-mRNA-splicing factor SYF1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
10	M	706	3553	2141	706	706	0	0

- Molecule 11 is a protein called Pre-mRNA-splicing factor SYF2.

Mol	Chain	Residues	Atoms			AltConf	Trace	
11	N	115	Total	C	N	O	0	0
			579	349	115	115		

- Molecule 12 is a protein called Crooked neck-like protein 1.

Mol	Chain	Residues	Atoms			AltConf	Trace	
12	O	564	Total	C	N	O	0	0
			2842	1715	564	563		

- Molecule 13 is a protein called Pre-mRNA-splicing factor RBM22.

Mol	Chain	Residues	Atoms			AltConf	Trace	
13	P	277	Total	C	N	O	0	0
			1396	842	277	277		

- Molecule 14 is a protein called Protein BUD31 homolog.

Mol	Chain	Residues	Atoms			AltConf	Trace	
14	Q	142	Total	C	N	O	0	0
			713	429	142	142		

- Molecule 15 is a protein called Spliceosome-associated protein CWC15 homolog.

Mol	Chain	Residues	Atoms			AltConf	Trace	
15	R	98	Total	C	N	O	0	0
			493	297	98	98		

- Molecule 16 is a protein called Serine/arginine repetitive matrix protein 2.

Mol	Chain	Residues	Atoms			AltConf	Trace	
16	S	30	Total	C	N	O	0	0
			148	88	30	30		

- Molecule 17 is a protein called Pre-mRNA-splicing factor CWC22 homolog.

Mol	Chain	Residues	Atoms			AltConf	Trace	
17	T	395	Total	C	N	O	2	197
			1191	795	198	198		

- Molecule 18 is a protein called Intron-binding protein aquarius.

Mol	Chain	Residues	Atoms		AltConf	Trace
18	U	1287	Total	C	6	1287
			1287	1287		

- Molecule 19 is a protein called Peptidyl-prolyl cis-trans isomerase-like 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
19	V	155	Total	C	N	O	0	0
			774	464	155	155		

- Molecule 20 is a protein called U2 small nuclear ribonucleoprotein A'.

Mol	Chain	Residues	Atoms				AltConf	Trace
20	W	162	Total	C	N	O	0	0
			816	492	162	162		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
W	89	ASP	CYS	conflict	UNP P09661
W	119	CYS	SER	conflict	UNP P09661

- Molecule 21 is a protein called U2 small nuclear ribonucleoprotein B'.

Mol	Chain	Residues	Atoms				AltConf	Trace
21	X	94	Total	C	N	O	0	0
			470	282	94	94		

- Molecule 22 is a protein called Small nuclear ribonucleoprotein Sm D2.

Mol	Chain	Residues	Atoms				AltConf	Trace
22	a	99	Total	C	N	O	0	0
			497	299	99	99		
22	h	98	Total	C	N	O	0	0
			493	297	98	98		

- Molecule 23 is a protein called Small nuclear ribonucleoprotein F.

Mol	Chain	Residues	Atoms				AltConf	Trace
23	b	76	Total	C	N	O	0	0
			379	227	76	76		
23	i	74	Total	C	N	O	0	0
			369	221	74	74		



- Molecule 24 is a protein called Small nuclear ribonucleoprotein E.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
24	c	79	393	235	79	79	0	0
24	j	79	393	235	79	79	0	0

- Molecule 25 is a protein called Small nuclear ribonucleoprotein G.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
25	d	74	369	221	74	74	0	0
25	k	74	369	221	74	74	0	0

- Molecule 26 is a protein called Small nuclear ribonucleoprotein Sm D3.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
26	e	84	420	252	84	84	0	0
26	l	83	415	249	83	83	0	0

- Molecule 27 is a protein called Small nuclear ribonucleoprotein-associated proteins B and B'.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
27	f	71	355	213	71	71	0	0
27	m	71	355	213	71	71	0	0

- Molecule 28 is a protein called Small nuclear ribonucleoprotein Sm D1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
28	g	82	412	248	82	82	0	0
28	n	82	412	248	82	82	0	0

- Molecule 29 is a protein called Peptidyl-prolyl cis-trans isomerase E.

Mol	Chain	Residues	Atoms		AltConf	Trace
29	o	223	Total	C	0	223
			223	223		

- Molecule 30 is a protein called Eukaryotic initiation factor 4A-III.

Mol	Chain	Residues	Atoms		AltConf	Trace
30	p	385	Total	C	8	385
			385	385		

- Molecule 31 is a protein called ATP-dependent RNA helicase DHX8.

Mol	Chain	Residues	Atoms		AltConf	Trace
31	q	627	Total	C	5	627
			627	627		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
q	?	-	MET	deletion	UNP Q14562
q	?	-	ASP	deletion	UNP Q14562
q	948	GLY	ALA	conflict	UNP Q14562
q	949	SER	PRO	conflict	UNP Q14562

- Molecule 32 is a RNA chain called MINX pre-mRNA (intron).

Mol	Chain	Residues	Atoms					AltConf	Trace
32	Y	32	Total	C	N	O	P	0	0
			677	303	115	227	32		
32	Z	12	Total	C	N	O	P	0	0
			256	114	47	83	12		

- Molecule 33 is a RNA chain called Human gene for small nuclear RNA U2 (snRNA U2).

Mol	Chain	Residues	Atoms					AltConf	Trace
33	2	140	Total	C	N	O	P	0	0
			2968	1327	510	991	140		

- Molecule 34 is a RNA chain called Homo sapiens U5 A small nuclear RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	5	114	Total	C	N	O	P	0	0
			2397	1074	399	810	114		


- Molecule 35 is a RNA chain called Homo sapiens RNA, U6 small nuclear 1 (RNU6-1), small nuclear RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
35	6	89	1903	851	350	613	89	0	0

### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-processing-splicing factor 8





























• Molecule 18: Intron-binding protein aquarius



GLY  
ALA  
THR  
SER  
THR  
PRO  
GLU  
GLU  
ALA  
ILE  
PRO  
ALA  
LEU  
SER  
GLU  
THR  
THR  
PRO  
THR  
VAL  
VAL  
GLY  
VAL  
ALA  
ALA  
SER  
SER  
ALA  
PRO  
ALA  
GLU  
ALA  
ASN  
ASN  
THR  
PRO  
GLN

- Molecule 19: Peptidyl-prolyl cis-trans isomerase-like 1



MET  
ALA  
ALA  
ILE  
ASN  
PRO  
PRO  
ASP  
SER  
TRP  
GLN  
P11  
L16  
E17  
T18  
I22  
W29  
T35  
C36  
F39  
A40  
I97  
L98  
A99  
M100  
F113  
V114  
T115  
T127  
I128  
F129  
I160  
S165  
GLY

- Molecule 20: U2 small nuclear ribonucleoprotein A'



MET  
Y2  
T5  
R27  
P32  
I47  
D58  
P61  
A86  
S99  
K160  
K163  
GLY  
LYS  
ARG  
GLY  
ALA  
GLN  
LEU  
ALA  
LYS  
ARG  
ASP  
ILE  
ARG  
SER  
GLY  
PRO  
THR  
LYS  
PHE  
ASN  
PRO  
GLU  
GLY  
ALA  
GLU  
MET  
GLY  
LEU  
THR  
ASP  
THR  
VAL  
LYS  
LYS  
GLY  
SER  
SER

GLU  
ALA  
ILE  
LYS  
ASN  
ALA  
ILE  
ALA  
ASN  
SER  
THR  
LEU  
ALA  
GLU  
VAL  
GLU  
ARG  
LEU  
LYS  
GLY  
LEU  
LEU  
SER  
GLY  
GLN  
ILE  
PRO  
GLY  
ALA  
ARG  
ARG  
SER  
GLY  
PRO  
THR  
ASP  
GLY  
GLU  
GLU  
MET  
GLU  
THR  
ASP  
THR  
VAL  
LYS  
LYS  
GLY  
SER

- Molecule 21: U2 small nuclear ribonucleoprotein B''




MET  
ASP  
I6  
F75  
P76  
G99  
THR  
PHE  
ALA  
LYS  
GLY  
LYS  
LYS  
LYS  
LYS  
LYS  
LYS  
LEU  
SER  
MET  
LYS  
ALA  
PHE  
LYS  
VAL  
GLN  
PHE  
PRO  
GLY  
THR  
THR  
VAL  
ARG  
ASN  
LYS  
VAL  
VAL  
PRO  
GLY  
ARG  
GLY  
HIS  
ASP  
ILE  
ALA  
PHE  
VAL  
ASN  
GLU  
PHE  
GLN  
ASN  
ASN  
THR  
THR  
VAL  
VAL  
PRO  
ASP  
VAL  
ASP  
VAL  
TYR  
PRO

PRO  
ASN  
TYR  
ILE  
LEU  
PHE  
LEU  
ASN  
ASN  
PRO  
GLU  
GLU  
THR  
ASN  
GLU  
MET  
LYS  
SER  
MET  
PHE  
ASN  
GLN  
PHE  
PRO  
GLY  
LYS  
VAL  
ARG  
ASN  
VAL  
VAL  
PRO  
GLY  
ARG  
HIS  
ASP  
ILE  
ALA  
PHE  
VAL  
ASN  
GLU  
PHE  
GLN  
ASN  
ASN  
THR  
THR  
VAL  
VAL  
PRO  
ASP  
VAL  
ASP  
VAL  
TYR  
PRO


GLY  
PHE  
LYS  
ILE  
THR  
PRO  
SER  
HIS  
ALA  
HIS  
LYS  
MET  
GLU  
THR  
ILE  
THR  
TYR  
ALA  
LYS  
LYS

- Molecule 22: Small nuclear ribonucleoprotein Sm D2




MET  
SER  
LEU  
LEU  
ASN  
LYS  
PRO  
LYS  
SER  
SER  
GLU  
MET  
MET  
PRO  
GLU  
GLU  
GLU  
GLN  
LYS  
ARG  
E20  
K84  
K118


- Molecule 22: Small nuclear ribonucleoprotein Sm D2





MET  
SER  
LEU  
LEU  
ASN  
LYS  
PRO  
LYS  
SER  
SER  
GLU  
MET  
MET  
PRO  
GLU  
GLU  
GLU  
GLN  
LYS  
R19  
E20  
V77  
P78  
K79  
S80  
G81  
S87  
K88  
P89  
N112  
A116  
GLY  
LYS

- Molecule 23: Small nuclear ribonucleoprotein F








- Molecule 23: Small nuclear ribonucleoprotein F



- Molecule 24: Small nuclear ribonucleoprotein E



- Molecule 24: Small nuclear ribonucleoprotein E




- Molecule 25: Small nuclear ribonucleoprotein G



- Molecule 25: Small nuclear ribonucleoprotein G




- Molecule 26: Small nuclear ribonucleoprotein Sm D3






- Molecule 26: Small nuclear ribonucleoprotein Sm D3











- Molecule 27: Small nuclear ribonucleoprotein-associated proteins B and B'




- Molecule 27: Small nuclear ribonucleoprotein-associated proteins B and B'




- Molecule 28: Small nuclear ribonucleoprotein Sm D1



- Molecule 28: Small nuclear ribonucleoprotein Sm D1




- Molecule 29: Peptidyl-prolyl cis-trans isomerase E












- Molecule 34: Homo sapiens U5 A small nuclear RNA



- Molecule 35: Homo sapiens RNA, U6 small nuclear 1 (RNU6-1), small nuclear RNA



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	136534	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	2.1	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	0.316	Depositor
Minimum map value	-0.146	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.016	Depositor
Map size (Å)	572.4, 572.4, 572.4	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.59, 1.59, 1.59	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.34	5/10059 (0.0%)	0.46	2/14115 (0.0%)
2	B	0.24	0/4619	0.44	1/6475 (0.0%)
3	C	0.27	0/1249	0.52	0/1758
4	D	0.25	0/1587	0.55	1/2215 (0.0%)
5	E	0.25	0/1655	0.49	0/2309
6	F	0.25	0/1540	0.49	0/2148
7	G	0.23	0/688	0.38	0/969
7	H	0.26	0/706	0.40	0/995
7	I	0.23	0/701	0.38	0/988
7	J	0.25	0/706	0.39	0/995
8	K	0.23	0/963	0.36	0/1346
9	L	0.23	0/1683	0.40	0/2349
10	M	0.23	0/3575	0.43	0/5006
11	N	0.27	0/582	0.52	0/814
12	O	0.50	1/2858 (0.0%)	0.42	0/4007
13	P	0.24	0/1408	0.44	0/1969
14	Q	0.23	0/718	0.39	0/1003
15	R	0.22	0/493	0.40	0/688
16	S	0.28	0/149	0.52	0/205
17	T	0.22	0/999	0.39	0/1397
19	V	0.25	0/782	0.45	0/1088
20	W	0.64	0/821	1.48	5/1149 (0.4%)
21	X	0.61	0/472	1.26	0/658
22	a	0.23	0/500	0.47	0/698
22	h	0.62	0/496	1.00	1/693 (0.1%)
23	b	0.23	0/382	0.44	0/531
23	i	0.67	0/372	0.78	0/517
24	c	0.22	0/393	0.43	0/547
24	j	0.64	0/393	0.81	0/547
25	d	0.23	0/371	0.47	0/516
25	k	0.50	0/371	0.69	0/516
26	e	0.23	0/422	0.50	0/588
26	l	0.46	0/417	0.60	0/581
27	f	0.23	0/356	0.47	0/494

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
27	m	0.46	0/356	0.63	0/494
28	g	0.41	1/414 (0.2%)	0.48	0/578
28	n	0.55	0/414	0.71	0/578
32	Y	0.45	1/754 (0.1%)	0.83	0/1169
32	Z	0.16	0/285	0.71	0/442
33	2	0.15	0/3312	0.68	0/5151
34	5	0.16	0/2672	0.76	0/4154
35	6	0.19	0/2129	0.78	0/3314
All	All	0.32	8/53822 (0.0%)	0.57	10/76754 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	B	0	1
4	D	0	1
5	E	0	1
6	F	0	1
9	L	0	1
10	M	0	2
11	N	0	2
12	O	0	2
22	h	0	1
All	All	0	16

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	O	209	PRO	C-N	23.47	1.78	1.34
1	A	1134	TRP	C-N	15.33	1.63	1.34
1	A	323	LEU	C-N	10.94	1.55	1.34
1	A	369	GLU	C-N	9.57	1.52	1.34
1	A	174	PRO	C-N	9.19	1.51	1.34

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
22	h	77	VAL	C-N-CD	-17.59	81.91	120.60
2	B	349	PHE	C-N-CA	6.40	137.71	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
20	W	5	THR	N-CA-CB	-6.30	98.32	110.30
20	W	27	ARG	CB-CA-C	-5.85	98.70	110.40
1	A	1945[A]	VAL	C-N-CA	-5.79	107.23	121.70

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	172	GLU	Peptide
1	A	1852	LEU	Peptide
1	A	1899	VAL	Peptide
1	A	1946	ASN	Peptide
2	B	352	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9955	0	4850	175	0
2	B	4566	0	2269	89	0
3	C	1231	0	665	18	0
4	D	1576	0	774	78	0
5	E	1646	0	787	82	0
6	F	1531	0	747	71	0
7	G	679	0	356	7	0
7	H	696	0	367	13	0
7	I	691	0	365	10	0
7	J	696	0	367	11	0
8	K	960	0	461	14	0
9	L	1680	0	807	12	0
10	M	3553	0	1738	42	0
11	N	579	0	277	5	0
12	O	2842	0	1355	20	0
13	P	1396	0	684	11	0
14	Q	713	0	337	3	0
15	R	493	0	220	5	0
16	S	148	0	78	29	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	T	1191	0	457	15	0
18	U	1287	0	0	20	0
19	V	774	0	404	17	0
20	W	816	0	386	1	0
21	X	470	0	223	1	0
22	a	497	0	229	0	0
22	h	493	0	226	0	0
23	b	379	0	186	0	0
23	i	369	0	182	0	0
24	c	393	0	169	0	0
24	j	393	0	169	0	0
25	d	369	0	178	0	0
25	k	369	0	178	0	0
26	e	420	0	200	0	0
26	l	415	0	196	0	0
27	f	355	0	164	0	0
27	m	355	0	164	0	0
28	g	412	0	188	0	0
28	n	412	0	188	0	0
29	o	223	0	0	0	0
30	p	385	0	0	0	0
31	q	627	0	0	0	0
32	Y	677	0	342	8	0
32	Z	256	0	132	11	0
33	2	2968	0	1494	41	0
34	5	2397	0	1216	32	0
35	6	1903	0	964	28	0
All	All	55236	0	25739	813	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 813 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1355:SER:CB	16:S:17:GLY:HA3	1.26	1.57
1:A:1355:SER:CB	16:S:17:GLY:CA	1.88	1.49
10:M:309:ALA:HB3	18:U:358:GLU:CA	1.42	1.48
10:M:309:ALA:CB	18:U:358:GLU:CA	1.91	1.45
1:A:1355:SER:CA	16:S:17:GLY:HA3	1.43	1.43

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1961/2335 (84%)	1618 (82%)	328 (17%)	15 (1%)	19	60
2	B	898/972 (92%)	778 (87%)	108 (12%)	12 (1%)	12	48
3	C	238/536 (44%)	193 (81%)	44 (18%)	1 (0%)	34	72
4	D	313/515 (61%)	282 (90%)	30 (10%)	1 (0%)	41	76
5	E	325/579 (56%)	287 (88%)	37 (11%)	1 (0%)	41	76
6	F	305/357 (85%)	274 (90%)	30 (10%)	1 (0%)	41	76
7	G	130/504 (26%)	120 (92%)	9 (7%)	1 (1%)	19	60
7	H	133/504 (26%)	120 (90%)	12 (9%)	1 (1%)	19	60
7	I	132/504 (26%)	122 (92%)	10 (8%)	0	100	100
7	J	133/504 (26%)	116 (87%)	14 (10%)	3 (2%)	6	34
8	K	187/225 (83%)	167 (89%)	19 (10%)	1 (0%)	29	69
9	L	330/802 (41%)	297 (90%)	32 (10%)	1 (0%)	41	76
10	M	703/855 (82%)	600 (85%)	97 (14%)	6 (1%)	17	56
11	N	113/243 (46%)	87 (77%)	23 (20%)	3 (3%)	5	31
12	O	562/848 (66%)	476 (85%)	78 (14%)	8 (1%)	11	46
13	P	273/420 (65%)	232 (85%)	39 (14%)	2 (1%)	22	62
14	Q	140/144 (97%)	127 (91%)	12 (9%)	1 (1%)	22	62
15	R	94/229 (41%)	84 (89%)	10 (11%)	0	100	100
16	S	28/2752 (1%)	20 (71%)	5 (18%)	3 (11%)	0	8
17	T	196/908 (22%)	177 (90%)	18 (9%)	1 (0%)	29	69
19	V	153/166 (92%)	145 (95%)	7 (5%)	1 (1%)	22	62
20	W	160/255 (63%)	146 (91%)	12 (8%)	2 (1%)	12	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
21	X	92/225 (41%)	90 (98%)	2 (2%)	0	100	100
22	a	97/118 (82%)	95 (98%)	2 (2%)	0	100	100
22	h	96/118 (81%)	83 (86%)	8 (8%)	5 (5%)	2	19
23	b	74/86 (86%)	72 (97%)	2 (3%)	0	100	100
23	i	72/86 (84%)	69 (96%)	3 (4%)	0	100	100
24	c	77/92 (84%)	75 (97%)	2 (3%)	0	100	100
24	j	77/92 (84%)	75 (97%)	2 (3%)	0	100	100
25	d	72/76 (95%)	71 (99%)	1 (1%)	0	100	100
25	k	72/76 (95%)	70 (97%)	2 (3%)	0	100	100
26	e	82/126 (65%)	81 (99%)	1 (1%)	0	100	100
26	l	81/126 (64%)	80 (99%)	1 (1%)	0	100	100
27	f	67/240 (28%)	63 (94%)	4 (6%)	0	100	100
27	m	67/240 (28%)	66 (98%)	1 (2%)	0	100	100
28	g	80/119 (67%)	76 (95%)	4 (5%)	0	100	100
28	n	80/119 (67%)	77 (96%)	3 (4%)	0	100	100
All	All	8693/17096 (51%)	7611 (88%)	1012 (12%)	70 (1%)	24	60

5 of 70 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	988	ILE
1	A	1777	ILE
2	B	182	LYS
2	B	947	VAL
10	M	720	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/2108 (5%)	106 (100%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	55/866 (6%)	55 (100%)	0	100	100
3	C	19/459 (4%)	19 (100%)	0	100	100
4	D	12/441 (3%)	12 (100%)	0	100	100
5	E	11/502 (2%)	11 (100%)	0	100	100
6	F	10/300 (3%)	10 (100%)	0	100	100
7	G	10/435 (2%)	10 (100%)	0	100	100
7	H	11/435 (2%)	11 (100%)	0	100	100
7	I	11/435 (2%)	11 (100%)	0	100	100
7	J	11/435 (2%)	11 (100%)	0	100	100
8	K	5/196 (3%)	5 (100%)	0	100	100
9	L	6/709 (1%)	6 (100%)	0	100	100
10	M	24/749 (3%)	24 (100%)	0	100	100
11	N	4/209 (2%)	4 (100%)	0	100	100
12	O	17/751 (2%)	17 (100%)	0	100	100
13	P	14/361 (4%)	14 (100%)	0	100	100
14	Q	6/130 (5%)	6 (100%)	0	100	100
15	R	2/203 (1%)	2 (100%)	0	100	100
16	S	2/2432 (0%)	2 (100%)	0	100	100
17	T	6/838 (1%)	6 (100%)	0	100	100
19	V	9/134 (7%)	9 (100%)	0	100	100
20	W	6/218 (3%)	6 (100%)	0	100	100
21	X	3/195 (2%)	3 (100%)	0	100	100
22	a	4/110 (4%)	4 (100%)	0	100	100
22	h	4/110 (4%)	4 (100%)	0	100	100
23	b	4/74 (5%)	4 (100%)	0	100	100
23	i	4/74 (5%)	4 (100%)	0	100	100
24	c	1/84 (1%)	1 (100%)	0	100	100
24	j	1/84 (1%)	1 (100%)	0	100	100
25	d	3/66 (4%)	3 (100%)	0	100	100
25	k	3/66 (4%)	3 (100%)	0	100	100
26	e	3/101 (3%)	3 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
26	l	3/101 (3%)	3 (100%)	0	100	100
27	f	3/177 (2%)	3 (100%)	0	100	100
27	m	3/177 (2%)	3 (100%)	0	100	100
28	g	3/101 (3%)	3 (100%)	0	100	100
28	n	3/101 (3%)	3 (100%)	0	100	100
All	All	402/14967 (3%)	402 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
32	Y	30/324 (9%)	21 (70%)	1 (3%)
32	Z	11/324 (3%)	9 (81%)	0
33	2	137/188 (72%)	34 (24%)	2 (1%)
34	5	113/116 (97%)	42 (37%)	2 (1%)
35	6	87/106 (82%)	34 (39%)	1 (1%)
All	All	378/1058 (35%)	140 (37%)	6 (1%)

5 of 140 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
32	Y	60	U
32	Y	61	A
32	Y	62	A
32	Y	64	A
32	Y	65	G

5 of 6 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
34	5	78	U
34	5	93	U
35	6	53	A
33	2	23	A
32	Y	65	G

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2
10	M	1
2	B	1
12	O	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	M	33:PHE	C	34:SER	N	4.52
1	B	797:ALA	C	798:GLN	N	3.21
1	A	1914:MET	C	1915:VAL	N	3.11
1	O	209:PRO	C	210:PRO	N	1.78
1	A	1134:TRP	C	1135:PRO	N	1.63


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3545. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


### 6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180

The images above show central slices of the map in three orthogonal directions.



## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 175




Y Index: 171




Z Index: 189


### 6.3.2 Raw map



X Index: 174



Y Index: 180




Z Index: 188

The images above show the largest variance slices of the map in three orthogonal directions.


## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.016. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 5939  $\text{nm}^3$ ; this corresponds to an approximate mass of 5364 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.169 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.169 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.90	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	7.96	9.16	8.18

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.96 differs from the reported value 5.9 by more than 10 %

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-3545 and PDB model 5MQF. Per-residue inclusion information can be found in section [3](#) on page [12](#).


### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.016 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.016).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

























The table lists the average atom inclusion at the recommended contour level (0.016) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9871	 0.2020
2	 0.9835	 0.1140
5	 0.9946	 0.2160
6	 0.9958	 0.2300
A	 0.9951	 0.2760
B	 0.9993	 0.2670
C	 0.9854	 0.2710
D	 0.9918	 0.2840
E	 0.9982	 0.2420
F	 1.0000	 0.2340
G	 0.9573	 0.1250
H	 0.9353	 0.0670
I	 0.9986	 0.1550
J	 0.9670	 0.1200
K	 0.9990	 0.1880
L	 0.9946	 0.2530
M	 0.9916	 0.1740
N	 1.0000	 0.2930
O	 0.9989	 0.1990
P	 0.9893	 0.2280
Q	 1.0000	 0.2610
R	 0.9838	 0.2630
S	 1.0000	 0.2700
T	 0.9370	 0.2110
U	 0.9324	 0.0170
V	 1.0000	 0.2380
W	 1.0000	 0.0650
X	 1.0000	 0.1190
Y	 0.9852	 0.1590
Z	 0.9922	 0.2530
a	 0.9779	 0.1460
b	 0.9868	 0.1210
c	 1.0000	 0.1090
d	 0.9973	 0.1450
e	 0.9952	 0.1930



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
f	 1.0000	 0.1710
g	 1.0000	 0.1660
h	 0.9594	 0.0760
i	 0.9892	 0.0740
j	 0.9975	 0.0650
k	 0.9946	 0.0200
l	 0.9976	 0.1370
m	 1.0000	 0.1100
n	 0.9976	 0.1280
o	 0.9731	 0.0280
p	 0.6649	 0.0040
q	 0.9474	 0.0080