

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 2, 2023 – 01:29 AM EDT

PDB ID	:	6MQ6
Title	:	Mapping the binding trajectory of a suicide inhibitor in human indoleamine
		2,3-dioxygenase 1
Authors	:	Pham, K.N.; Yeh, S.R.
Deposited on	:	2018-10-09
Resolution	:	3.05  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 3.05 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6114 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	271	Total	С	Ν	0	$\mathbf{S}$	0	0	0
1	1 A	371	2943	1892	501	533	17	0	0	U
1	р	266	Total	С	Ν	0	S	0	0	0
I B	300	2915	1878	496	524	17	0	0	0	

• Molecule 1 is a protein called Indoleamine 2,3-dioxygenase 1.

Residue	Modelled	Actual	Comment	Reference
11	MET	-	initiating methionine	UNP P14902
404	LYS	-	expression tag	UNP P14902
405	GLY	-	expression tag	UNP P14902
406	GLU	-	expression tag	UNP P14902
407	LEU	-	expression tag	UNP P14902
408	ASN	-	expression tag	UNP P14902
409	SER	-	expression tag	UNP P14902
410	LYS	-	expression tag	UNP P14902
411	LEU	-	expression tag	UNP P14902
412	GLU	-	expression tag	UNP P14902
413	GLY	-	expression tag	UNP P14902
414	LYS	-	expression tag	UNP P14902
415	PRO	-	expression tag	UNP P14902
416	ILE	-	expression tag	UNP P14902
417	PRO	-	expression tag	UNP P14902
418	ASN	-	expression tag	UNP P14902
419	PRO	-	expression tag	UNP P14902
420	LEU	-	expression tag	UNP P14902
421	LEU	-	expression tag	UNP P14902
422	GLY	-	expression tag	UNP P14902
423	LEU	-	expression tag	UNP P14902
424	ASP	-	expression tag	UNP P14902
425	SER	-	expression tag	UNP P14902
426	THR	-	expression tag	UNP P14902
427	ARG	-	expression tag	UNP P14902
	Residue   11   404   405   406   407   408   409   410   411   412   413   414   415   416   417   418   419   420   421   422   423   424   425   426   427	Residue   Modelled     11   MET     404   LYS     405   GLY     406   GLU     407   LEU     408   ASN     409   SER     410   LYS     410   LYS     411   LEU     412   GLU     413   GLY     414   LYS     415   PRO     416   ILE     417   PRO     418   ASN     419   PRO     420   LEU     421   LEU     422   GLY     423   LEU     424   ASP     425   SER     426   THR     427   ARG	Residue   Modelled   Actual     11   MET   -     404   LYS   -     405   GLY   -     406   GLU   -     407   LEU   -     408   ASN   -     409   SER   -     410   LYS   -     410   LYS   -     411   LEU   -     412   GLU   -     413   GLY   -     414   LYS   -     415   PRO   -     415   PRO   -     416   ILE   -     417   PRO   -     418   ASN   -     419   PRO   -     420   LEU   -     421   LEU   -     422   GLY   -     423   LEU   -     423   LEU   -     424	ResidueModelledActualComment11MET-initiating methionine404LYS-expression tag405GLY-expression tag406GLU-expression tag407LEU-expression tag408ASN-expression tag409SER-expression tag410LYS-expression tag411LEU-expression tag412GLU-expression tag413GLY-expression tag414LYS-expression tag415PRO-expression tag416ILE-expression tag417PRO-expression tag418ASN-expression tag419PRO-expression tag412LEU-expression tag413ASN-expression tag414HXS-expression tag415PRO-expression tag416ILE-expression tag417PRO-expression tag418ASN-expression tag419PRO-expression tag420LEU-expression tag421LEU-expression tag422GLY-expression tag423LEU-expression tag424ASP-ex

There are 66 discrepancies between the modelled and reference sequences:

Continued on next page...



eference	

Chain	Residue	Modelled	Actual	Comment	Reference
А	428	THR	-	expression tag	UNP P14902
А	429	GLY	-	expression tag	UNP P14902
А	430	HIS	-	expression tag	UNP P14902
А	431	HIS	-	expression tag	UNP P14902
А	432	HIS	-	expression tag	UNP P14902
А	433	HIS	-	expression tag	UNP P14902
А	434	HIS	-	expression tag	UNP P14902
А	435	HIS	-	expression tag	UNP P14902
В	11	MET	-	initiating methionine	UNP P14902
В	404	LYS	-	expression tag	UNP P14902
В	405	GLY	-	expression tag	UNP P14902
В	406	GLU	-	expression tag	UNP P14902
В	407	LEU	-	expression tag	UNP P14902
В	408	ASN	-	expression tag	UNP P14902
В	409	SER	-	expression tag	UNP P14902
В	410	LYS	-	expression tag	UNP P14902
В	411	LEU	-	expression tag	UNP P14902
В	412	GLU	-	expression tag	UNP P14902
В	413	GLY	-	expression tag	UNP P14902
В	414	LYS	-	expression tag	UNP P14902
В	415	PRO	-	expression tag	UNP P14902
В	416	ILE	-	expression tag	UNP P14902
В	417	PRO	-	expression tag	UNP P14902
В	418	ASN	-	expression tag	UNP P14902
В	419	PRO	-	expression tag	UNP P14902
В	420	LEU	-	expression tag	UNP P14902
В	421	LEU	-	expression tag	UNP P14902
В	422	GLY	-	expression tag	UNP P14902
В	423	LEU	-	expression tag	UNP P14902
В	424	ASP	-	expression tag	UNP P14902
В	425	SER	-	expression tag	UNP P14902
В	426	THR	-	expression tag	UNP P14902
В	427	ARG	-	expression tag	UNP P14902
В	428	THR	-	expression tag	UNP P14902
В	429	GLY	-	expression tag	UNP P14902
В	430	HIS	-	expression tag	UNP P14902
В	431	HIS	-	expression tag	UNP P14902
В	432	HIS	-	expression tag	UNP P14902
В	433	HIS	-	expression tag	UNP P14902
В	434	HIS	-	expression tag	UNP P14902
В	435	HIS	-	expression tag	UNP P14902

Continued from previous page...

• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (for-



mula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	А	1	Total 43	C 34	Fe 1	N 4	0 4	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 4 is (2R)-N-(4-chlorophenyl)-2-[cis-4-(6-fluoroquinolin-4-yl)cyclohexyl]propanam ide (three-letter code: H7P) (formula: C<sub>24</sub>H<sub>24</sub>ClFN<sub>2</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
4	В	1	Total 29	С 24	Cl 1	F 1	N 2	0 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	70	Total   O     70   70	0	0
5	В	78	Total   O     78   78	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



## 3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	85.33Å 96.41Å 129.93Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	29.00 - 3.05	Depositor
% Data completeness	99.6 (29.00-3.05)	Depositor
(in resolution range)	33.0 (23.00 3.03)	Берозног
R <sub>merge</sub>	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.01 (at 3.05 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
$R, R_{free}$	0.184 , $0.229$	Depositor
Wilson B-factor $(Å^2)$	105.2	Xtriage
Anisotropy	0.032	Xtriage
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6114	wwPDB-VP
Average B, all atoms $(Å^2)$	114.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 4.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	Turne	Chain	Res	Tink	Bond lengths			Bond angles		
	туре	Unam		nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ
3	GOL	А	503	-	$5,\!5,\!5$	0.26	0	$5,\!5,\!5$	0.26	0
3	GOL	В	504	-	$5,\!5,\!5$	0.28	0	$5,\!5,\!5$	0.21	0
2	HEM	А	501	5,1	41,50,50	1.31	4 (9%)	45,82,82	1.74	7 (15%)
3	GOL	А	502	-	$5,\!5,\!5$	0.29	0	$5,\!5,\!5$	0.21	0
3	GOL	В	505	-	$5,\!5,\!5$	0.30	0	$5,\!5,\!5$	0.20	0
3	GOL	В	502	-	$5,\!5,\!5$	0.26	0	$5,\!5,\!5$	0.23	0
3	GOL	В	503	-	$5,\!5,\!5$	0.30	0	$5,\!5,\!5$	0.14	0
4	H7P	В	501	-	32,32,32	1.45	5 (15%)	45,45,45	1.33	6 (13%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	А	503	-	-	2/4/4/4	-
3	GOL	В	504	-	-	2/4/4/4	-
2	HEM	А	501	5,1	-	4/12/54/54	-
3	GOL	А	502	-	-	2/4/4/4	-
3	GOL	В	505	-	-	2/4/4/4	-
3	GOL	В	502	-	-	2/4/4/4	-
3	GOL	В	503	-	-	2/4/4/4	-
4	H7P	В	501	-	-	3/16/26/26	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	В	501	H7P	C9-C14	3.87	1.48	1.42
4	В	501	H7P	C8-C9	3.86	1.49	1.43
2	А	501	HEM	C1B-NB	-3.22	1.34	1.40
2	А	501	HEM	C4D-ND	-3.09	1.35	1.40
2	А	501	HEM	FE-NB	2.93	2.11	1.96
4	В	501	H7P	C18-N1	-2.82	1.35	1.41
4	В	501	H7P	C21-CL	2.45	1.79	1.74
4	В	501	H7P	C10-C11	2.35	1.40	1.36
2	А	501	HEM	CHB-C1B	2.16	1.40	1.35



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	А	501	HEM	CHC-C4B-NB	5.05	129.91	124.43
4	В	501	H7P	C5-C8-C9	4.12	124.97	120.67
2	А	501	HEM	CHD-C1D-ND	4.02	128.79	124.43
2	А	501	HEM	C1B-NB-C4B	3.67	108.86	105.07
2	А	501	HEM	CHA-C4D-ND	3.35	128.52	124.38
4	В	501	H7P	C15-N-C14	3.23	121.94	116.93
2	А	501	HEM	CHB-C1B-NB	3.19	128.32	124.38
4	В	501	H7P	C9-C14-N	-2.74	119.91	122.83
2	А	501	HEM	CHD-C1D-C2D	-2.58	120.94	124.98
4	В	501	H7P	C16-C8-C5	-2.42	118.05	120.55
4	B	501	H7P	C3-C2-C7	2.33	113.55	109.44
2	А	501	HEM	CHA-C4D-C3D	-2.15	121.30	125.33
4	B	501	H7P	C18-N1-C17	-2.07	122.45	127.40

All (13) bond angle outliers are listed below:

There are no chirality outliers.

All $(19)$	) torsion	outliers	are	listed	below:

Mol	Chain	Res	Type	Atoms
3	В	502	GOL	O1-C1-C2-O2
3	В	502	GOL	O1-C1-C2-C3
3	В	503	GOL	O1-C1-C2-O2
3	В	503	GOL	O1-C1-C2-C3
3	В	504	GOL	O1-C1-C2-C3
3	В	505	GOL	O1-C1-C2-C3
4	В	501	H7P	C6-C5-C8-C9
4	В	501	H7P	C6-C5-C8-C16
3	А	503	GOL	O1-C1-C2-O2
3	В	504	GOL	O1-C1-C2-O2
3	А	502	GOL	C1-C2-C3-O3
3	А	503	GOL	O1-C1-C2-C3
3	А	502	GOL	O2-C2-C3-O3
3	В	505	GOL	O1-C1-C2-O2
4	В	501	H7P	C-C1-C17-O
2	А	501	HEM	CAA-CBA-CGA-O1A
2	A	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	А	501	HEM	CAA-CBA-CGA-O2A

There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

## 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

## 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

