



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 02:30 am GMT

PDB ID : 5MRA  
Title : human SCBD (sorcin calcium binding domain) in complex with doxorubicin  
Authors : Ilari, A.; Fiorillo, A.; Colotti, G.; Genovese, I.  
Deposited on : 2016-12-22  
Resolution : 3.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1001 (3.90-3.58)
Clashscore	141614	1063 (3.90-3.58)
Ramachandran outliers	138981	1027 (3.90-3.58)
Sidechain outliers	138945	1023 (3.90-3.58)
RSRZ outliers	127900	1006 (3.92-3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	
1	B	167	
1	C	167	
1	D	167	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	B	203	-	-	-	X
4	DM2	B	204	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorcin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	167	1313	825	224	252	12	0	0	0
1	B	167	1299	813	222	252	12	0	0	0
1	C	167	1309	821	225	251	12	0	0	0
1	D	167	1305	820	222	251	12	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	MET	-	initiating methionine	UNP P30626
B	32	MET	-	initiating methionine	UNP P30626
C	32	MET	-	initiating methionine	UNP P30626
D	32	MET	-	initiating methionine	UNP P30626

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Mg 3	0	0
2	B	2	Total 2	Mg 2	0	0
2	C	3	Total 3	Mg 3	0	0
2	D	2	Total 2	Mg 2	0	0

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
3	B	1	4	2	1	1	0	0

- Molecule 4 is DOXORUBICIN (three-letter code: DM2) (formula:  $C_{27}H_{29}NO_{11}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	B	1	39	27	1	11	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total O 1 1	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Sorcin





- Molecule 1: Sorcin



- Molecule 1: Sorcin



- Molecule 1: Sorcin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.23Å 104.83Å 113.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.94 – 3.74 48.26 – 3.74	Depositor EDS
% Data completeness (in resolution range)	99.0 (76.94-3.74) 99.1 (48.26-3.74)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 3.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.192 , 0.285 0.199 , 0.293	Depositor DCC
$R_{free}$ test set	569 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	162.2	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 166.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5280	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	180.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DM2, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1339	0.66	0/1807
1	B	0.48	0/1325	0.67	0/1791
1	C	0.49	0/1335	0.75	2/1803 (0.1%)
1	D	0.53	0/1331	0.71	0/1798
All	All	0.49	0/5330	0.70	2/7199 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	2
1	D	0	5
All	All	0	10

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	149	SER	N-CA-C	5.63	126.21	111.00
1	C	149	SER	CB-CA-C	-5.24	100.14	110.10

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	178	THR	Peptide
1	A	46	ASP	Peptide
1	B	132	MET	Peptide
1	C	133	GLY	Peptide
1	C	134	PHE	Peptide
1	D	117	SER	Peptide
1	D	119	THR	Peptide
1	D	148	TYR	Peptide
1	D	170	THR	Peptide
1	D	65	GLY	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1313	0	1262	7	0
1	B	1299	0	1222	10	0
1	C	1309	0	1246	11	0
1	D	1305	0	1245	7	0
2	A	3	0	0	0	0
2	B	2	0	0	0	0
2	C	3	0	0	0	0
2	D	2	0	0	0	0
3	B	4	0	6	0	0
4	B	39	0	28	3	0
5	C	1	0	0	0	0
All	All	5280	0	5009	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:78:MET:HE2	1:D:102:LEU:HD21	1.78	0.66
1:C:120:VAL:HG13	1:C:125:LEU:HD11	1.86	0.58
1:A:116:ARG:O	1:A:116:ARG:HG2	2.03	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:148:TYR:O	1:C:150:THR:OG1	2.21	0.55
1:D:38:TYR:CZ	1:D:61:SER:HA	2.43	0.54
1:D:74:THR:HG23	1:D:161:ALA:HB2	1.89	0.54
1:C:74:THR:HG23	1:C:161:ALA:HA	1.90	0.53
1:A:43:ALA:HB3	1:A:49:ILE:HG22	1.89	0.53
1:C:176:ARG:HG2	1:C:184:VAL:HG13	1.90	0.52
1:B:166:LEU:O	1:B:170:THR:HG23	2.10	0.52
1:A:113:ASP:OD1	1:A:117:SER:OG	2.27	0.52
1:B:170:THR:HB	4:B:204:DM2:H212	1.92	0.51
1:D:48:GLN:CG	1:D:89:THR:HB	2.41	0.51
1:C:169:LEU:HD13	1:D:196:MET:SD	2.53	0.48
1:B:74:THR:HG23	1:B:161:ALA:HB2	1.95	0.47
1:B:138:PRO:HA	1:B:141:VAL:HG12	1.95	0.47
1:C:83:ASP:OD2	1:C:86:MET:HA	2.16	0.46
1:B:109:PHE:HB2	1:B:159:TYR:CE2	2.50	0.46
1:C:150:THR:HG22	1:C:150:THR:O	2.15	0.46
1:B:112:PHE:CD1	1:B:128:ALA:HB2	2.51	0.46
1:A:55:GLN:NE2	1:A:70:PHE:O	2.48	0.46
1:A:38:TYR:CE1	1:A:61:SER:HA	2.51	0.45
1:A:198:VAL:HG23	1:A:198:VAL:OXT	2.17	0.45
1:B:134:PHE:HB3	4:B:204:DM2:H211	1.99	0.43
1:C:67:TYR:OH	1:C:116:ARG:NH2	2.51	0.43
1:A:188:TYR:OH	1:B:170:THR:HG22	2.20	0.42
1:D:125:LEU:O	1:D:129:LEU:HD22	2.20	0.42
1:C:73:GLU:O	1:C:74:THR:C	2.59	0.41
1:B:174:ARG:NE	4:B:204:DM2:H3	2.35	0.41
1:C:148:TYR:CD1	1:C:148:TYR:N	2.89	0.41
1:C:120:VAL:HG22	1:C:125:LEU:CD1	2.51	0.41
1:B:126:GLN:HA	1:B:129:LEU:HD22	2.02	0.41
1:D:196:MET:SD	1:D:196:MET:N	2.94	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/167 (99%)	154 (93%)	11 (7%)	0	100	100
1	B	165/167 (99%)	149 (90%)	16 (10%)	0	100	100
1	C	165/167 (99%)	141 (86%)	20 (12%)	4 (2%)	6	37
1	D	165/167 (99%)	148 (90%)	16 (10%)	1 (1%)	25	61
All	All	660/668 (99%)	592 (90%)	63 (10%)	5 (1%)	19	56

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	69	PRO
1	C	83	ASP
1	C	149	SER
1	C	72	LEU
1	D	79	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/142 (99%)	131 (93%)	10 (7%)	14	46
1	B	137/142 (96%)	122 (89%)	15 (11%)	6	30
1	C	139/142 (98%)	119 (86%)	20 (14%)	3	20
1	D	139/142 (98%)	123 (88%)	16 (12%)	5	28
All	All	556/568 (98%)	495 (89%)	61 (11%)	6	29

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	55	GLN
1	A	81	MET

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	103	ASN
1	A	121	ASP
1	A	126	GLN
1	A	130	THR
1	A	132	MET
1	A	142	ASN
1	A	155	THR
1	B	45	GLN
1	B	56	ARG
1	B	60	GLN
1	B	76	ARG
1	B	103	ASN
1	B	111	SER
1	B	126	GLN
1	B	129	LEU
1	B	130	THR
1	B	132	MET
1	B	139	GLN
1	B	151	ASN
1	B	167	ARG
1	B	170	THR
1	B	183	VAL
1	C	60	GLN
1	C	69	PRO
1	C	85	ASP
1	C	90	MET
1	C	93	ASN
1	C	107	GLN
1	C	123	GLN
1	C	130	THR
1	C	135	ARG
1	C	139	GLN
1	C	149	SER
1	C	153	LYS
1	C	155	THR
1	C	157	ASP
1	C	162	CYS
1	C	166	LEU
1	C	174	ARG
1	C	175	ARG
1	C	177	ASP
1	C	180	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	48	GLN
1	D	50	ASP
1	D	57	CYS
1	D	61	SER
1	D	71	ASN
1	D	79	VAL
1	D	83	ASP
1	D	89	THR
1	D	123	GLN
1	D	126	GLN
1	D	129	LEU
1	D	135	ARG
1	D	136	LEU
1	D	155	THR
1	D	194	CYS
1	D	196	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	193	GLN
1	B	103	ASN
1	B	126	GLN
1	D	71	ASN
1	D	107	GLN
1	D	108	HIS
1	D	123	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DMS	B	203	-	3,3,3	0.44	0	3,3,3	0.70	0
4	DM2	B	204	-	41,43,43	2.22	8 (19%)	55,67,67	1.71	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DM2	B	204	-	-	11/13/60/60	0/5/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	204	DM2	C20-C5	5.58	1.49	1.41
4	B	204	DM2	C17-C16	5.52	1.48	1.40
4	B	204	DM2	C8-C9	5.29	1.49	1.40
4	B	204	DM2	C9-C16	5.22	1.48	1.39
4	B	204	DM2	C7-C8	4.41	1.50	1.41
4	B	204	DM2	C5-C4	4.12	1.48	1.40
4	B	204	DM2	C18-C7	3.91	1.50	1.41
4	B	204	DM2	C18-C17	3.87	1.49	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	204	DM2	C14-C13-C12	7.42	126.19	117.73
4	B	204	DM2	O4-C4-C5	4.43	122.02	115.85
4	B	204	DM2	C8-C9-C10	3.52	124.10	118.59
4	B	204	DM2	C16-C9-C10	-3.20	115.75	120.96

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	204	DM2	O4-C4-C3	-2.80	119.58	124.37
4	B	204	DM2	O10-C1'-C2'	2.67	113.57	108.41
4	B	204	DM2	O13-C13-C12	-2.52	118.05	122.09
4	B	204	DM2	O5'-C1'-C2'	-2.12	107.67	110.87
4	B	204	DM2	O13-C13-C14	-2.08	115.76	119.71

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	204	DM2	C2'-C1'-O10-C10
4	B	204	DM2	O5'-C1'-O10-C10
4	B	204	DM2	O12-C12-C13-O13
4	B	204	DM2	C15-C12-C13-O13
4	B	204	DM2	C15-C12-C13-C14
4	B	204	DM2	C12-C13-C14-O14
4	B	204	DM2	O13-C13-C14-O14
4	B	204	DM2	C3-C4-O4-C21
4	B	204	DM2	C5-C4-O4-C21
4	B	204	DM2	C11-C10-O10-C1'
4	B	204	DM2	C11-C12-C13-C14


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	204	DM2	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	167/167 (100%)	-0.01	2 (1%) 79 74	117, 163, 214, 237	0
1	B	167/167 (100%)	0.09	4 (2%) 59 52	135, 177, 230, 247	0
1	C	167/167 (100%)	0.23	8 (4%) 30 26	122, 193, 254, 283	0
1	D	167/167 (100%)	-0.08	1 (0%) 89 87	119, 173, 223, 258	0
All	All	668/668 (100%)	0.06	15 (2%) 62 55	117, 176, 239, 283	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	152	GLY	5.5
1	C	184	VAL	4.8
1	C	156	PHE	3.7
1	C	154	ILE	3.5
1	C	183	VAL	3.0
1	C	153	LYS	2.9
1	B	70	PHE	2.7
1	A	49	ILE	2.6
1	B	134	PHE	2.6
1	C	32	MET	2.5
1	C	109	PHE	2.4
1	B	135	ARG	2.4
1	A	188	TYR	2.2
1	D	184	VAL	2.2
1	B	198	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	201	1/1	0.38	0.20	106,106,106,106	0
2	MG	A	203	1/1	0.52	0.16	125,125,125,125	0
4	DM2	B	204	39/39	0.68	0.61	223,263,292,309	0
2	MG	B	202	1/1	0.70	0.18	146,146,146,146	0
3	DMS	B	203	4/4	0.72	0.65	151,163,168,170	0
2	MG	C	201	1/1	0.73	0.12	114,114,114,114	0
2	MG	D	201	1/1	0.74	0.26	103,103,103,103	0
2	MG	C	203	1/1	0.79	0.10	125,125,125,125	0
2	MG	A	202	1/1	0.83	0.22	149,149,149,149	0
2	MG	B	201	1/1	0.89	0.09	117,117,117,117	0
2	MG	C	202	1/1	0.90	0.06	128,128,128,128	0
2	MG	D	202	1/1	0.97	0.14	147,147,147,147	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.