

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 06:56 AM EDT

PDB ID	:	6MS2
Title	:	Crystal structure of the GH43 BlXynB protein from Bacillus licheniformis
Authors	:	Zanphorlin, L.M.; Morais, M.A.B.; Diogo, J.A.; Murakami, M.T.
Deposited on		
Resolution	:	2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	FAILED
:	1.13
:	FAILED
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.35.1
	: : : :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.49 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glycoside Hydrolase Family 43.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	511	Total 4073	C 2590	N 706	O 765	S 12	0	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	-22	MET	-	initiating methionine	UNP Q65MB6
А	-21	GLY	-	expression tag	UNP Q65MB6
А	-20	SER	-	expression tag	UNP Q65MB6
А	-19	SER	-	expression tag	UNP Q65MB6
А	-18	HIS	-	expression tag	UNP Q65MB6
А	-17	HIS	-	expression tag	UNP Q65MB6
А	-16	HIS	-	expression tag	UNP Q65MB6
А	-15	HIS	-	expression tag	UNP Q65MB6
А	-14	HIS	-	expression tag	UNP Q65MB6
А	-13	HIS	-	expression tag	UNP Q65MB6
А	-12	SER	-	expression tag	UNP Q65MB6
А	-11	SER	-	expression tag	UNP Q65MB6
А	-10	GLY	-	expression tag	UNP Q65MB6
А	-9	LEU	-	expression tag	UNP Q65MB6
А	-8	VAL	-	expression tag	UNP Q65MB6
А	-7	PRO	-	expression tag	UNP Q65MB6
А	-6	ARG	-	expression tag	UNP Q65MB6
А	-5	GLY	-	expression tag	UNP Q65MB6
А	-4	SER	-	expression tag	UNP Q65MB6
А	-3	HIS	-	expression tag	UNP Q65MB6
А	-2	MET	-	expression tag	UNP Q65MB6
А	-1	ALA	-	expression tag	UNP Q65MB6
А	0	SER	-	expression tag	UNP Q65MB6

There are 23 discrepancies between the modelled and reference sequences:

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).



Mol	Chain	Residues	Atom	IS	ZeroOcc	AltConf
2	А	1	Total 1	Ca 1	0	0

• Molecule 3 is water.

Mo	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	58	Total O 58 58	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	152.86Å 41.90Å 71.81Å	Depositor
a, b, c, α , β , γ	90.00° 91.70° 90.00°	Depositor
Resolution (Å)	33.31 - 2.49	Depositor
% Data completeness	87.8 (33.31-2.49)	Depositor
(in resolution range)	,	-
R_{merge}	0.09	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.06 (at 2.48 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.10_2148	Depositor
R, R_{free}	0.197 , 0.242	Depositor
Wilson B-factor $(Å^2)$	24.5	Xtriage
Anisotropy	0.544	Xtriage
L-test for twinning ²	$< L >=0.46, < L^2>=0.28$	Xtriage
Estimated twinning fraction	0.055 for -h,-k,l	Xtriage
Total number of atoms	4132	wwPDB-VP
Average B, all atoms $(Å^2)$	38.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

