

wwPDB EM Validation Summary Report (i)

Nov 7, 2022 – 04:08 PM EST

PDB ID : 6MSG EMDB ID : EMD-9219

Title: Cryo-EM structures and dynamics of substrate-engaged human 26S protea-

some

Authors : Mao, Y.D. Deposited on : 2018-10-16

Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at $\frac{\text{https://www.wwpdb.org/validation/2017/EMValidationReportHelp}}{\text{with specific help available everywhere you see the } \widehat{\textbf{i}} \text{ symbol.}$

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	U	953	86%	• 14%
2	V	533	12% 89%	• 10%
3	W	456	31%	
4	X	422	90%	10%
5	Y	389	96%	• •
6	Z	324	• 87%	12%
7	a	376	98%	••
8	b	377	51% 49%	
9	c	309	88%	5% 7%



 $Continued\ from\ previous\ page...$





Continued from previous page...

Mol	Chain	$oxed{ egin{array}{c c} \mathbf{Length} \end{array} }$	Quality of chain	
28	N	238	80%	20%
28	n	238	80%	20%
29	О	276	80%	20%
29	О	276	80%	20%
30	Р	204	100%	
30	p	204	100%	
31	Q	201	99%	
31	q	201	99%	
32	R	262	77%	23%
32	r	262	77%	23%
33	S	240	89%	11%
33	S	240	89%	11%
34	Т	263	81%	18%
34	t	263	81%	18%



2 Entry composition (i)

There are 38 unique types of molecules in this entry. The entry contains 104399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 26S proteasome non-ATPase regulatory subunit 1.

Mol	Chain	Residues		A	AltConf	Trace			
1	U	822	Total 6414	C 4072	N 1088	O 1210	S 44	0	0

• Molecule 2 is a protein called 26S proteasome non-ATPase regulatory subunit 3.

Mol	Chain	Residues		At	AltConf	Trace			
2	V	480	Total 3852	C 2444	N 684	O 710	S 14	0	0

• Molecule 3 is a protein called 26S proteasome non-ATPase regulatory subunit 12.

Mol	Chain	Residues		At	AltConf	Trace			
3	W	456	Total 3703	C 2339	N 635	O 704	S 25	0	0

• Molecule 4 is a protein called 26S proteasome non-ATPase regulatory subunit 11.

Mol	Chain	Residues		At	AltConf	Trace			
4	X	380	Total 3009	C 1918	N 509	O 570	S 12	0	0

• Molecule 5 is a protein called 26S proteasome non-ATPase regulatory subunit 6.

Mol	Chain	Residues		At	AltConf	Trace			
5	Y	378	Total 3115	C 1987	N 533	O 578	S 17	0	0

• Molecule 6 is a protein called 26S proteasome non-ATPase regulatory subunit 7.

Mol	Chain	Residues		Ato	AltConf	Trace			
6	Z	286	Total 2281	C 1457	N 392	O 427	S 5	0	0



• Molecule 7 is a protein called 26S proteasome non-ATPase regulatory subunit 13.

Mo	l Chain	Residues		At	AltConf	Trace			
7	a	373	Total 2995	C 1911	N 510	O 559	S 15	0	0

• Molecule 8 is a protein called 26S proteasome non-ATPase regulatory subunit 4.

Mol	Chain	Residues		Atoms					Trace
Q	h	191	Total	С	N	О	S	0	0
0	D	191	1458	910	261	279	8	0	U

• Molecule 9 is a protein called 26S proteasome non-ATPase regulatory subunit 14.

Mol	Chain	Residues		At	AltConf	Trace			
9	c	287	Total 2260	C 1430	N 389	O 422	S 19	0	0

• Molecule 10 is a protein called 26S proteasome non-ATPase regulatory subunit 8.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	d	257	Total 2116	C 1371	N 346	O 390	S 9	0	0

• Molecule 11 is a protein called 26S proteasome complex subunit SEM1.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$			AltConf	Trace
11	е	40	Total 334	C 200	N 55	O 77	S 2	0	0

• Molecule 12 is a protein called 26S proteasome non-ATPase regulatory subunit 2.

Mol	Chain	Residues		A	toms			AltConf	Trace
12	f	889	Total 6866	C 4315	N 1174	O 1331	S 46	0	0

• Molecule 13 is a protein called 26S proteasome regulatory subunit 7.

Mol	Chain	Residues		At	oms			AltConf	Trace
13	A	356	Total 2767	C 1741	N 488	O 521	S 17	0	0

• Molecule 14 is a protein called 26S proteasome regulatory subunit 4.



Mol	Chain	Residues		At	oms			AltConf	Trace
14	В	347	Total 2691	C 1694	N 455	O 530	S 12	0	0

• Molecule 15 is a protein called 26S proteasome regulatory subunit 8.

\mathbf{N}	/Iol	Chain	Residues		At	oms			AltConf	Trace
1	15	С	396	Total 3105	C 1954	N 558	O 576	S 17	0	0

• Molecule 16 is a protein called 26S proteasome regulatory subunit 6B.

Mol	Chain	Residues		At	oms			AltConf	Trace
16	D	380	Total 3040	C 1923	N 524	O 580	S 13	0	0

• Molecule 17 is a protein called 26S proteasome regulatory subunit 10B.

Mol	Chain	Residues		At	oms			AltConf	Trace
17	E	389	Total 3097	C 1947	N 552	O 581	S 17	0	0

• Molecule 18 is a protein called 26S proteasome regulatory subunit 6A.

Mol	Chain	Residues		At	oms			AltConf	Trace
18	F	395	Total 3098	C 1951	N 533	O 596	S 18	0	0

• Molecule 19 is a protein called Ubiquitin.

Mol	Chain	Residues		At	oms			AltConf	Trace
19	u	76	Total 603	C 378	N 107	O 117	S 1	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
u	63	ARG	LYS	conflict	UNP P0CG47

• Molecule 20 is a protein called substrate.



Mol	Chain	Residues	L	Ator	ns		AltConf	Trace
20	V	2/	Total	С	N	О	0	0
20	V	24	120	72	24	24		

• Molecule 21 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues		At		AltConf	Trace		
21	С	240	Total	С	N	О	S	0	0
21	G	240	1826	1160	305	348	13	U	U
21	G.	240	Total	С	N	О	S	0	0
21	g	240	1826	1160	305	348	13	U	U

• Molecule 22 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues		Ato		AltConf	Trace		
22	Н	232	20001	C 1081	- '		S 5	0	0
22	h	232	Total 1708	C 1081		O 333	S 5	0	0

• Molecule 23 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues		Ato	oms		AltConf	Trace	
23	I	250	Total 1912	C 1204	N 320	O 371	S 8	0	0
99	:	250				0	S	0	0
23		250	1912	1204	329	371	8	U	U

• Molecule 24 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues		Ato	oms		AltConf	Trace	
24	Ţ	239	Total	С	N	О	S	0	0
24	J	239	1704	1056	308	335	5	0	
24	;	239	Total	С	N	О	S	0	0
24	J	239	1704	1056	308	335	5		U

• Molecule 25 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues		\mathbf{At}	oms		AltConf	Trace	
25	K	228	Total 1722	C 1080		O 348	\sim	0	0
25	k	228	Total 1722	C 1080		O 348	S 10	0	0



• Molecule 26 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues		At	oms		AltConf	Trace	
26	Т	238	Total	С	N	О	S	0	0
20	ь	230	1850	1159	334	346	11	0	0
26	1	238	Total	С	N	О	S	0	0
20	1	230	1850	1159	334	346	11	0	U

• Molecule 27 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues		At	oms		AltConf	Trace	
27	М	240	Total	С	N	О	S	0	0
21	1V1	240	1856	1178	314	353	11	0	U
27	772	240	Total	С	N	О	S	0	0
21	m	Z40	1856	1178	314	353	11	0	U

• Molecule 28 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues		A	toms		AltConf	Trace	
28	N	191	Total 1430	_		_		0	0
28	n	191	Total 1430		N 245		S 12	0	0

• Molecule 29 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues		At	oms		AltConf	Trace	
29	О	220	Total 1643	C 1033	- '	O 318	D	0	0
29	О	220	Total 1643	C 1033		O 318	S 12	0	0

• Molecule 30 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues		At		AltConf	Trace		
30	D	204	Total	С	N	О	S	0	0
30	1	204	1585	1010	262	294	19	0	
30	n	204	Total	С	N	О	S	0	0
30	þ	204	1585	1010	262	294	19	0	U

• Molecule 31 is a protein called Proteasome subunit beta type-2.



Mol	Chain	Residues		Ato		AltConf	Trace		
31	Q	199	Total	_	= :	0	S	0	0
			1570	1006	265	290	9		
21		199	Total	\mathbf{C}	N	O	\mathbf{S}	0	0
31	q	199	1570	1006	265	290	9		U

• Molecule 32 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues		At	oms		AltConf	Trace	
32	D	201	Total	С	N	О	S	0	0
32	π	201	1548	974	273	292	9	0	U
32		201	Total	С	N	О	S	0	0
32	1	201	1548	974	273	292	9	0	U

• Molecule 33 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms			AltConf	Trace		
33	S	213	Total	С	N	О	S	0	0
00	S	210	1641	1036	282	313	10		
33	g	213	Total	С	N	О	S	0	0
- 55	S	213	1641	1036	282	313	10		U

• Molecule 34 is a protein called Proteasome subunit beta type-4.


Mol	Chain	Residues	Atoms			AltConf	Trace		
34	т	215	Total	С	N	О	S	0	0
04	1	210	1667	1052	285	318	12	0	0
34	+	215	Total	С	N	О	S	0	0
34	U	210	1667	1052	285	318	12	0	U

• Molecule 35 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
35	c	1	Total Zn 1 1	0


• Molecule 36 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).





Mol	Chain	Residues		Ato	oms			AltConf	
36	Λ	1	Total	С	N	О	Р	0	
30	A	1	27	10	5	10	2	U	
36	E	1	Total	С	N	О	Р	0	
30	Г	1	27	10	5	10	2	U	

• Molecule 37 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues		${f Atoms}$			AltConf	
37	С	1	Total 31	C 10		O 13	P 3	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues		Ato	oms			AltConf
37	D	1	Total	С	N	О	Р	0
31	D	1	31	10	5	13	3	U
37	E	1	Total	С	N	О	Р	0
31	E	1	31	10	5	13	3	U

 \bullet Molecule 38 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	AltConf
38	С	1	Total Mg 1 1	0
38	D	1	Total Mg 1 1	0
38	E	1	Total Mg 1 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 26S proteasome non-ATPase regulatory subunit 1



• Molecule 2. 200 proteasome non 1111 ase regulatory subtint 9



• Molecule 3: 26S proteasome non-ATPase regulatory subunit 12










• Molecule 8: 26S proteasome non-ATPase regulatory subunit 4





















• Molecule 22: Proteasome subunit alpha type-2











• Molecule 29	9: Proteasome subunit beta type-1	
Chain o:	80%	20%
ALA VALA VALA TYRA PRO VAL	617 PHE	E220 ILE GLU VAL LEU GLU GLU THR MET ASP THR SER
• Molecule 30): Proteasome subunit beta type-3	
Chain P:	100%	
There are no	outlier residues recorded for this chain.	
• Molecule 30): Proteasome subunit beta type-3	
Chain p:	100%	
There are no	outlier residues recorded for this chain.	
• Molecule 31	1: Proteasome subunit beta type-2	
Chain Q:	99%	
M1 L102 Q199 GLY SER		
• Molecule 31	1: Proteasome subunit beta type-2	
Chain q:	99%	:
M1 L102 Q199 GLY SER		
• Molecule 32	2: Proteasome subunit beta type-5	
Chain R:	77%	23%
ALA LEU ALA SER VAL LEU GLU ARG PRO	PRO ASN ASN ASN GLN GLN GLN GLY	TRP GLY VAL PRO GLU
G201 SER THR PRO		
• Molecule 32	2: Proteasome subunit beta type-5	
Chain r:	77%	23%
ALA LEU ALA SER VAL LEU GLU ARG PRO	PRO A VAL A VAL A KAG GLN GLN GLN GLN GLN GLN GLN GLN GLN GL	TRP GLY VAL VAL PRO GLU GLU GLU GLU GLY GLY HLE HLE HTS





• Molecule 33: Proteasome subunit beta type-1

Chain S:

89%

11%

• Molecule 33: Proteasome subunit beta type-1

Chain s: 89% 11%

• Molecule 34: Proteasome subunit beta type-4

Chain T: 81% 18%

• Molecule 34: Proteasome subunit beta type-4

Chain t: 81% 18%



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	112776	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	44	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.014	Depositor
Minimum map value	-0.004	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0035	Depositor
Map size (Å)	411.0, 411.0, 411.0	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.685, 0.685, 0.685	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, ATP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	ond lengths	В	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	U	0.30	0/6530	0.54	1/8840 (0.0%)
2	V	0.32	0/3929	0.59	0/5309
3	W	0.29	0/3751	0.56	0/5042
4	X	0.28	0/3053	0.52	0/4115
5	Y	0.31	0/3173	0.57	0/4273
6	Z	0.32	0/2324	0.59	2/3150 (0.1%)
7	a	0.29	0/3053	0.57	1/4133 (0.0%)
8	b	0.28	0/1478	0.55	0/2001
9	c	0.34	0/2302	0.65	3/3110 (0.1%)
10	d	0.30	0/2162	0.61	1/2919 (0.0%)
11	е	0.32	0/338	0.73	0/450
12	f	0.33	1/6980 (0.0%)	0.71	6/9433 (0.1%)
13	A	0.29	0/2814	0.53	0/3801
14	В	0.28	0/2730	0.53	0/3688
15	С	0.30	0/3146	0.54	0/4226
16	D	0.31	0/3090	0.61	0/4168
17	Е	0.31	0/3145	0.59	1/4233 (0.0%)
18	F	0.30	0/3137	0.62	6/4223 (0.1%)
19	u	0.23	0/609	0.39	0/819
21	G	0.34	0/1859	0.55	$1/2523 \ (0.0\%)$
21	g	0.34	0/1859	0.55	$1/2523 \ (0.0\%)$
22	Н	0.37	0/1743	0.53	0/2372
22	h	0.37	0/1743	0.53	0/2372
23	I	0.32	0/1942	0.53	0/2628
23	i	0.32	0/1942	0.53	0/2628
24	J	0.31	0/1728	0.51	0/2358
24	j	0.31	0/1728	0.51	0/2358
25	K	0.30	0/1747	0.53	0/2364
25	k	0.30	0/1747	0.53	0/2364
26	L	0.32	0/1885	0.54	0/2552
26	1	0.32	0/1885	0.54	0/2552
27	M	0.35	0/1891	0.53	0/2552



Mol	Chain	Во	ond lengths	В	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
27	m	0.35	0/1891	0.53	0/2552
28	N	0.32	0/1454	0.49	0/1967
28	n	0.32	0/1454	0.49	0/1967
29	О	0.32	0/1670	0.51	0/2265
29	О	0.32	0/1670	0.51	0/2265
30	Р	0.32	0/1614	0.50	0/2177
30	p	0.32	0/1614	0.50	0/2177
31	Q	0.34	0/1603	0.56	0/2174
31	q	0.34	0/1603	0.56	0/2174
32	R	0.35	0/1579	0.48	0/2134
32	r	0.35	0/1579	0.48	0/2134
33	S	0.32	0/1671	0.50	0/2253
33	s	0.32	0/1671	0.50	0/2253
34	Т	0.33	0/1700	0.50	0/2305
34	t	0.33	0/1700	0.50	0/2305
All	All	0.32	1/105916 (0.0%)	0.56	23/143181 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	V	0	1
4	X	0	1
6	Z	0	2
7	a	0	1
9	c	0	2
10	d	0	2
11	е	0	2
12	f	0	12
13	A	0	1
15	С	0	1
16	D	0	4
18	F	0	1
All	All	0	30

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
12	f	192	VAL	C-N	5.94	1.45	1.34



The worst	5	of 2	23	bond	angle	outliers	are	listed	below:
110 WOID	$\overline{}$	01.		OIIG	211510	Cathere	COL C	IID CCC	OCIO III.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
21	g	210	PHE	C-N-CA	8.26	142.34	121.70
21	G	210	PHE	C-N-CA	8.25	142.34	121.70
12	f	830	LEU	CA-CB-CG	7.96	133.61	115.30
18	F	348	LEU	CA-CB-CG	7.24	131.94	115.30
12	f	759	LEU	CA-CB-CG	6.75	130.81	115.30

There are no chirality outliers.

5 of 30 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	V	29	PRO	Peptide
4	X	393	VAL	Peptide
6	Z	144	VAL	Peptide
6	Z	183	THR	Peptide
7	a	214	GLY	Peptide

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	U	816/953~(86%)	753 (92%)	62 (8%)	1 (0%)	51 84
2	V	478/533~(90%)	426 (89%)	52 (11%)	0	100 100
3	W	454/456 (100%)	408 (90%)	46 (10%)	0	100 100
4	X	$378/422 \ (90\%)$	357 (94%)	20 (5%)	1 (0%)	41 75
5	Y	$376/389 \; (97\%)$	345 (92%)	31 (8%)	0	100 100
6	Z	284/324~(88%)	247 (87%)	34 (12%)	3 (1%)	14 52



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentil	les
7	a	371/376~(99%)	341 (92%)	28 (8%)	2 (0%)	29 68	3
8	b	189/377~(50%)	167 (88%)	22 (12%)	0	100 10)0
9	c	$285/309\ (92\%)$	238 (84%)	44 (15%)	3 (1%)	14 52	2
10	d	255/349~(73%)	212 (83%)	42 (16%)	1 (0%)	34 72	2
11	e	36/70~(51%)	24 (67%)	10 (28%)	2 (6%)	2 17	
12	f	887/892~(99%)	712 (80%)	164 (18%)	11 (1%)	13 50)
13	A	354/433~(82%)	306 (86%)	47 (13%)	1 (0%)	41 75	5
14	В	345/440~(78%)	305 (88%)	40 (12%)	0	100 10)0
15	С	394/398~(99%)	344 (87%)	50 (13%)	0	100 10)0
16	D	378/418 (90%)	326 (86%)	49 (13%)	3 (1%)	19 58	3
17	Е	387/403 (96%)	340 (88%)	42 (11%)	5 (1%)	12 48	3
18	F	391/439 (89%)	358 (92%)	29 (7%)	4 (1%)	15 54	Ł
19	u	74/76 (97%)	67 (90%)	6 (8%)	1 (1%)	11 46	j
21	G	238/245 (97%)	227 (95%)	11 (5%)	0	100 10)0
21	g	238/245 (97%)	227 (95%)	11 (5%)	0	100 10)0
22	Н	230/233 (99%)	216 (94%)	14 (6%)	0	100 10)0
22	h	230/233~(99%)	217 (94%)	13 (6%)	0	100 10)0
23	I	248/260 (95%)	229 (92%)	19 (8%)	0	100 10)0
23	i	248/260 (95%)	229 (92%)	19 (8%)	0	100 10)0
24	J	237/247 (96%)	223 (94%)	14 (6%)	0	100 10)0
24	j	237/247 (96%)	223 (94%)	14 (6%)	0	100 10)0
25	K	224/240 (93%)	208 (93%)	16 (7%)	0	100 10)0
25	k	224/240 (93%)	208 (93%)	16 (7%)	0	100 10)0
26	L	236/268 (88%)	225 (95%)	11 (5%)	0	100 10)0
26	1	236/268 (88%)	225 (95%)	11 (5%)	0	100 10)0
27	M	238/254 (94%)	219 (92%)	19 (8%)	0	100 10)0
27	m	238/254 (94%)	219 (92%)	19 (8%)	0	100 10)0
28	N	189/238 (79%)	184 (97%)	5 (3%)	0	100 10)0
28	n	189/238 (79%)	184 (97%)	5 (3%)	0	100 10)0
29	О	218/276 (79%)	211 (97%)	7 (3%)	0	100 10)0
29	О	218/276 (79%)	211 (97%)	7 (3%)	0	100 10)0



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
30	Р	$202/204\ (99\%)$	191 (95%)	11 (5%)	0	100	100
30	p	$202/204\ (99\%)$	191 (95%)	11 (5%)	0	100	100
31	Q	197/201 (98%)	184 (93%)	13 (7%)	0	100	100
31	q	197/201 (98%)	184 (93%)	13 (7%)	0	100	100
32	R	$199/262 \ (76\%)$	194 (98%)	5 (2%)	0	100	100
32	r	199/262~(76%)	194 (98%)	5 (2%)	0	100	100
33	S	211/240 (88%)	203 (96%)	8 (4%)	0	100	100
33	s	211/240 (88%)	203 (96%)	8 (4%)	0	100	100
34	Т	213/263 (81%)	202 (95%)	11 (5%)	0	100	100
34	t	213/263 (81%)	202 (95%)	11 (5%)	0	100	100
All	All	13292/14919 (89%)	12109 (91%)	1145 (9%)	38 (0%)	44	75

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
9	c	271	ALA
9	c	272	ILE
10	d	200	PHE
12	f	808	ASN
12	f	853	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	Percentiles	
1	U	702/816~(86%)	699 (100%)	3 (0%)	91	96	
2	V	414/459 (90%)	412 (100%)	2 (0%)	88	94	
3	W	416/416 (100%)	411 (99%)	5 (1%)	71	87	
4	X	327/362~(90%)	327 (100%)	0	100	100	
5	Y	334/344~(97%)	331 (99%)	3 (1%)	78	90	
6	Z	$257/295\ (87\%)$	257 (100%)	0	100	100	



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
7	a	333/336~(99%)	331 (99%)	2 (1%)	86	94
8	b	$167/312\ (54\%)$	167 (100%)	0	100	100
9	c	$252/267\ (94\%)$	241 (96%)	11 (4%)	28	62
10	d	$231/293\ (79\%)$	231 (100%)	0	100	100
11	e	38/63~(60%)	38 (100%)	0	100	100
12	f	745/748~(100%)	727 (98%)	18 (2%)	49	76
13	A	$297/372\ (80\%)$	293 (99%)	4 (1%)	69	86
14	В	$300/385\ (78\%)$	297 (99%)	3 (1%)	76	88
15	С	340/346~(98%)	336 (99%)	4 (1%)	71	87
16	D	333/366~(91%)	325 (98%)	8 (2%)	49	76
17	E	$341/353\ (97\%)$	335 (98%)	6 (2%)	59	81
18	F	340/379~(90%)	337 (99%)	3 (1%)	78	90
19	u	68/68 (100%)	68 (100%)	0	100	100
21	G	193/209 (92%)	193 (100%)	0	100	100
21	g	193/209 (92%)	193 (100%)	0	100	100
22	Н	164/190 (86%)	164 (100%)	0	100	100
22	h	164/190 (86%)	164 (100%)	0	100	100
23	I	193/220 (88%)	193 (100%)	0	100	100
23	i	193/220 (88%)	193 (100%)	0	100	100
24	J	152/210 (72%)	152 (100%)	0	100	100
24	j	152/210 (72%)	152 (100%)	0	100	100
25	K	$186/202 \ (92\%)$	185 (100%)	1 (0%)	88	94
25	k	$186/202 \ (92\%)$	185 (100%)	1 (0%)	88	94
26	L	$198/229\ (86\%)$	197 (100%)	1 (0%)	88	94
26	1	198/229 (86%)	197 (100%)	1 (0%)	88	94
27	M	$192/211 \ (91\%)$	192 (100%)	0	100	100
27	m	192/211 (91%)	192 (100%)	0	100	100
28	N	148/180 (82%)	148 (100%)	0	100	100
28	n	148/180 (82%)	148 (100%)	0	100	100
29	О	177/227 (78%)	177 (100%)	0	100	100
29	О	177/227 (78%)	177 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
30	Р	172/173~(99%)	172 (100%)	0	100	100
30	p	172/173~(99%)	172 (100%)	0	100	100
31	Q	164/171~(96%)	163 (99%)	1 (1%)	86	94
31	q	164/171 (96%)	163 (99%)	1 (1%)	86	94
32	R	153/201 (76%)	153 (100%)	0	100	100
32	r	153/201 (76%)	153 (100%)	0	100	100
33	S	174/198~(88%)	174 (100%)	0	100	100
33	s	174/198~(88%)	174 (100%)	0	100	100
34	Т	175/214~(82%)	174 (99%)	1 (1%)	86	94
34	t	175/214~(82%)	174 (99%)	1 (1%)	86	94
All	All	11117/12650 (88%)	11037 (99%)	80 (1%)	84	93

5 of 80 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
16	D	273	LYS
18	F	344	ARG
16	D	312	ASN
17	Е	280	ASN
34	Т	37	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 112 such side chains are listed below:

Mol	Chain	Res	Type
12	f	815	HIS
31	q	101	ASN
17	Е	262	ASN
31	q	82	ASN
24	j	18	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Clasia	Das	Link	Во	ond leng	ths	Bond angles		
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
37	ATP	D	501	38	26,33,33	0.94	1 (3%)	31,52,52	1.71	5 (16%)
37	ATP	Е	401	38	26,33,33	0.94	1 (3%)	31,52,52	1.68	5 (16%)
36	ADP	F	501	-	24,29,29	0.95	1 (4%)	29,45,45	1.60	4 (13%)
36	ADP	A	501	-	24,29,29	0.96	1 (4%)	29,45,45	1.47	4 (13%)
37	ATP	С	501	38	26,33,33	0.93	1 (3%)	31,52,52	1.52	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
37	ATP	D	501	38	-	1/18/38/38	0/3/3/3
37	ATP	E	401	38	-	2/18/38/38	0/3/3/3
36	ADP	F	501	-	-	2/12/32/32	0/3/3/3
36	ADP	A	501	-	-	2/12/32/32	0/3/3/3
37	ATP	С	501	38	-	2/18/38/38	0/3/3/3

All (5) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
36	A	501	ADP	C5-C4	2.53	1.47	1.40
37	С	501	ATP	C5-C4	2.47	1.47	1.40
37	Е	401	ATP	C5-C4	2.42	1.47	1.40
36	F	501	ADP	C5-C4	2.41	1.47	1.40
37	D	501	ATP	C5-C4	2.39	1.47	1.40

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
37	D	501	ATP	PA-O3A-PB	-4.83	116.24	132.83
36	F	501	ADP	PA-O3A-PB	-4.22	118.33	132.83
37	Е	401	ATP	PA-O3A-PB	-4.14	118.61	132.83
37	D	501	ATP	C3'-C2'-C1'	4.08	107.12	100.98
37	Е	401	ATP	PB-O3B-PG	-3.99	119.15	132.83

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
36	F	501	ADP	C5'-O5'-PA-O1A
37	D	501	ATP	PG-O3B-PB-O3A
36	A	501	ADP	C5'-O5'-PA-O3A
36	F	501	ADP	C5'-O5'-PA-O3A
37	Е	401	ATP	C5'-O5'-PA-O3A

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-9219. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map




The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map







Y Index: 300




Z Index: 300





The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map







X Index: 362 Y Index: 284

Z Index: 278

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map







 \mathbf{Z}

The images above show the 3D surface view of the map at the recommended contour level 0.0035. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $1352~\mathrm{nm}^3$; this corresponds to an approximate mass of $1222~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.286 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-9219 and PDB model 6MSG. Per-residue inclusion information can be found in section 3 on page 13.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0035 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0035).



9.4 Atom inclusion (i)



At the recommended contour level, 90% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0035) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8118	0.3400
A	0.7840	0.2280
В	0.7096	0.1780
С	0.7902	0.2810
D	0.8188	0.3180
E	0.8411	0.3340
F	0.8109	0.2800
G	0.9298	0.4630
Н	0.9237	0.4670
I	0.8981	0.4230
J	0.9160	0.4030
K	0.8992	0.4320
L	0.9402	0.4680
M	0.9086	0.4370
N	0.9592	0.5140
O	0.9591	0.4980
Р	0.9602	0.5120
Q	0.9394	0.4780
R	0.9660	0.5100
S	0.9470	0.5030
Т	0.9618	0.5140
U	0.8541	0.2990
V	0.7461	0.2340
W	0.6096	0.2230
X	0.5849	0.2290
Y	0.8360	0.2470
Z	0.8457	0.3130
a	0.8264	0.2290
b	0.8487	0.2310
С	0.8342	0.3370
d	0.7461	0.2130
e	0.8061	0.2130
f	0.3013	0.0250
g	0.9115	0.4520
h	0.9046	0.4530



Continued on next page...



 $Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
i	0.8613	0.4120
j	0.8745	0.3810
k	0.8834	0.4350
1	0.9286	0.4660
m	0.8921	0.4310
n	0.9542	0.5100
О	0.9517	0.5030
р	0.9551	0.5070
q	0.9401	0.4830
r	0.9626	0.5200
\mathbf{s}	0.9502	0.5070
t	0.9643	0.5150
u	0.1858	0.1850
V	0.3083	0.2510

