



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 07:41 AM EDT

PDB ID : 6MSU
Title : Integrin alphaVBeta3 in complex with EETI-II 2.5F
Authors : van Agthoven, J.F.; Arnaout, M.A.
Deposited on : 2018-10-18
Resolution : 3.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.11 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 13158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	924	7196	4556	1221	1384	35	0	0	0

- Molecule 2 is a protein called Integrin beta-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	690	5294	3250	904	1070	70	0	0	0

- Molecule 3 is a protein called Engineered EETI-II 2.5F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	32	224	132	42	44	6	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	4	50	28	2	20	0	0	0
4	G	4	50	28	2	20	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	E	2	28	16	2	10	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	K	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	J	3	Total	C	N	O	0	0	0
			39	22	2	15			
7	L	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	B	1	Total	C	N	O	0	0
			14	8	1	5		
8	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	5	Total	Mn	0	0
			5	5		
9	B	3	Total	Mn	0	0
			3	3		

- Molecule 10 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	1	Total	Cl	0	0
			1	1		

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	1	Total	O	0	0
			1	1		
11	B	2	Total	O	0	0
			2	2		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.89Å 129.89Å 305.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	3.21 – 3.11	Depositor
% Data completeness (in resolution range)	100.0 (3.21-3.11)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Wilson B-factor (Å ²)	58.0	Xtrriage
Anisotropy	0.007	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
Total number of atoms	13158	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

28 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	D	1	4,1	14,14,15	0.39	0	17,19,21	0.62	0
4	NAG	D	2	4	14,14,15	0.25	0	17,19,21	0.41	0
4	BMA	D	3	4	11,11,12	1.06	0	15,15,17	1.17	2 (13%)
4	MAN	D	4	4	11,11,12	0.90	0	15,15,17	1.17	1 (6%)
5	NAG	E	1	5,1	14,14,15	0.46	0	17,19,21	1.32	2 (11%)
5	NAG	E	2	5	14,14,15	0.40	0	17,19,21	0.55	0
6	NAG	F	1	6,1	14,14,15	0.33	0	17,19,21	0.45	0
6	NAG	F	2	6	14,14,15	0.21	0	17,19,21	0.50	0
6	BMA	F	3	6	11,11,12	0.81	0	15,15,17	1.04	1 (6%)
6	BMA	F	4	6	11,11,12	1.82	4 (36%)	15,15,17	1.22	2 (13%)
6	MAN	F	5	6	11,11,12	1.12	2 (18%)	15,15,17	1.48	4 (26%)
6	MAN	F	6	6	11,11,12	1.08	1 (9%)	15,15,17	1.39	3 (20%)
4	NAG	G	1	4,1	14,14,15	0.37	0	17,19,21	0.57	0
4	NAG	G	2	4	14,14,15	0.62	1 (7%)	17,19,21	0.63	0
4	BMA	G	3	4	11,11,12	0.94	0	15,15,17	0.76	0
4	MAN	G	4	4	11,11,12	0.87	0	15,15,17	1.15	2 (13%)
5	NAG	H	1	5,1	14,14,15	0.36	0	17,19,21	0.51	0
5	NAG	H	2	5	14,14,15	0.43	0	17,19,21	0.42	0
5	NAG	I	1	5,1	14,14,15	0.37	0	17,19,21	0.44	0
5	NAG	I	2	5	14,14,15	0.41	0	17,19,21	0.92	2 (11%)
7	NAG	J	1	7,1	14,14,15	0.54	0	17,19,21	0.59	0
7	NAG	J	2	7	14,14,15	0.41	0	17,19,21	0.43	0
7	BMA	J	3	7	11,11,12	0.83	0	15,15,17	0.82	0
5	NAG	K	1	2,5	14,14,15	0.32	0	17,19,21	0.48	0
5	NAG	K	2	5	14,14,15	0.28	0	17,19,21	0.85	1 (5%)
7	NAG	L	1	2,7	14,14,15	0.36	0	17,19,21	0.61	0
7	NAG	L	2	7	14,14,15	1.01	1 (7%)	17,19,21	1.45	1 (5%)
7	BMA	L	3	7	11,11,12	1.29	2 (18%)	15,15,17	1.30	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	4/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	MAN	D	4	4	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	5,1	-	5/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
6	NAG	F	1	6,1	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	1/6/23/26	0/1/1/1
6	BMA	F	3	6	-	1/2/19/22	0/1/1/1
6	BMA	F	4	6	-	2/2/19/22	0/1/1/1
6	MAN	F	5	6	-	0/2/19/22	0/1/1/1
6	MAN	F	6	6	-	0/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	1/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
5	NAG	H	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	NAG	I	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	I	2	5	-	2/6/23/26	0/1/1/1
7	NAG	J	1	7,1	-	2/6/23/26	0/1/1/1
7	NAG	J	2	7	-	4/6/23/26	0/1/1/1
7	BMA	J	3	7	-	0/2/19/22	0/1/1/1
5	NAG	K	1	2,5	-	1/6/23/26	0/1/1/1
5	NAG	K	2	5	-	4/6/23/26	0/1/1/1
7	NAG	L	1	2,7	-	2/6/23/26	0/1/1/1
7	NAG	L	2	7	-	0/6/23/26	0/1/1/1
7	BMA	L	3	7	-	1/2/19/22	1/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	2	NAG	O5-C1	3.62	1.49	1.43
6	F	4	BMA	C4-C5	3.23	1.59	1.53
6	F	4	BMA	C4-C3	2.64	1.59	1.52
6	F	5	MAN	C1-C2	2.57	1.58	1.52
6	F	4	BMA	O5-C5	2.55	1.48	1.43
6	F	4	BMA	C2-C3	2.53	1.56	1.52
6	F	6	MAN	C1-C2	2.46	1.57	1.52
7	L	3	BMA	O5-C5	2.39	1.48	1.43
4	G	2	NAG	O5-C1	-2.24	1.40	1.43
6	F	5	MAN	C2-C3	2.22	1.55	1.52
7	L	3	BMA	C1-C2	2.19	1.57	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	L	2	NAG	C1-O5-C5	5.42	119.53	112.19
5	E	1	NAG	C2-N2-C7	4.43	129.21	122.90
7	L	3	BMA	C1-O5-C5	3.60	117.08	112.19
4	D	4	MAN	C1-O5-C5	3.12	116.42	112.19
6	F	5	MAN	C1-C2-C3	2.96	113.31	109.67
6	F	6	MAN	O5-C1-C2	2.73	114.99	110.77
6	F	3	BMA	C1-O5-C5	2.56	115.67	112.19
4	G	4	MAN	C1-O5-C5	2.54	115.63	112.19
4	D	3	BMA	O3-C3-C2	2.53	114.83	109.99
5	K	2	NAG	C1-O5-C5	2.47	115.54	112.19
5	I	2	NAG	C1-O5-C5	2.40	115.44	112.19
6	F	5	MAN	O5-C1-C2	2.38	114.45	110.77
6	F	5	MAN	C1-O5-C5	2.31	115.32	112.19
5	I	2	NAG	C3-C4-C5	2.27	114.28	110.24
6	F	6	MAN	O2-C2-C3	-2.20	105.74	110.14
5	E	1	NAG	C1-C2-N2	2.18	114.21	110.49
6	F	5	MAN	O2-C2-C3	-2.12	105.89	110.14
4	G	4	MAN	O2-C2-C3	-2.11	105.91	110.14
4	D	3	BMA	C1-O5-C5	2.07	115.00	112.19
6	F	6	MAN	C1-C2-C3	2.03	112.16	109.67
6	F	4	BMA	C1-O5-C5	2.03	114.94	112.19
6	F	4	BMA	C3-C4-C5	2.00	113.81	110.24

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	1	NAG	O5-C5-C6-O6
6	F	1	NAG	O5-C5-C6-O6
7	J	1	NAG	C4-C5-C6-O6
5	K	2	NAG	O5-C5-C6-O6
7	J	1	NAG	O5-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6
5	H	2	NAG	O5-C5-C6-O6
5	H	1	NAG	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
5	H	2	NAG	C4-C5-C6-O6
6	F	1	NAG	C4-C5-C6-O6
5	E	1	NAG	O5-C5-C6-O6
7	J	2	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	D	4	MAN	C4-C5-C6-O6
5	E	1	NAG	C4-C5-C6-O6
5	K	2	NAG	C4-C5-C6-O6
7	L	1	NAG	O5-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
7	L	1	NAG	C4-C5-C6-O6
7	J	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C8-C7-N2-C2
4	D	2	NAG	O7-C7-N2-C2
5	E	1	NAG	C8-C7-N2-C2
5	E	1	NAG	O7-C7-N2-C2
5	E	2	NAG	C8-C7-N2-C2
5	E	2	NAG	O7-C7-N2-C2
5	I	2	NAG	C8-C7-N2-C2
5	I	2	NAG	O7-C7-N2-C2
5	K	2	NAG	C8-C7-N2-C2
5	K	2	NAG	O7-C7-N2-C2
4	D	2	NAG	O5-C5-C6-O6
5	H	1	NAG	C4-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
6	F	4	BMA	C4-C5-C6-O6
4	D	4	MAN	O5-C5-C6-O6
6	F	4	BMA	O5-C5-C6-O6
5	I	1	NAG	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
6	F	3	BMA	O5-C5-C6-O6
7	L	3	BMA	O5-C5-C6-O6
5	K	1	NAG	O5-C5-C6-O6
4	G	3	BMA	O5-C5-C6-O6
7	J	2	NAG	C1-C2-N2-C7
5	E	1	NAG	C3-C2-N2-C7
5	I	1	NAG	C4-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
7	J	2	NAG	C3-C2-N2-C7


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	L	3	BMA	C1-C2-C3-C4-C5-O5


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for oligosaccharide.


















4.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 9 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	NAG	B	702	2	14,14,15	0.47	0	17,19,21	0.78	1 (5%)
8	NAG	A	1020	1	14,14,15	0.65	0	17,19,21	0.61	0
8	NAG	A	1021	1	14,14,15	0.46	0	17,19,21	0.58	0
8	NAG	B	701	2	14,14,15	0.85	1 (7%)	17,19,21	0.92	1 (5%)
8	NAG	A	1017	1	14,14,15	0.33	0	17,19,21	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	B	702	2	-	1/6/23/26	0/1/1/1
8	NAG	A	1020	1	-	4/6/23/26	0/1/1/1
8	NAG	A	1021	1	-	2/6/23/26	0/1/1/1
8	NAG	B	701	2	-	1/6/23/26	0/1/1/1
8	NAG	A	1017	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	701	NAG	O5-C1	2.52	1.47	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	701	NAG	C1-O5-C5	3.52	116.96	112.19
8	B	702	NAG	C1-O5-C5	2.69	115.84	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	1017	NAG	C4-C5-C6-O6
8	A	1017	NAG	O5-C5-C6-O6
8	A	1020	NAG	O5-C5-C6-O6
8	A	1020	NAG	C4-C5-C6-O6
8	B	701	NAG	O5-C5-C6-O6
8	A	1020	NAG	C1-C2-N2-C7
8	A	1021	NAG	C4-C5-C6-O6
8	A	1021	NAG	O5-C5-C6-O6
8	B	702	NAG	O5-C5-C6-O6
8	A	1020	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.