

wwPDB X-ray Structure Validation Summary Report (i)

May 25, 2020 – 05:04 am BST

PDB ID	:	5MTR
Title	:	Crystal structure of M. tuberculosis InhA inhibited by PT512
Authors	:	Eltschkner, S.; Pschibul, A.; Spagnuolo, L.A.; Yu, W.; Tonge, P.J.; Kisker, C.
Deposited on		
Resolution	:	2.00 Å(reported)
Deposited on	:	2017-01-10

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4.02b-467 1.8.5 (274361), CSD as541be (2020)
9		
Xtriage (Phenix)		1.13
EDS	:	2.11
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R _{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	289	82%	9% • 7%
1	В	289	3% 84%	8% • 7%
1	С	289	83%	8% • 7%
1	D	289	4%	9% 7%
1	Е	289	% • 81%	8% • 8%
1	F	289	% • 83%	9% • 7%



Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	G	289	80%	9%	•	10%
1	Н	289	% 	8%		10%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 17729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	268	Total	С	Ν	0	S	0	2	0
	A	208	2004	1270	348	375	11	0	2	0
1	В	268	Total	С	Ν	Ο	S	0	2	0
	D	208	2002	1268	348	376	10	0	<u>ک</u>	0
1	Е	265	Total	С	Ν	0	S	0	2	0
		205	1985	1257	345	373	10		2	0
1	G	260	Total	С	Ν	Ο	S	0	2	0
	G	200	1954	1240	340	363	11			0
1	С	268	Total	С	Ν	0	S	0	1	0
		200	1999	1266	348	375	10	0		
1	D	268	Total	С	Ν	0	S	0	2	0
		200	2002	1268	348	376	10	0	2	0
1	F	268	Total	С	Ν	0	S	0	3	0
	T,	200	2007	1271	348	378	10	0	5	0
1	Н	261	Total	С	Ν	Ο	S	0	2	0
	11	201	1960	1244	341	364	11	U	2	0

• Molecule 1 is a protein called Enoyl-[acyl-carrier-protein] reductase [NADH].

There are 160 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-19	MET	-	initiating methionine	UNP P9WGR1
А	-18	GLY	-	expression tag	UNP P9WGR1
А	-17	SER	-	expression tag	UNP P9WGR1
А	-16	SER	-	expression tag	UNP P9WGR1
А	-15	HIS	-	expression tag	UNP P9WGR1
А	-14	HIS	-	expression tag	UNP P9WGR1
А	-13	HIS	-	expression tag	UNP P9WGR1
А	-12	HIS	-	expression tag	UNP P9WGR1
А	-11	HIS	-	expression tag	UNP P9WGR1
А	-10	HIS	-	expression tag	UNP P9WGR1
А	-9	SER	-	expression tag	UNP P9WGR1
А	-8	SER	-	expression tag	UNP P9WGR1
А	-7	GLY	-	expression tag	UNP P9WGR1



5MTR

		vious page			
Chain	Residue	Modelled	Actual	$\operatorname{Comment}$	Reference
A	-6	LEU	-	expression tag	UNP P9WGR1
A	-5	VAL	-	expression tag	UNP P9WGR1
A	-4	PRO	-	expression tag	UNP P9WGR1
A	-3	ARG	-	expression tag	UNP P9WGR1
A	-2	GLY	-	expression tag	UNP P9WGR1
A	-1	SER	-	expression tag	UNP P9WGR1
A	0	HIS	-	expression tag	UNP P9WGR1
В	-19	MET	-	initiating methionine	UNP P9WGR1
В	-18	GLY	-	expression tag	UNP P9WGR1
В	-17	SER	-	expression tag	UNP P9WGR1
В	-16	SER	-	expression tag	UNP P9WGR1
В	-15	HIS	-	expression tag	UNP P9WGR1
В	-14	HIS	-	expression tag	UNP P9WGR1
В	-13	HIS	-	expression tag	UNP P9WGR1
В	-12	HIS	-	expression tag	UNP P9WGR1
В	-11	HIS	-	expression tag	UNP P9WGR1
В	-10	HIS	-	expression tag	UNP P9WGR1
В	-9	SER	-	expression tag	UNP P9WGR1
В	-8	SER	-	expression tag	UNP P9WGR1
В	-7	GLY	-	expression tag	UNP P9WGR1
В	-6	LEU	-	expression tag	UNP P9WGR1
В	-5	VAL	-	expression tag	UNP P9WGR1
В	-4	PRO	-	expression tag	UNP P9WGR1
В	-3	ARG	-	expression tag	UNP P9WGR1
В	-2	GLY	-	expression tag	UNP P9WGR1
В	-1	SER	-	expression tag	UNP P9WGR1
В	0	HIS	-	expression tag	UNP P9WGR1
E	-19	MET	-	initiating methionine	UNP P9WGR1
Е	-18	GLY	-	expression tag	UNP P9WGR1
E	-17	SER	-	expression tag	UNP P9WGR1
E	-16	SER	-	expression tag	UNP P9WGR1
Е	-15	HIS	-	expression tag	UNP P9WGR1
Е	-14	HIS	-	expression tag	UNP P9WGR1
Е	-13	HIS	-	expression tag	UNP P9WGR1
Е	-12	HIS	-	expression tag	UNP P9WGR1
Е	-11	HIS	-	expression tag	UNP P9WGR1
Е	-10	HIS	-	expression tag	UNP P9WGR1
Е	-9	SER	-	expression tag	UNP P9WGR1
Е	-8	SER	-	expression tag	UNP P9WGR1
Е	-7	GLY	-	expression tag	UNP P9WGR1
Е	-6	LEU	-	expression tag	UNP P9WGR1
Е	-5	VAL	_	expression tag	UNP P9WGR1



5MTR

Continu	° 1	vious page			
Chain	Residue	Modelled	Actual	Comment	Reference
Ε	-4	PRO	-	expression tag	UNP P9WGR1
Е	-3	ARG	-	expression tag	UNP P9WGR1
Ε	-2	GLY	-	expression tag	UNP P9WGR1
Ε	-1	SER	-	expression tag	UNP P9WGR1
Е	0	HIS	-	expression tag	UNP P9WGR1
G	-19	MET	-	initiating methionine	UNP P9WGR1
G	-18	GLY	-	expression tag	UNP P9WGR1
G	-17	SER	-	expression tag	UNP P9WGR1
G	-16	SER	-	expression tag	UNP P9WGR1
G	-15	HIS	-	expression tag	UNP P9WGR1
G	-14	HIS	-	expression tag	UNP P9WGR1
G	-13	HIS	-	expression tag	UNP P9WGR1
G	-12	HIS	-	expression tag	UNP P9WGR1
G	-11	HIS	-	expression tag	UNP P9WGR1
G	-10	HIS	-	expression tag	UNP P9WGR1
G	-9	SER	-	expression tag	UNP P9WGR1
G	-8	SER	-	expression tag	UNP P9WGR1
G	-7	GLY	-	expression tag	UNP P9WGR1
G	-6	LEU	-	expression tag	UNP P9WGR1
G	-5	VAL	-	expression tag	UNP P9WGR1
G	-4	PRO	-	expression tag	UNP P9WGR1
G	-3	ARG	-	expression tag	UNP P9WGR1
G	-2	GLY	-	expression tag	UNP P9WGR1
G	-1	SER	-	expression tag	UNP P9WGR1
G	0	HIS	-	expression tag	UNP P9WGR1
С	-19	MET	-	initiating methionine	UNP P9WGR1
С	-18	GLY	-	expression tag	UNP P9WGR1
С	-17	SER	-	expression tag	UNP P9WGR1
С	-16	SER	-	expression tag	UNP P9WGR1
С	-15	HIS	-	expression tag	UNP P9WGR1
С	-14	HIS	-	expression tag	UNP P9WGR1
С	-13	HIS	-	expression tag	UNP P9WGR1
С	-12	HIS	-	expression tag	UNP P9WGR1
С	-11	HIS	-	expression tag	UNP P9WGR1
С	-10	HIS	_	expression tag	UNP P9WGR1
С	-9	SER	-	expression tag	UNP P9WGR1
С	-8	SER	-	expression tag	UNP P9WGR1
С	-7	GLY	-	expression tag	UNP P9WGR1
С	-6	LEU	-	expression tag	UNP P9WGR1
С	-5	VAL	-	expression tag	UNP P9WGR1
С	-4	PRO	-	expression tag	UNP P9WGR1
С	-3	ARG	_	expression tag	UNP P9WGR1




5MTR

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	GLY	-	expression tag	UNP P9WGR1
С	-1	SER	-	expression tag	UNP P9WGR1
С	0	HIS	-	expression tag	UNP P9WGR1
D	-19	MET	-	initiating methionine	UNP P9WGR1
D	-18	GLY	-	expression tag	UNP P9WGR1
D	-17	SER	-	expression tag	UNP P9WGR1
D	-16	SER	-	expression tag	UNP P9WGR1
D	-15	HIS	-	expression tag	UNP P9WGR1
D	-14	HIS	-	expression tag	UNP P9WGR1
D	-13	HIS	-	expression tag	UNP P9WGR1
D	-12	HIS	-	expression tag	UNP P9WGR1
D	-11	HIS	-	expression tag	UNP P9WGR1
D	-10	HIS	-	expression tag	UNP P9WGR1
D	-9	SER	-	expression tag	UNP P9WGR1
D	-8	SER	-	expression tag	UNP P9WGR1
D	-7	GLY	-	expression tag	UNP P9WGR1
D	-6	LEU	-	expression tag	UNP P9WGR1
D	-5	VAL	-	expression tag	UNP P9WGR1
D	-4	PRO	-	expression tag	UNP P9WGR1
D	-3	ARG	-	expression tag	UNP P9WGR1
D	-2	GLY	-	expression tag	UNP P9WGR1
D	-1	SER	-	expression tag	UNP P9WGR1
D	0	HIS	-	expression tag	UNP P9WGR1
F	-19	MET	-	initiating methionine	UNP P9WGR1
F	-18	GLY	-	expression tag	UNP P9WGR1
F	-17	SER	-	expression tag	UNP P9WGR1
F	-16	SER	-	expression tag	UNP P9WGR1
F	-15	HIS	-	expression tag	UNP P9WGR1
F	-14	HIS	-	expression tag	UNP P9WGR1
F	-13	HIS	-	expression tag	UNP P9WGR1
F	-12	HIS	-	expression tag	UNP P9WGR1
F	-11	HIS	-	expression tag	UNP P9WGR1
F	-10	HIS	-	expression tag	UNP P9WGR1
F	-9	SER	-	expression tag	UNP P9WGR1
F	-8	SER	-	expression tag	UNP P9WGR1
F	-7	GLY	-	expression tag	UNP P9WGR1
F	-6	LEU	-	expression tag	UNP P9WGR1
F	-5	VAL	-	expression tag	UNP P9WGR1
F	-4	PRO	-	expression tag	UNP P9WGR1
F	-3	ARG	-	expression tag	UNP P9WGR1
F	-2	GLY	-	expression tag	UNP P9WGR1
F	-1	SER	-	expression tag	UNP P9WGR1



Chain	Residue	Modelled	Actual Comment		Reference
F	0	HIS	-	expression tag	UNP P9WGR1
Н	-19	MET	-	initiating methionine	UNP P9WGR1
Н	-18	GLY	-	expression tag	UNP P9WGR1
Н	-17	SER	-	expression tag	UNP P9WGR1
Н	-16	SER	-	expression tag	UNP P9WGR1
Н	-15	HIS	-	expression tag	UNP P9WGR1
Н	-14	HIS	-	expression tag	UNP P9WGR1
Н	-13	HIS	-	expression tag	UNP P9WGR1
Н	-12	HIS	-	expression tag	UNP P9WGR1
Н	-11	HIS	-	expression tag	UNP P9WGR1
Н	-10	HIS	-	expression tag	UNP P9WGR1
Н	-9	SER	-	expression tag	UNP P9WGR1
Н	-8	SER	-	expression tag	UNP P9WGR1
Н	-7	GLY	-	expression tag	UNP P9WGR1
Н	-6	LEU	-	expression tag	UNP P9WGR1
Н	-5	VAL	-	expression tag	UNP P9WGR1
Н	-4	PRO	-	expression tag	UNP P9WGR1
Н	-3	ARG	-	expression tag	UNP P9WGR1
Н	-2	GLY	-	expression tag	UNP P9WGR1
Н	-1	SER	-	expression tag	UNP P9WGR1
Н	0	HIS	-	expression tag	UNP P9WGR1


• Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
2	А	1	Total	С	Ν	Ο	Р	0	0
	A	1	44	21	7	14	2	0	0
2	В	1	Total	С	Ν	Ο	Р	0	0
	D	1	44	21	7	14	2	0	0
2	Е	1	Total	С	Ν	Ο	Р	0	0
		T	44	21	7	14	2	0	0
2	G	1	Total	С	Ν	Ο	Р	0	0
	G	T	44	21	7	14	2	0	0
2	С	1	Total	С	Ν	Ο	Р	0	0
	U	I	44	21	7	14	2	0	0
2	D	1	Total	С	Ν	Ο	Р	0	0
		T	44	21	7	14	2	0	0
2	F	1	Total	С	Ν	Ο	Р	0	0
	Ľ		44	21	7	14	2	0	0
2	Н	1	Total	С	Ν	Ο	Р	0	0
	11		44	21	7	14	2	0	0

• Molecule 3 is 2-[4-[(4-cyclopentyl-1,2,3-triazol-1-yl)methyl]-2-oxidanyl-phenoxy]benzenecarb onitrile (three-letter code: XT0) (formula: C₂₁H₂₀N₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C N O 27 21 4 2	0	0
3	В	1	Total C N O 27 21 4 2	0	0
3	Е	1	Total C N O 27 21 4 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total C N O	0	0
	u	L	27 21 4 2	0	0
3	С	1	Total C N O	0	0
	U	T	27 21 4 2	0	0
3	п	1	Total C N O	0	0
	D	T	27 21 4 2	0	0
3	F	1	Total C N O	0	0
J J	Г	T	27 21 4 2	0	0
3	Н	1	Total C N O	0	0
0	11	1	27 21 4 2	U	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Н	1	Total Cl 1 1	0	0
4	С	1	Total Cl 1 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	173	Total O 173 173	0	0
5	В	174	Total O 174 174	0	0
5	Е	149	Total O 149 149	0	0
5	G	136	Total O 136 136	0	0
5	С	160	Total O 160 160	0	0
5	D	153	Total O 153 153	0	0
5	F	149	Total O 149 149	0	0
5	Н	152	Total O 152 152	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]








• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]

Chain C:		E	33%									8	3%	•	7%			
MET SER SER SER SER SER SER SER SER SER SER	12 D6	D18	R45 L46	I50	R53	K57	E68	T79 E80	<mark>(85</mark>	<mark>6</mark> 90	<mark>Y113</mark>	<mark>S126</mark>	S129	K132	S143	T162	R177	P193


• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]



• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]

•	6			
Chain F:	83%	9%	•	7%

• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants	87.59Å 92.23Å 180.53Å	Depositor
a, b, c, α , β , γ	90.00° 96.05° 90.00°	Depositor
Resolution (Å)	58.33 - 2.00	Depositor
Resolution (A)	58.33 - 2.00	EDS
% Data completeness	80.3 (58.33-2.00)	Depositor
(in resolution range)	80.4(58.33-2.00)	EDS
R _{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.18 (at 2.00 \text{\AA})$	Xtriage
Refinement program	REFMAC $5.8.0155$	Depositor
R, R_{free}	0.205 , 0.234	Depositor
III, IIIfree	0.205 , 0.234	DCC
R_{free} test set	7736 reflections (4.99%)	wwPDB-VP
Wilson B-factor $(Å^2)$	22.2	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 46.8	EDS
L-test for twinning ²	$ \langle L \rangle = 0.46, \langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17729	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 29.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4360e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NAD, XT0 $\,$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	ond lengths	E	Sond angles
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	1.11	10/2048~(0.5%)	1.22	24/2779~(0.9%)
1	В	0.95	9/2046~(0.4%)	1.00	15/2777~(0.5%)
1	С	1.06	5/2040~(0.2%)	1.01	9/2769~(0.3%)
1	D	0.93	4/2046~(0.2%)	1.02	12/2777~(0.4%)
1	Е	0.91	3/2028~(0.1%)	1.02	15/2751~(0.5%)
1	F	1.03	8/2054~(0.4%)	1.06	18/2788~(0.6%)
1	G	0.93	3/1997~(0.2%)	1.06	14/2709~(0.5%)
1	Н	0.96	3/2003~(0.1%)	1.03	16/2717~(0.6%)
All	All	0.99	45/16262~(0.3%)	1.06	123/22067~(0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2
1	В	0	1
1	С	0	2
All	All	0	5

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	А	219	GLU	CD-OE1	13.10	1.40	1.25
1	С	219	GLU	CD-OE2	10.46	1.37	1.25
1	F	220	GLU	CD-OE2	10.36	1.37	1.25
1	А	219	GLU	CD-OE2	9.54	1.36	1.25
1	А	153	ARG	CZ-NH2	9.47	1.45	1.33

The worst 5 of 123 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	225	ARG	NE-CZ-NH2	-17.24	111.68	120.30
1	А	225	ARG	NE-CZ-NH1	15.36	127.98	120.30
1	А	153	ARG	NE-CZ-NH2	12.87	126.74	120.30
1	G	225	ARG	NE-CZ-NH1	12.53	126.56	120.30
1	Н	225	ARG	NE-CZ-NH1	12.09	126.35	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	106	ASN	Sidechain
1	А	225	ARG	Sidechain
1	В	225	ARG	Sidechain
1	С	211	ALA	Mainchain
1	С	85	GLY	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2004	0	2027	10	0
1	В	2002	0	2023	8	0
1	С	1999	0	2018	14	0
1	D	2002	0	2023	11	0
1	Е	1985	0	2003	15	0
1	F	2007	0	2027	7	0
1	G	1954	0	1980	10	0
1	Н	1960	0	1987	6	0
2	А	44	0	26	1	0
2	В	44	0	26	0	0
2	С	44	0	26	1	0
2	D	44	0	26	1	0
2	Е	44	0	26	1	0
2	F	44	0	26	1	0
2	G	44	0	26	0	0
2	Н	44	0	26	1	0
3	А	27	0	0	0	0
3	В	27	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	27	0	0	0	0
3	D	27	0	0	0	0
3	Ε	27	0	0	0	0
3	F	27	0	0	0	0
3	G	27	0	0	0	0
3	Н	27	0	0	0	0
4	С	1	0	0	0	0
4	Н	1	0	0	0	0
5	A	173	0	0	3	1
5	В	174	0	0	1	0
5	С	160	0	0	3	0
5	D	153	0	0	6	0
5	Е	149	0	0	7	0
5	F	149	0	0	2	1
5	G	136	0	0	2	0
5	Н	152	0	0	0	0
All	All	17729	0	16296	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:106:ASN:ND2	5:G:401:HOH:O	1.98	0.94
1:E:106:ASN:OD1	5:E:401:HOH:O	1.85	0.92
1:G:195:ARG:HA	1:G:199[A]:MET:HE2	1.56	0.88
1:D:106:ASN:ND2	5:D:401:HOH:O	2.05	0.87
1:B:225:ARG:HD2	5:B:475:HOH:O	1.79	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:528:HOH:O	5:F:404:HOH:O[2_656]	2.13	0.07



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	268/289~(93%)	256~(96%)	12~(4%)	0	100	100
1	В	268/289~(93%)	255~(95%)	13~(5%)	0	100	100
1	С	267/289~(92%)	256~(96%)	11 (4%)	0	100	100
1	D	268/289~(93%)	255~(95%)	13~(5%)	0	100	100
1	Ε	263/289~(91%)	250~(95%)	13~(5%)	0	100	100
1	F	269/289~(93%)	257~(96%)	12 (4%)	0	100	100
1	G	258/289~(89%)	246~(95%)	12~(5%)	0	100	100
1	Η	259/289~(90%)	247 (95%)	12 (5%)	0	100	100
All	All	2120/2312~(92%)	2022~(95%)	98~(5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	es
1	А	206/222~(93%)	201~(98%)	5(2%)	49 51	
1	В	206/222~(93%)	200~(97%)	6 (3%)	42 43	
1	С	205/222 (92%)	201 (98%)	4 (2%)	55 58	
1	D	206/222~(93%)	202~(98%)	4 (2%)	57 61	
1	Ε	205/222 (92%)	200~(98%)	5 (2%)	49 51	
1	F	207/222~(93%)	202~(98%)	5(2%)	49 51	



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	G	202/222~(91%)	197~(98%)	5(2%)	47 49
1	Η	203/222~(91%)	200~(98%)	3(2%)	65 69
All	All	1640/1776~(92%)	1603~(98%)	37~(2%)	52 53

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	6	ASP
1	G	162	THR
1	Н	6	ASP
1	G	49	ARG
1	G	68	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	48	GLN
1	Н	214	GLN
1	F	214	GLN
1	А	216	GLN
1	G	214	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Туре	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	gles
	Type	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	XT0	Н	302	-	$30,\!30,\!30$	1.66	7 (23%)	$39,\!41,\!41$	1.49	4(10%)
2	NAD	D	301	-	42,48,48	1.48	2 (4%)	$50,\!73,\!73$	1.18	4 (8%)
3	XT0	G	302	-	$30,\!30,\!30$	1.61	4 (13%)	$39,\!41,\!41$	1.25	<mark>3 (7%)</mark>
2	NAD	В	301	-	42,48,48	0.73	0	50,73,73	1.21	7 (14%)
3	XT0	А	302	-	30,30,30	1.95	8 (26%)	39,41,41	1.58	10(25%)
3	XT0	С	302	-	30,30,30	1.87	7 (23%)	$39,\!41,\!41$	1.91	15 (38%)
2	NAD	Н	301	-	42,48,48	1.01	1 (2%)	50,73,73	1.20	2 (4%)
2	NAD	F	301	-	42,48,48	1.04	1 (2%)	$50,\!73,\!73$	1.32	7 (14%)
3	XT0	Е	302	-	$30,\!30,\!30$	2.42	8 (26%)	$39,\!41,\!41$	1.44	<mark>6 (15%)</mark>
3	XT0	D	302	-	$30,\!30,\!30$	1.58	4 (13%)	$39,\!41,\!41$	1.53	7 (17%)
2	NAD	G	301	-	42,48,48	0.92	3 (7%)	$50,\!73,\!73$	1.01	<mark>3 (6%)</mark>
3	XT0	F	302	-	$30,\!30,\!30$	2.39	7 (23%)	$39,\!41,\!41$	1.45	<mark>5 (12%)</mark>
2	NAD	Е	301	-	42,48,48	1.39	3 (7%)	$50,\!73,\!73$	1.11	5(10%)
2	NAD	С	301	-	42,48,48	0.86	2 (4%)	50,73,73	1.38	3 (6%)
3	XT0	В	302	-	$30,\!30,\!30$	2.31	9 (30%)	$39,\!41,\!41$	2.38	12 (30%)
2	NAD	А	301	-	42,48,48	1.15	3 (7%)	$50,\!73,\!73$	1.18	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
3	XT0	Н	302	-	-	2/14/21/21	0/4/4/4
2	NAD	D	301	-	-	7/26/62/62	0/5/5/5
3	XT0	G	302	-	-	2/14/21/21	0/4/4/4
2	NAD	В	301	-	-	7/26/62/62	0/5/5/5
3	XT0	А	302	-	-	2/14/21/21	0/4/4/4
3	XT0	С	302	-	-	3/14/21/21	0/4/4/4
2	NAD	Н	301	-	-	9/26/62/62	0/5/5/5



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	F	301	-	-	7/26/62/62	0/5/5/5
3	XT0	Е	302	-	-	2/14/21/21	0/4/4/4
3	XT0	D	302	-	-	2/14/21/21	0/4/4/4
2	NAD	G	301	-	-	7/26/62/62	0/5/5/5
3	XT0	F	302	-	-	3/14/21/21	0/4/4/4
2	NAD	Е	301	-	-	7/26/62/62	0/5/5/5
2	NAD	С	301	-	-	6/26/62/62	0/5/5/5
3	XT0	В	302	-	-	3/14/21/21	0/4/4/4
2	NAD	А	301	-	-	7/26/62/62	0/5/5/5

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	Е	302	XT0	NAQ-NAR	-9.04	1.18	1.34
3	F	302	XT0	NAR-NBA	7.84	1.49	1.34
2	D	301	NAD	O7N-C7N	6.87	1.37	1.24
2	Е	301	NAD	C2N-N1N	6.76	1.43	1.35
3	А	302	XT0	CAV-CAC	-6.58	1.34	1.44

The worst 5 of 97 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	В	302	XT0	CAM-CAO-CAZ	-7.44	91.26	104.71
2	С	301	NAD	C3D-C2D-C1D	-6.63	90.99	100.98
3	В	302	XT0	CAO-CAZ-CAW	5.36	124.10	114.28
3	D	302	XT0	CAY-CAV-CAC	5.29	124.73	119.57
3	F	302	XT0	NAQ-NAR-NBA	-5.22	103.38	107.31

There are no chirality outliers.

5 of 76 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
3	Н	302	XT0	CAK-CAW-CAZ-CAN
3	Н	302	XT0	NAQ-CAW-CAZ-CAN
2	D	301	NAD	C5D-O5D-PN-O2N
2	D	301	NAD	O4D-C1D-N1N-C2N
3	G	302	XT0	CAK-CAW-CAZ-CAN


There are no ring outliers.



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	NAD	1	0
2	Н	301	NAD	1	0
2	F	301	NAD	1	0
2	Е	301	NAD	1	0
2	С	301	NAD	1	0
2	А	301	NAD	1	0


6 monomers are involved in 6 short contacts:

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(A^2)$	Q<0.9
1	А	268/289~(92%)	-0.18	1 (0%) 92 92	22, 25, 29, 60	0
1	В	268/289~(92%)	0.03	8 (2%) 50 49	22, 25, 37, 58	0
1	С	268/289~(92%)	-0.18	0 100 100	22, 25, 32, 44	0
1	D	268/289~(92%)	0.04	11 (4%) 37 36	23, 26, 41, 69	0
1	Е	265/289~(91%)	-0.08	4 (1%) 73 72	23, 26, 38, 82	0
1	F	268/289~(92%)	-0.21	3 (1%) 80 79	23, 25, 36, 55	0
1	G	260/289~(89%)	-0.05	5 (1%) 66 65	23, 26, 39, 52	0
1	Н	261/289~(90%)	-0.16	3 (1%) 80 79	23, 25, 35, 54	0
All	All	2126/2312 (91%)	-0.10	35 (1%) 72 70	22, 25, 37, 82	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	THR	5.2
1	D	84	ALA	5.0
1	F	2	THR	4.8
1	Е	2	THR	4.1
1	D	85	GLY	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$\mathbf{Q}{<}0.9$
3	XT0	Н	302	27/27	0.97	0.08	$15,\!18,\!24,\!25$	0
2	NAD	D	301	44/44	0.97	0.07	$10,\!13,\!16,\!17$	0
3	XT0	G	302	27/27	0.97	0.06	12,14,17,18	0
2	NAD	В	301	44/44	0.97	0.07	$9,\!13,\!15,\!16$	0
4	CL	Н	303	1/1	0.97	0.10	$31,\!31,\!31,\!31$	0
3	XT0	Е	302	27/27	0.97	0.07	$11,\!14,\!18,\!20$	0
3	XT0	А	302	27/27	0.98	0.07	$14,\!16,\!19,\!19$	0
2	NAD	Н	301	44/44	0.98	0.08	12,17,20,21	0
2	NAD	F	301	44/44	0.98	0.06	$10,\!14,\!17,\!19$	0
3	XT0	С	302	27/27	0.98	0.07	12,13,21,21	0
3	XT0	D	302	27/27	0.98	0.07	12,14,17,19	0
2	NAD	G	301	44/44	0.98	0.07	$12,\!17,\!19,\!22$	0
3	XT0	F	302	27/27	0.98	0.06	$12,\!13,\!18,\!19$	0
2	NAD	Е	301	44/44	0.98	0.06	11,16,19,21	0
2	NAD	С	301	44/44	0.98	0.07	$10,\!12,\!15,\!16$	0
3	XT0	В	302	27/27	0.98	0.07	$10,\!12,\!16,\!17$	0
2	NAD	А	301	44/44	0.98	0.06	$10,\!13,\!15,\!17$	0
4	CL	С	303	1/1	0.99	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









































6.5 Other polymers (i)

There are no such residues in this entry.

