

Full wwPDB X-ray Structure Validation Report (i)

Aug 10, 2020 - 08:39 AM BST

 Title : Crystal Structure of HIV-1 BG505 SOSIP.664 Prefusion Env Trimer Bou to Small Molecule HIV-1 Entry Inhibitor Compound 484 in Complex with Human Antibodies 3H109L and 35O22 at 3.0 Angstrom Authors : Lai, YT.; Kwong, P.D. 	nd
to Small Molecule HIV-1 Entry Inhibitor Compound 484 in Complex wi Human Antibodies 3H109L and 35O22 at 3.0 Angstrom Authors : Lai, YT.; Kwong, P.D.	
Human Antibodies 3H109L and 35O22 at 3.0 Angstrom Authors : Lai, YT.; Kwong, P.D.	$^{\mathrm{th}}$
Authors : Lai, YT.; Kwong, P.D.	
Deposited on : $2018-10-20$	
${\rm Resolution} : 2.50 \ {\rm \AA}({\rm reported})$	

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	÷	4.02b-467
Mogul		1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries}, { m resolution\ range}({ m \AA}))$		
R _{free}	130704	4661 (2.50-2.50)		
Clashscore	141614	$5346\ (2.50-2.50)$		
Ramachandran outliers	138981	5231(2.50-2.50)		
Sidechain outliers	138945	5233 (2.50-2.50)		
RSRZ outliers	127900	4559(2.50-2.50)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	В	153	2% 68%	15%	16%
		100	18%		10/0
2	D	134	66%	27%	• •
3	Е	114	<u>6%</u> 69%	21%	• 8%
			2%		
4	G	481	66%	25%	• 9%
5	Н	244	67%	25%	• 7%
6	L	217	% 72%	23%	• •



Mol	Chain	Length		Quality of chain	
7	А	2		100%	
7	J	2		100%	
7	K	2	50%		50%
7	N	2		100%	
8	С	6	33%	50%	17%
9	F	3		100%	
9	Ι	3	33%	67%	
10	М	10	20%	70%	10%



2 Entry composition (i)

There are 13 unique types of molecules in this entry. The entry contains 10157 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	В	128	Total 1018	$\begin{array}{c} \mathrm{C} \\ 647 \end{array}$	N 175	O 190	S 6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	559	PRO	ILE	engineered mutation	UNP Q2N0S6
В	605	CYS	THR	engineered mutation	UNP Q2N0S6

• Molecule 2 is a protein called 35O22 scFv heavy chain portion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	128	Total 994	C 628	N 169	O 192	${ m S}{ m 5}$	0	0	0

• Molecule 3 is a protein called 35O22 scFv light chain portion.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	105	Total 805	$ m C \ 506$	N 133	O 160	${ m S}{ m 6}$	0	0	0

• Molecule 4 is a protein called Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	439	Total 3461	C 2177	N 613	O 643	S 28	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	137	ALA	ASN	engineered mutation	UNP Q2N0S6
G	332	ASN	THR	conflict	UNP Q2N0S6



Chain	Residue	Modelled	Actual	Comment	Reference
G	501	CYS	ALA	conflict	UNP Q2N0S6
G	509	ARG	-	expression tag	UNP Q2N0S6
G	510	ARG	-	expression tag	UNP Q2N0S6
G	511	ARG	-	expression tag	UNP Q2N0S6
G	512	ARG	-	expression tag	UNP Q2N0S6
G	513	ARG	-	expression tag	UNP Q2N0S6

• Molecule 5 is a protein called 3H109L Fab heavy chain.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
5	Н	226	Total	С	Ν	Ο	S	0	0	0
			1715	1093	278	338	6		0	

• Molecule 6 is a protein called 3H109L Fab light chain.

Mol	Chain	Residues		Ate	\mathbf{oms}			ZeroOcc	AltConf	Trace
6	L	211	Total 1604	C 1009	N 276	0 312	S 7	0	0	0

• Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
7	Λ	2	Total C N O	0	0	0
	<u>ک</u>	28 16 2 10	0	0	0	
7	Т	9	Total C N O	0	0	0
4	1 1	2	28 16 2 10	0	0	0
7	K	2	Total C N O	0	0	0
1	17		28 16 2 10	0		
7	7 N	2	Total C N O	0	0	0
7			28 16 2 10			U

• Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyran ose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
8	С	6	Total 72	C 40	N 2	O 30	0	0	0

• Molecule 9 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
9	F	3	Total C N 39 22 2	O 15	0	0	0
9	Ι	3	Total C N 39 22 2	O 15	0	0	0

• Molecule 10 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyra nose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyr anose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyr anose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-g lucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
10	М	10	Total 116	C N 64 2	NO 250)	0	0	0

• Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	В	1	Total C N O 14 8 1 5	0	0
11	В	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0
11	G	1	Total C N O 14 8 1 5	0	0

• Molecule 12 is {4-[1-(3-chlorophenyl)cyclopropane-1-carbonyl]piperazin-1-yl}(thiophen-3 -yl)methanone (three-letter code: JYJ) (formula: C₁₉H₁₉ClN₂O₂S) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
12	G	1	Total 25	C 19	Cl 1	N 2	О 2	${ m S}$	0	0

• Molecule 13 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	В	1	Total O 1 1	0	0
13	G	1	Total O 1 1	0	0
13	L	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Envelope glycoprotein gp160





• Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain A:

100%

NAG 1 NAG 2

• Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain J:		100%	
MAG 1 NAG 2			
• Molecule 7: opyranose	2-acetamido-2-deoxy-beta	a-D-glucopyranose-(1-4)-2-acetamic	lo-2-deoxy-beta-D-gluc
Chain K:	50%	50%	•
NAG 1 NAG 2			
• Molecule 7: opyranose	2-acetamido-2-deoxy-beta	a-D-glucopyranose-(1-4)-2-acetamic	lo-2-deoxy-beta-D-gluc
Chain N:		100%	•
NAG 1 NAG 2			
• Malaanla 9.	- l- h - D	$(1,2)$ alaba D $\dots \dots $	

• Molecule 8: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:	33%	50%	17%
NAG1 NAG2 BMA3 MAN4 MAN5 MAN6 MAN6			

• Molecule 9: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:	100%		
NAG1 BMA2 BMA3			
• Moloculo 0: bota D mannopur	nnoso(1.4) 2 ncotan	nida 2 daavy hata D al	$u_{convrance} (1.4)$

• Molecule 9: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:	33%	67%

NAG1 NAG2 BMAG

 $\label{eq:solution} \bullet \mbox{ Molecule 10: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)] alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-g lucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranoy-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetam$



Chain M:	20%	70%	10%
NAG1 NAG2 BNA3 BNA3 MAN4 MAN4 MAN5 MAN5 MAN5 MAN3 MAN3 MAN10 MAN10			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants	131.50Å 131.50 Å 315.74 Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	43.04 - 2.50	Depositor
Resolution (A)	43.04 - 2.48	EDS
% Data completeness	39.1(43.04-2.50)	Depositor
(in resolution range)	38.2(43.04-2.48)	EDS
R _{merge}	0.13	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.66 (at 2.48 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
B B.	0.223 , 0.263	Depositor
Π, Π_{free}	0.223 , 0.263	DCC
R_{free} test set	2077 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor ($Å^2$)	39.5	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.27 , 31.2	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.057 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	10157	wwPDB-VP
Average B, all atoms $(Å^2)$	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.15% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, JYJ, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5
1	В	0.23	0/1036	0.41	0/1404
2	D	0.25	0/1021	0.47	0/1390
3	Е	0.25	0/829	0.48	0/1133
4	G	0.25	0/3532	0.46	0/4791
5	Н	0.25	0/1758	0.48	0/2397
6	L	0.24	0/1647	0.45	0/2247
All	All	0.25	0/9823	0.46	0/13362

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1018	0	1007	18	0
2	D	994	0	952	20	0
3	Е	805	0	752	13	0
4	G	3461	0	3408	72	0
5	Н	1715	0	1687	35	0
6	L	1604	0	1553	32	0
7	А	28	0	25	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	J	28	0	25	0	0
7	K	28	0	25	0	0
7	N	28	0	25	0	0
8	С	72	0	61	1	0
9	F	39	0	34	0	0
9	Ι	39	0	34	1	0
10	М	116	0	97	1	0
11	В	28	0	26	1	0
11	G	126	0	117	1	0
12	G	25	0	0	0	0
13	В	1	0	0	0	0
13	G	1	0	0	0	0
13	L	1	0	0	0	0
All	All	10157	0	9828	184	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (184) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Λ)	Clash overlap $(Å)$
4.C.202.VAL.HC23	4.C.449.ILE.HB	1.66	$\frac{0.78}{0.78}$
4.G.232.VIID.HG23	4.G.977.ILE.HG12	1.00	0.77
5·H·137·CLV·HA3	$5 \cdot \text{H} \cdot 179 \cdot \text{V} \Delta \text{L} \cdot \text{H} \text{C} 12$	1.00	0.77
6.I. 83.CI U.OF1	6.L.167.LVS.NZ	2.12	0.72
4.C.163.THR.HC23	4.C.165.I FU.H	1.56	0.71
6.1.34.CI N.HC3	4.G.105.DEU.II 6·L·40·TVR·HΔ	1.50	0.70
1.B.617.ABC.HH22	$11 \cdot B \cdot 702 \cdot N \Delta C \cdot H 4$	1.75	0.70
$2 \cdot D \cdot 72 (F) \cdot THP \cdot HC 23$	2.D.72(H).DHE.H	1.09	0.08
4.C.358.II F.HC12	4.C.306.II F.HA	1.00	0.00
4.G.358.ILE.IIG12 4.C.477.ASP.OD1	4.G.390.11E.11A	2.30	0.05
9.D.51.II F.UD11	2.D.71.TUD.UC22	2.30	0.05
	2:D:71:1 III.IIG25	1.79	0.03
	2:D:97:LEU:HG	1.70	0.04
0:L:13:VAL:HG22		1.80	0.03
5:E:54:ARG:NHI	5:E:58:1LE:U	2.31	0.63
D:H:140:GLU:HG3	5:H:147:PRO:HA	1.81	0.62
5:H:24:VAL:HB	5:H:76:ASN:HB3	1.81	0.62
3:E:29:SER:HA	3:E:66:LY S:HZ1	1.64	0.62
4:G:09:TKP:HZ3	4:G:108:1LE:HG23	1.05	0.61
3:E:85:THR:HG22	3:E:103:LYS:HG2	1.81	0.61
4:G:136:ASN:HB3	4:G:151:ARG:HE	1.65	0.61



6MTN

		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
5:H:63:LEU:HD13	5:H:67:VAL:HG21	1.83	0.61	
6:L:185:GLN:HA	6:L:188:MET:HG2	1.83	0.61	
1:B:585:ARG:NH2	4:G:491:ILE:O	2.34	0.60	
4:G:476:ARG:HA	4:G:479:TRP:CD1	2.37	0.60	
6:L:19:ALA:HB3	6:L:75:ILE:HB	1.84	0.60	
4:G:93:PHE:HB2	4:G:233:PHE:HZ	1.66	0.60	
2:D:57:LYS:HE3	2:D:69:MET:HG3	1.83	0.59	
6:L:83:GLU:HG3	6:L:106:VAL:HG12	1.82	0.59	
2:D:72(F):THR:HG22	2:D:73:THR:HB	1.85	0.59	
6:L:119:PHE:HB2	6:L:134:VAL:HG13	1.84	0.58	
4:G:426:MET:HG2	4:G:428:GLN:H	1.68	0.58	
4:G:321(A):ASP:OD1	4:G:322:ILE:N	2.36	0.58	
6:L:34:GLN:NE2	6:L:49:TYR:O	2.37	0.57	
8:C:2:NAG:H83	8:C:2:NAG:H3	1.87	0.56	
6:L:37:GLN:NE2	6:L:86:TYR:OH	2.38	0.56	
6:L:83:GLU:HG2	6:L:105:THR:HA	1.88	0.56	
4:G:360:ARG:HB2	4:G:394:THR:HG22	1.88	0.56	
4:G:350:ARG:NH1	4:G:357:THR:O	2.39	0.55	
1:B:629:LEU:HD23	4:G:44:VAL:HG23	1.87	0.55	
2:D:4:LEU:HG	2:D:24:THR:HG22	1.87	0.55	
2:D:94:LYS:HG2	2:D:102:LEU:HB3	1.89	0.55	
4:G:90:THR:HG22	4:G:240:PRO:HA	1.89	0.55	
4:G:376:PHE:CE2	4:G:378:CYS:HB2	2.42	0.55	
6:L:50:ASN:O	6:L:52:GLN:N	2.40	0.54	
5:H:6:GLU:HG3	5:H:22:CYS:SG	2.47	0.54	
4:G:361:PHE:HB3	4:G:391:PHE:HB3	1.90	0.54	
4:G:387:THR:HG22	4:G:390:LEU:HD12	1.89	0.54	
3:E:19:VAL:HG23	3:E:78:LEU:HD11	1.89	0.54	
3:E:47:ILE:HD12	3:E:58:ILE:HD12	1.89	0.53	
5:H:139:LEU:HG	5:H:141:LYS:HG3	1.91	0.53	
2:D:9:ALA:HB1	2:D:108:LEU:HB2	1.91	0.52	
5:H:163:THR:HG23	5:H:178:SER:HB2	1.91	0.52	
2:D:37:ILE:HD12	2:D:103:TRP:CH2	2.45	0.52	
4:G:101:VAL:HG13	4:G:479:TRP:HB2	1.91	0.52	
5:H:36:TRP:HB3	5:H:48:ILE:HD12	1.92	0.52	
6:L:22:SER:HA	6:L:72:THR:HG22	1.92	0.52	
4:G:47:ASP:N	4:G:47:ASP:OD1	2.43	0.51	
6:L:37:GLN:HB2	6:L:47:LEU:HD11	1.92	0.51	
6:L:80:ALA:HA	6:L:106:VAL:HG11	1.92	0.51	
2:D:82(A):ARG:O	2:D:82(C):LEU:N	2.44	0.51	
1:B:604:CYS:SG	4:G:38:VAL:HB	2.51	0.51	



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
5:H:33:TYR:CD2	5:H:52:SEB:HA	2.45	0.51	
4:G:456:ARG:HD2	4:G:466:GLU:HG2	1.93	0.50	
5:H:138:CYS:SG	5:H:194:CYS:HB2	2.51	0.50	
5:H:36:TRP:CG	5:H:80:LEU:HD12	2.47	0.50	
4:G:134:VAL:HG22	4:G:135:THR:H	1.77	0.50	
5:H:191:THR:HB	5:H:208:LYS:HD2	1.93	0.50	
1:B:610:TRP:CD2	4:G:498:PRO:HB3	2.47	0.49	
4:G:257:THR:O	4:G:259:LEU:N	2.43	0.49	
4:G:297:THR:HB	4:G:444:ARG:HG3	1.94	0.49	
5:H:35:SER:HB3	5:H:47:TRP:HE1	1.78	0.49	
6:L:39:ARG:HG2	6:L:84:ALA:HB2	1.95	0.49	
6:L:86:TYR:HB2	6:L:102:THR:HG23	1.95	0.49	
3:E:28:CYS:HB3	3:E:66:LYS:HZ3	1.78	0.48	
4:G:370:GLU:HG3	4:G:384:TYR:HE1	1.79	0.48	
4:G:299:PRO:HG2	4:G:327:ARG:HB2	1.96	0.48	
3:E:4:LEU:HB2	3:E:27(B):VAL:HG11	1.95	0.48	
3:E:38:TRP:O	3:E:85:THR:OG1	2.31	0.48	
2:D:37:ILE:HD11	2:D:100(F):PRO:HG2	1.96	0.47	
3:E:11:VAL:HG23	3:E:104:VAL:HG12	1.96	0.47	
4:G:358:ILE:O	4:G:465:THR:HA	2.13	0.47	
5:H:53:ASP:OD1	5:H:54:SER:N	2.45	0.47	
6:L:61:ARG:NH1	6:L:82:ASP:OD2	2.47	0.47	
3:E:38:TRP:CE2	3:E:44:PRO:HG3	2.50	0.47	
5:H:6:GLU:N	5:H:6:GLU:OE1	2.46	0.47	
1:B:569:THR:HG23	1:B:572:GLY:H	1.80	0.47	
4:G:86:LEU:HD21	4:G:244:THR:HG22	1.97	0.47	
4:G:358:ILE:HG23	4:G:395:TRP:O	2.14	0.47	
4:G:42:VAL:HG23	4:G:44:VAL:HG12	1.97	0.47	
4:G:474:ASP:OD1	4:G:476:ARG:HG2	2.14	0.47	
4:G:484:TYR:CE2	4:G:485:LYS:HG2	2.51	0.46	
4:G:32:GLU:OE1	4:G:32:GLU:N	2.49	0.46	
5:H:194:CYS:O	5:H:206:ASP:HB2	2.16	0.46	
10:M:1:NAG:H83	10:M:1:NAG:H3	1.97	0.46	
4:G:257:THR:HG22	4:G:258:GLN:HG3	1.98	0.46	
3:E:49:TYR:CD2	3:E:50:GLU:HG2	2.50	0.45	
5:H:94:ARG:NH2	5:H:100(Q):ASP:OD2	2.38	0.45	
5:H:33:TYR:HB2	5:H:95:ALA:O	2.17	0.45	
6:L:197:THR:HA	6:L:202:THR:HA	1.98	0.45	
1:B:567:LYS:HD3	1:B:567:LYS:HA	1.81	0.45	
4:G:139:THR:HG23	4:G:150:MET:HB2	1.98	0.45	
4:G:376:PHE:HE2	4:G:378:CYS:HB2	1.80	0.45	



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
4:G:384:TYR:CE2	4:G:421:LYS:HB2	2.50	0.45
1:B:593:LEU:HD13	4:G:494:LEU:HD21	1.98	0.45
4:G:205:CYS:O	4:G:208:VAL:HG22	2.17	0.45
5:H:87:SER:HB3	5:H:109:VAL:H	1.81	0.45
2:D:13:LYS:HD2	2:D:13:LYS:HA	1.64	0.44
2:D:88:GLY:O	2:D:109:LEU:HB2	2.17	0.44
4:G:234:ASN:HD21	11:G:613:NAG:C7	2.30	0.44
4:G:91:GLU:OE1	4:G:487:LYS:NZ	2.46	0.44
4:G:129:LEU:HD13	4:G:192:ARG:HA	1.99	0.44
4:G:201:ILE:HD11	4:G:435:TYR:HB2	1.98	0.44
6:L:35:TRP:CE2	6:L:73:LEU:HB2	2.52	0.44
5:H:144:PHE:HA	5:H:145:PRO:HA	1.72	0.44
4:G:298:ARG:NH2	4:G:441:GLY:O	2.50	0.44
4:G:101:VAL:HG21	4:G:480:ARG:HG2	1.98	0.44
4:G:298:ARG:NH1	4:G:381:GLU:OE2	2.50	0.44
4:G:175:LEU:HB2	4:G:320:THR:HB	1.99	0.44
5:H:18:LEU:HD11	5:H:107:VAL:HG21	2.00	0.44
4:G:206:PRO:HG3	4:G:318:TYR:CE2	2.53	0.44
1:B:537:LEU:HB2	1:B:602:LEU:HD23	1.99	0.44
4:G:358:ILE:HD11	4:G:396:ILE:HG23	1.99	0.44
4:G:98:ASN:OD1	4:G:99:ASN:N	2.50	0.44
2:D:63:PHE:HB3	2:D:67:VAL:HG21	2.00	0.43
1:B:608:VAL:HG11	1:B:645:LEU:HB2	1.99	0.43
4:G:69:TRP:HB2	4:G:71:THR:HG23	2.00	0.43
4:G:135:THR:OG1	6:L:94:ARG:HD2	2.18	0.43
1:B:549:VAL:HG12	4:G:53:PHE:CE1	2.53	0.43
1:B:537:LEU:CB	1:B:602:LEU:HD23	2.49	0.43
6:L:125:GLU:OE1	6:L:132:THR:N	2.52	0.43
6:L:170:ASN:HB2	6:L:172:LYS:HG3	1.99	0.43
4:G:123:THR:N	4:G:124:PRO:HD2	2.34	0.43
4:G:335:LYS:H	4:G:413:SER:HA	1.83	0.43
3:E:28:CYS:HB3	3:E:66:LYS:NZ	2.33	0.43
6:L:181:LEU:HD11	6:L:186:TRP:HB2	2.01	0.43
4:G:193:LEU:HB2	4:G:196:CYS:SG	2.59	0.43
5:H:103:LYS:HD2	5:H:103:LYS:N	2.34	0.43
6:L:34:GLN:HB2	6:L:89:HIS:HB3	2.01	0.43
5:H:188:GLY:HA2	5:H:189:THR:HA	1.61	0.42
2:D:35:ASN:OD1	2:D:47:TRP:NE1	2.45	0.42
5:H:150:VAL:HA	5:H:195:ASN:O	2.19	0.42
4:G:276:ASN:ND2	4:G:279:ASN:HB2	2.34	0.42
6:L:13:VAL:HG21	6:L:78:VAL:HG21	2.01	0.42



	ne pagem	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:523:LEU:O	4:G:86:LEU:HD22	2.20	0.42	
4:G:333:VAL:HG11	4:G:390:LEU:HD21	2.01	0.42	
4:G:260:LEU:HD21	4:G:453:ILE:HD11	2.00	0.42	
6:L:131:ALA:HB3	6:L:181:LEU:HD12	2.00	0.42	
4:G:259:LEU:HD13	4:G:449:ILE:HD13	2.01	0.42	
1:B:614:TRP:HH2	1:B:645:LEU:HD11	1.84	0.42	
4:G:105:HIS:CE1	4:G:109:ILE:HD11	2.55	0.42	
3:E:11:VAL:O	3:E:104:VAL:HA	2.19	0.41	
4:G:305:LYS:HE3	4:G:321:GLY:HA2	2.02	0.41	
4:G:346:VAL:HG23	4:G:359:ILE:HD11	2.02	0.41	
6:L:37:GLN:OE1	6:L:45:ILE:HD11	2.20	0.41	
6:L:54:ARG:HD2	6:L:58:ILE:HG22	2.00	0.41	
6:L:18:THR:HG23	6:L:76:SER:HA	2.02	0.41	
2:D:52:SER:C	2:D:53:TYR:H	2.22	0.41	
2:D:47:TRP:CZ2	2:D:49:GLY:HA2	2.55	0.41	
6:L:145:VAL:HG12	6:L:198:HIS:HB2	2.03	0.41	
1:B:655:LYS:HE3	1:B:655:LYS:HB2	1.80	0.41	
5:H:194:CYS:SG	5:H:196:VAL:HG23	2.61	0.41	
6:L:90:MET:O	6:L:95(C):SER:OG	2.28	0.41	
5:H:43:LYS:HD2	6:L:6:SER:N	2.34	0.41	
2:D:20:ILE:HD11	2:D:109:LEU:HD11	2.03	0.41	
2:D:36:TRP:CZ2	2:D:78:ALA:HB1	2.56	0.41	
4:G:384:TYR:HE2	4:G:421:LYS:HB2	1.85	0.41	
5:H:143:TYR:O	5:H:174:TYR:N	2.53	0.41	
5:H:11:LEU:HG	5:H:145:PRO:HG3	2.01	0.41	
5:H:83:THR:C	5:H:109:VAL:HG11	2.41	0.41	
5:H:4:LEU:HB3	5:H:102:GLY:HA2	2.03	0.41	
5:H:36:TRP:CZ3	5:H:92:CYS:HB3	2.55	0.41	
5:H:51:ILE:HD11	5:H:69:ILE:HB	2.02	0.41	
1:B:525:ALA:HB1	1:B:528:SER:HB2	2.03	0.41	
4:G:93:PHE:HB2	4:G:233:PHE:CZ	2.50	0.41	
5:H:204:LYS:HB2	5:H:204:LYS:HE2	1.73	0.41	
5:H:59:TYR:HD2	5:H:64:LYS:HD2	1.85	0.41	
1:B:593:LEU:HA	1:B:593:LEU:HD12	1.91	0.40	
5:H:100(P):MET:SD	5:H:100(P):MET:N	2.94	0.40	
4:G:212:PRO:HG3	4:G:254:VAL:HG22	2.03	0.40	
1:B:592:LEU:HA	1:B:595:ILE:HG22	2.04	0.40	
4:G:163:THR:HG22	4:G:168:LYS:O	2.21	0.40	
9:I:1:NAG:H61	9:I:2:NAG:N2	2.36	0.40	
2:D:72(C):VAL:HB	2:D:75:THR:OG1	2.22	0.40	
4:G:272:ILE:HG23	4:G:284:ILE:HG23	2.02	0.40	



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perc	$\mathbf{entiles}$
1	В	124/153~(81%)	117 (94%)	6~(5%)	1 (1%)	19	35
2	D	126/134~(94%)	111 (88%)	14 (11%)	1 (1%)	19	35
3	Ε	103/114~(90%)	82~(80%)	20 (19%)	1 (1%)	15	28
4	G	429/481~(89%)	401 (94%)	26~(6%)	2(0%)	29	48
5	Η	222/244~(91%)	202~(91%)	19~(9%)	1 (0%)	29	48
6	L	209/217~(96%)	196~(94%)	12~(6%)	1 (0%)	29	48
All	All	1213/1343~(90%)	1109 (91%)	97 (8%)	7 (1%)	25	43

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	G	397	SER
2	D	82(B)	ASN
3	Е	52	ASN
6	L	51	ASN
1	В	548	ILE
4	G	138	ILE
5	Н	117	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	В	110/129~(85%)	107~(97%)	3 (3%)	44 71
2	D	107/112~(96%)	101 (94%)	6 (6%)	21 40
3	Ε	92/100~(92%)	83 (90%)	9 (10%)	8 15
4	G	391/427~(92%)	377~(96%)	14 (4%)	35 61
5	Н	196/212~(92%)	189~(96%)	7 (4%)	35 61
6	L	175/181~(97%)	167~(95%)	8 (5%)	27 50
All	All	1071/1161~(92%)	1024 (96%)	47 (4%)	28 52

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	519	PHE
1	В	574	LYS
1	В	645	LEU
2	D	35	ASN
2	D	69	MET
2	D	82(A)	ARG
2	D	85	ASP
2	D	98	ARG
2	D	105	GLN
3	Е	6	GLN
3	Е	11	VAL
3	Е	27(C)	CYS
3	Е	34	SER
3	Е	47	ILE
3	Е	61	ARG
3	Ε	84	THR
3	Е	89	CYS
3	Е	96	CYS
4	G	47	ASP
4	G	54	CYS
4	G	107	ASP
4	G	139	THR
4	G	155	LYS
4	G	164	GLU
4	G	203	GLN
4	G	246	GLN
4	G	309	ILE
4	G	356	ASN
4	G	412	ASP
4	G	447	SER



Mol	Chain	Res	Type
4	G	466	GLU
4	G	469	ARG
5	Н	22	CYS
5	Н	38	ARG
5	Н	63	LEU
5	Н	75	LYS
5	Н	100(F)	SER
5	Н	136	LEU
5	Н	173	LEU
6	L	61	ARG
6	L	95(B)	PHE
6	L	102	THR
6	L	103	ARG
6	L	134	VAL
6	L	170	ASN
6	L	172	LYS
6	L	206	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	G	246	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

30 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



Mal	Trees	Chain	Dec	Tink	Bond lengths		Bond angles			
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	А	1	2,7	14, 14, 15	0.33	0	$17,\!19,\!21$	0.41	0
7	NAG	А	2	7	14,14,15	0.21	0	$17,\!19,\!21$	0.55	0
8	NAG	С	1	8,4	14,14,15	0.27	0	17,19,21	0.55	0
8	NAG	С	2	8	14,14,15	0.36	0	$17,\!19,\!21$	1.30	2 (11%)
8	BMA	С	3	8	11,11,12	0.74	0	$15,\!15,\!17$	0.79	0
8	MAN	С	4	8	11,11,12	1.17	2 (18%)	$15,\!15,\!17$	1.67	4 (26%)
8	MAN	С	5	8	11,11,12	1.56	2 (18%)	$15,\!15,\!17$	2.24	4 (26%)
8	MAN	С	6	8	11,11,12	0.75	0	$15,\!15,\!17$	1.03	2 (13%)
9	NAG	F	1	9,4	14,14,15	0.41	0	$17,\!19,\!21$	0.50	0
9	NAG	F	2	9	14,14,15	0.24	0	$17,\!19,\!21$	0.50	0
9	BMA	F	3	9	11,11,12	0.65	0	$15,\!15,\!17$	0.70	0
9	NAG	Ι	1	9,4	14,14,15	0.32	0	17,19,21	0.53	0
9	NAG	Ι	2	9	14,14,15	0.28	0	17, 19, 21	0.55	0
9	BMA	I	3	9	11,11,12	0.54	0	$15,\!15,\!17$	0.73	0
7	NAG	J	1	4,7	14,14,15	0.39	0	17,19,21	0.45	0
7	NAG	J	2	7	14,14,15	0.27	0	17,19,21	0.42	0
7	NAG	K	1	4,7	14,14,15	0.32	0	17,19,21	0.49	0
7	NAG	K	2	7	14,14,15	0.25	0	$17,\!19,\!21$	0.59	1 (5%)
10	NAG	М	1	10,4	14,14,15	0.34	0	$17,\!19,\!21$	1.46	2 (11%)
10	MAN	М	10	10	11,11,12	1.12	1(9%)	$15,\!15,\!17$	1.19	2 (13%)
10	NAG	М	2	10	14,14,15	0.34	0	17,19,21	0.43	0
10	BMA	М	3	10	11,11,12	0.83	0	$15,\!15,\!17$	0.94	0
10	MAN	М	4	10	11,11,12	0.74	1 (9%)	$15,\!15,\!17$	1.34	2 (13%)
10	MAN	М	5	10	11,11,12	0.64	0	$15,\!15,\!17$	0.96	1 (6%)
10	MAN	М	6	10	11,11,12	0.74	0	$15,\!15,\!17$	0.95	2 (13%)
10	MAN	М	7	10	11,11,12	0.82	0	$15,\!15,\!17$	0.96	1(6%)
10	MAN	М	8	10	11,11,12	0.71	0	$15,\!15,\!17$	1.04	1 (6%)
10	MAN	М	9	10	11,11,12	0.70	0	$15,\!15,\!17$	1.43	2(13%)
7	NAG	N	1	4,7	14,14,15	0.21	0	17,19,21	0.64	0
7	NAG	N	2	7	14,14,15	0.19	0	$17,\!19,\!21$	0.61	0

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



OTAL TTA	6	М	Л	ľ	I
----------	---	---	---	---	---

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	А	1	2,7	-	4/6/23/26	0/1/1/1
7	NAG	А	2	7	-	4/6/23/26	0/1/1/1
8	NAG	С	1	8,4	_	0/6/23/26	0/1/1/1
8	NAG	С	2	8	-	5/6/23/26	0/1/1/1
8	BMA	С	3	8	-	2/2/19/22	0/1/1/1
8	MAN	С	4	8	-	0/2/19/22	0/1/1/1
8	MAN	С	5	8	-	1/2/19/22	0/1/1/1
8	MAN	С	6	8	_	0/2/19/22	0/1/1/1
9	NAG	F	1	9,4	-	2/6/23/26	0/1/1/1
9	NAG	F	2	9	-	0/6/23/26	0/1/1/1
9	BMA	F	3	9	-	0/2/19/22	0/1/1/1
9	NAG	Ι	1	9,4	-	2/6/23/26	0/1/1/1
9	NAG	Ι	2	9	-	2/6/23/26	0/1/1/1
9	BMA	Ι	3	9	-	1/2/19/22	0/1/1/1
7	NAG	J	1	4,7	-	2/6/23/26	0/1/1/1
7	NAG	J	2	7	-	2/6/23/26	0/1/1/1
7	NAG	K	1	4,7	_	2/6/23/26	0/1/1/1
7	NAG	K	2	7	-	2/6/23/26	0/1/1/1
10	NAG	М	1	10,4	-	3/6/23/26	0/1/1/1
10	MAN	М	10	10	-	0/2/19/22	0/1/1/1
10	NAG	М	2	10	-	2/6/23/26	0/1/1/1
10	BMA	М	3	10	_	0/2/19/22	0/1/1/1
10	MAN	М	4	10	-	2/2/19/22	0/1/1/1
10	MAN	М	5	10	-	0/2/19/22	0/1/1/1
10	MAN	М	6	10	-	0/2/19/22	0/1/1/1
10	MAN	М	7	10	-	0/2/19/22	0/1/1/1
10	MAN	М	8	10	-	0/2/19/22	0/1/1/1
10	MAN	М	9	10	-	2/2/19/22	0/1/1/1
7	NAG	N	1	4,7	-	0/6/23/26	0/1/1/1
7	NAG	N	2	7	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\operatorname{\AA})$
8	С	5	MAN	C1-C2	3.91	1.61	1.52
8	С	5	MAN	O5-C1	2.98	1.48	1.43
8	С	4	MAN	C2-C3	2.51	1.56	1.52
8	С	4	MAN	C1-C2	2.50	1.57	1.52
10	М	4	MAN	C1-C2	2.10	1.57	1.52
10	М	10	MAN	C2-C3	2.03	1.55	1.52



6M	ΤN

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
8	С	5	MAN	C1-O5-C5	6.70	121.27	112.19
10	М	9	MAN	C1-O5-C5	4.64	118.48	112.19
10	М	1	NAG	C2-N2-C7	4.58	129.43	122.90
8	С	4	MAN	C1-C2-C3	4.42	115.11	109.67
8	С	2	NAG	C2-N2-C7	4.32	129.05	122.90
10	М	4	MAN	C1-O5-C5	3.66	117.16	112.19
8	С	5	MAN	O5-C1-C2	3.31	115.88	110.77
8	С	5	MAN	C1-C2-C3	3.02	113.38	109.67
10	М	1	NAG	C1-C2-N2	2.71	115.11	110.49
10	М	8	MAN	C1-O5-C5	2.64	115.76	112.19
8	С	6	MAN	C1-O5-C5	2.62	115.74	112.19
10	М	10	MAN	O2-C2-C3	-2.30	105.54	110.14
10	М	5	MAN	O2-C2-C3	-2.29	105.55	110.14
8	С	4	MAN	C2-C3-C4	2.28	114.84	110.89
10	М	4	MAN	O2-C2-C3	-2.28	105.57	110.14
10	М	9	MAN	O2-C2-C3	-2.24	105.64	110.14
8	С	5	MAN	O2-C2-C3	-2.24	105.65	110.14
8	С	6	MAN	O2-C2-C3	-2.24	105.66	110.14
8	С	4	MAN	O2-C2-C3	-2.20	105.73	110.14
10	М	6	MAN	O2-C2-C3	-2.18	105.78	110.14
8	С	4	MAN	C1-O5-C5	2.14	115.09	112.19
10	М	6	MAN	C1-O5-C5	2.13	115.08	112.19
10	М	10	MAN	C1-C2-C3	2.12	112.27	109.67
10	М	7	MAN	O2-C2-C3	-2.09	105.95	110.14
7	K	2	NAG	C1-O5-C5	2.03	114.94	112.19
8	С	2	NAG	C1-C2-N2	2.02	113.95	110.49

All (26) bond angle outliers are listed below:

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
8	С	3	BMA	O5-C5-C6-O6
7	J	1	NAG	O5-C5-C6-O6
10	М	2	NAG	O5-C5-C6-O6
8	С	3	BMA	C4-C5-C6-O6
8	С	2	NAG	O5-C5-C6-O6
7	А	1	NAG	O5-C5-C6-O6
7	J	2	NAG	O5-C5-C6-O6
7	Κ	2	NAG	O5-C5-C6-O6
7	А	1	NAG	C4-C5-C6-O6
7	J	1	NAG	C4-C5-C6-O6



Mol	Chain	Res	Type	Atoms
8	С	2	NAG	C8-C7-N2-C2
8	С	2	NAG	O7-C7-N2-C2
10	М	1	NAG	C8-C7-N2-C2
10	М	1	NAG	O7-C7-N2-C2
7	K	1	NAG	O5-C5-C6-O6
10	М	2	NAG	C4-C5-C6-O6
7	J	2	NAG	C4-C5-C6-O6
9	Ι	1	NAG	O5-C5-C6-O6
8	С	2	NAG	C4-C5-C6-O6
9	Ι	1	NAG	C4-C5-C6-O6
10	М	9	MAN	O5-C5-C6-O6
10	М	9	MAN	C4-C5-C6-O6
7	K	1	NAG	C4-C5-C6-O6
7	А	1	NAG	C1-C2-N2-C7
9	F	1	NAG	C4-C5-C6-O6
10	М	4	MAN	O5-C5-C6-O6
7	K	2	NAG	C4-C5-C6-O6
8	С	5	MAN	O5-C5-C6-O6
9	Ι	3	BMA	O5-C5-C6-O6
10	М	4	MAN	C4-C5-C6-O6
9	Ι	2	NAG	C4-C5-C6-O6
9	Ι	2	NAG	O5-C5-C6-O6
9	F	1	NAG	O5-C5-C6-O6
7	A	2	NAG	C1-C2-N2-C7
7	А	2	NAG	C4-C5-C6-O6
7	A	1	NAG	C3-C2-N2-C7
10	М	1	NAG	C3-C2-N2-C7
7	A	2	NAG	C3-C2-N2-C7
8	С	2	NAG	C3-C2-N2-C7
7	А	2	NAG	O5-C5-C6-O6

Continued from previous page...

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	Ι	1	NAG	1	0
9	Ι	2	NAG	1	0
8	С	2	NAG	1	0
10	М	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





















5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Tune	Chain	Dog	Tink	Bo	ond leng	$_{\rm sths}$	B	ond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	NAG	G	607	4	14,14,15	0.26	0	17,19,21	0.56	0
11	NAG	G	635	4	14,14,15	0.19	0	17,19,21	0.47	0
11	NAG	В	701	1	14,14,15	0.35	0	17,19,21	0.56	0
11	NAG	G	617	4	14,14,15	0.22	0	17,19,21	0.45	0
11	NAG	G	632	4	14,14,15	0.27	0	17,19,21	0.47	0
11	NAG	G	633	4	14,14,15	0.25	0	17,19,21	0.50	0
11	NAG	G	634	4	14,14,15	0.33	0	17,19,21	0.65	1 (5%)
12	JYJ	G	638	-	27,28,28	1.30	2 (7%)	33,41,41	2.36	7 (21%)
11	NAG	G	611	4	14,14,15	0.34	0	17,19,21	0.47	0
11	NAG	G	612	4	14,14,15	0.28	0	17,19,21	0.47	0
11	NAG	В	702	1	14,14,15	0.24	0	17,19,21	0.45	0
11	NAG	G	613	4	14,14,15	0.30	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
11	NAG	G	607	4	-	2/6/23/26	0/1/1/1
11	NAG	G	635	4	-	0/6/23/26	0/1/1/1
11	NAG	В	701	1	-	2/6/23/26	0/1/1/1
11	NAG	G	617	4	-	2/6/23/26	0/1/1/1
11	NAG	G	632	4	-	2/6/23/26	0/1/1/1
11	NAG	G	633	4	-	0/6/23/26	0/1/1/1
11	NAG	G	634	4	-	2/6/23/26	0/1/1/1
12	JYJ	G	638	-	-	10/22/38/38	0/4/4/4
11	NAG	G	611	4	-	2/6/23/26	0/1/1/1
11	NAG	G	612	4	-	2/6/23/26	0/1/1/1
11	NAG	В	702	1	-	2/6/23/26	0/1/1/1
11	NAG	G	613	4	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
12	G	638	JYJ	C12-C9	5.22	1.40	1.37
12	G	638	JYJ	C17-CL	2.40	1.79	1.74

All (8) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
12	G	638	JYJ	C9-C12-S	-9.12	106.41	112.29
12	G	638	JYJ	C13-C2-C3	5.72	121.77	113.91
12	G	638	JYJ	C-C2-C1	4.34	60.16	58.33
12	G	638	JYJ	C5-C4-N	-2.95	104.12	110.44
12	G	638	JYJ	C6-C7-N	-2.92	104.18	110.44
12	G	638	JYJ	C11-S-C12	2.46	97.38	92.37
11	G	634	NAG	C1-O5-C5	2.29	115.29	112.19
12	G	638	JYJ	C-C1-C2	-2.03	59.86	60.84

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
12	G	638	JYJ	O1-C3-N-C4
12	G	638	JYJ	O1-C3-N-C7
12	G	638	JYJ	O-C8-C9-C10
12	G	638	JYJ	O-C8-C9-C12
12	G	638	JYJ	C9-C8-N1-C5
11	В	702	NAG	C4-C5-C6-O6
11	В	702	NAG	O5-C5-C6-O6
12	G	638	JYJ	O-C8-N1-C5
11	G	613	NAG	C4-C5-C6-O6
11	В	701	NAG	O5-C5-C6-O6
11	G	634	NAG	O5-C5-C6-O6
11	G	617	NAG	O5-C5-C6-O6
11	В	701	NAG	C4-C5-C6-O6
11	G	611	NAG	O5-C5-C6-O6
11	G	634	NAG	C4-C5-C6-O6
11	G	611	NAG	C4-C5-C6-O6
11	G	632	NAG	O5-C5-C6-O6
11	G	613	NAG	O5-C5-C6-O6
12	G	638	JYJ	O-C8-N1-C6
12	G	638	JYJ	C9-C8-N1-C6
11	G	617	NAG	C4-C5-C6-O6
11	G	612	NAG	O5-C5-C6-O6
11	G	632	NAG	C4-C5-C6-O6
11	G	607	NAG	O5-C5-C6-O6
11	G	607	NAG	C4-C5-C6-O6
12	G	638	JYJ	C2-C3-N-C4
12	G	638	JYJ	C2-C3-N-C7
11	G	612	NAG	C4-C5-C6-O6

All (28) torsion outliers are listed below:

There are no ring outliers.



Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	В	702	NAG	1	0
11	G	613	NAG	1	0

2 monomers are involved in 2 short contacts:

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	$Q{<}0.9$
1	В	128/153~(83%)	-0.23	3 (2%) 60 63	12, 38, 64, 84	0
2	D	128/134~(95%)	0.58	24 (18%) 1 1	38, 76, 107, 119	0
3	Е	105/114~(92%)	0.20	7 (6%) 17 18	40, 64, 97, 109	0
4	G	439/481~(91%)	-0.19	11 (2%) 57 61	13, 34, 83, 134	0
5	Н	226/244~(92%)	0.04	8 (3%) 44 47	24, 54, 84, 114	0
6	L	211/217~(97%)	-0.48	2 (0%) 84 86	14, 38, 57, 107	0
All	All	1237/1343~(92%)	-0.09	55 (4%) 34 37	12, 44, 94, 134	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	Н	187	LEU	7.5
4	G	138	ILE	7.2
4	G	141	ASP	6.8
4	G	137	ALA	6.4
4	G	458	GLY	5.9
4	G	188	ASN	4.8
2	D	18	VAL	4.5
3	Ε	78	LEU	4.0
2	D	82(C)	LEU	4.0
2	D	82	ILE	3.9
5	Н	189	THR	3.9
6	L	7	TYR	3.8
3	Ε	8	ALA	3.8
3	Е	80	PRO	3.8
4	G	140	ASP	3.7
2	D	14	PRO	3.5
2	D	88	GLY	3.4
3	Е	15	LEU	3.3
2	D	108	LEU	3.3



Mol	Chain	Res	Type	RSRZ
2	D	10	THR	3.3
4	G	81	PRO	3.2
3	Е	76	SER	3.2
2	D	110	THR	3.1
2	D	65	ASP	3.1
2	D	109	LEU	3.1
2	D	43	ARG	3.0
2	D	84	SER	2.8
2	D	61	PRO	2.8
2	D	1	GLN	2.8
4	G	398	ASN	2.8
2	D	80	MET	2.5
2	D	41	ALA	2.5
5	Н	209	VAL	2.5
1	В	549	VAL	2.5
2	D	89	THR	2.5
4	G	185	ASN	2.5
2	D	59	LEU	2.4
6	L	6	SER	2.4
2	D	20	ILE	2.4
4	G	396	ILE	2.4
1	В	547	GLY	2.3
3	Е	11	VAL	2.3
2	D	85	ASP	2.3
5	Η	157	LEU	2.3
5	Н	182	VAL	2.3
1	В	548	ILE	2.2
2	D	13	LYS	2.2
5	H	71	VAL	2.2
5	Н	125	SER	2.2
2	D	42	GLY	2.2
5	H	78	LEU	2.2
2	D	87	THR	2.2
3	Е	79	ARG	2.1
4	G	60	ALA	2.0
2	2 D		SER	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
9	BMA	F	3	11/12	0.68	0.24	$92,\!113,\!122,\!124$	0
7	NAG	А	1	14/15	0.75	0.23	77,107,124,125	0
7	NAG	А	2	14/15	0.78	0.37	72,112,123,123	0
7	NAG	J	2	14/15	0.82	0.26	$65,\!96,\!118,\!126$	0
10	MAN	М	9	11/12	0.83	0.32	72,84,99,107	0
8	MAN	С	4	11/12	0.86	0.18	77,79,90,94	0
9	BMA	Ι	3	11/12	0.87	0.21	$64,\!83,\!94,\!95$	0
7	NAG	K	2	14/15	0.90	0.24	$53,\!83,\!100,\!101$	0
8	MAN	С	5	11/12	0.90	0.20	$84,\!90,\!105,\!105$	0
9	NAG	F	2	14/15	0.91	0.15	$50,\!88,\!106,\!119$	0
7	NAG	N	2	14/15	0.92	0.19	$90,\!95,\!100,\!103$	0
10	NAG	М	1	14/15	0.93	0.14	40,52,62,72	0
10	MAN	М	7	11/12	0.93	0.12	$46,\!58,\!70,\!71$	0
9	NAG	Ι	2	14/15	0.93	0.12	$25,\!53,\!68,\!84$	0
7	NAG	N	1	14/15	0.94	0.11	39,67,87,93	0
10	MAN	М	8	11/12	0.94	0.11	$45,\!56,\!65,\!70$	0
10	MAN	М	6	11/12	0.94	0.11	$30,\!53,\!56,\!57$	0
9	NAG	F	1	14/15	0.95	0.12	24,52,67,73	0
8	NAG	С	1	14/15	0.95	0.17	$18,\!33,\!45,\!48$	0
7	NAG	J	1	14/15	0.95	0.12	$32,\!61,\!73,\!91$	0
10	MAN	М	10	11/12	0.95	0.09	$51,\!64,\!76,\!77$	0
10	BMA	М	3	11/12	0.96	0.10	$34,\!41,\!48,\!55$	0
10	NAG	М	2	14/15	0.96	0.09	$10,\!52,\!62,\!64$	0
8	NAG	С	2	14/15	0.96	0.12	$35,\!44,\!57,\!61$	0
8	BMA	С	3	11/12	0.97	0.10	$40,\!50,\!60,\!77$	0
8	MAN	С	6	11/12	0.97	0.09	$23,\!41,\!53,\!60$	0
10	MAN	М	4	11/12	0.97	0.10	$2\overline{4,\!31,\!46,\!54}$	0
10	MAN	M	5	11/12	0.97	0.14	$21,\!34,\!44,\!47$	0
7	NAG	K	1	14/15	0.98	0.11	$26,\!33,\!49,\!58$	0
9	NAG	Ι	1	14/15	0.98	0.11	18,28,37,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
11	NAG	G	607	14/15	0.77	0.20	76, 96, 117, 126	0
11	NAG	В	701	14/15	0.86	0.39	59,108,116,128	0
11	NAG	В	702	14/15	0.87	0.41	$69,\!84,\!107,\!110$	0
11	NAG	G	632	14/15	0.89	0.21	$74,\!90,\!100,\!102$	0
11	NAG	G	635	14/15	0.90	0.26	$41,\!94,\!100,\!106$	0
11	NAG	G	612	14/15	0.91	0.13	$35,\!53,\!74,\!76$	0
11	NAG	G	617	14/15	0.92	0.25	$53,\!72,\!96,\!98$	0
11	NAG	G	611	14/15	0.92	0.17	33,54,73,98	0
11	NAG	G	613	14/15	0.92	0.13	62,78,91,94	0
12	JYJ	G	638	25/25	0.93	0.15	$15,\!44,\!61,\!64$	0
11	NAG	G	633	14/15	0.94	0.14	49,68,75,79	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q < 0.9
11	NAG	G	634	14/15	0.95	0.12	$36,\!51,\!59,\!82$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

