



Full wwPDB NMR Structure Validation Report ⓘ

Jun 3, 2023 – 05:10 PM EDT

PDB ID : 2MVS
BMRB ID : 25220
Title : N6-Methyladenosine RNA
Authors : Lynch, S.R.; Kool, E.
Deposited on : 2014-10-10

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 25%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashescore	158937	12864
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	1	10	
1	2	10	

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 646 atoms, of which 224 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called N-6_Methyl_Adenosine_RNA.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	1	10	323	96	112	38	68	9	0
1	2	10	323	96	112	38	68	9	0

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: N-6_Methyl_Adenosine_RNA

Chain 1:  90% 10%

G1	G2	A3	C4	U5	A6	G7	U8	C9	C10
----	----	----	----	----	----	----	----	----	-----

- Molecule 1: N-6_Methyl_Adenosine_RNA

Chain 2:  90% 10%

G11	G12	A13	C14	U15	A16	G17	U18	C19	C20
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 30 calculated structures, 1 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.36
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	120
Number of shifts mapped to atoms	120
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	25%

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
6MZ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	1	211	112	112	22
1	2	211	112	112	19
All	All	422	224	224	41

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 63.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:1:5:U:O2'	1:1:6:A:H5'	0.93	1.63
1:2:13:6MZ:H2'	1:2:14:C:C6	0.90	2.02
1:1:7:G:O2'	1:1:8:U:H5'	0.89	1.68
1:2:17:G:O2'	1:2:18:U:H5'	0.74	1.82
1:1:6:A:H2'	1:1:7:G:C8	0.63	2.29
1:1:3:6MZ:O2'	1:1:4:C:H5'	0.61	1.95
1:1:8:U:O2'	1:1:9:C:H5'	0.59	1.98
1:1:3:6MZ:H2'	1:1:4:C:C6	0.58	2.34
1:2:13:6MZ:H2'	1:2:14:C:H6	0.57	1.53
1:1:5:U:H2'	1:1:6:A:C8	0.57	2.34
1:1:7:G:O2'	1:1:8:U:C5'	0.57	2.50
1:2:11:G:H2'	1:2:12:G:C8	0.57	2.34

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:1:5:U:HO2'	1:1:6:A:H5'	0.55	1.60
1:1:5:U:O2'	1:1:6:A:C5'	0.54	2.48
1:2:13:6MZ:C2	1:2:14:C:C2	0.54	2.91
1:2:12:G:H2'	1:2:13:6MZ:C8	0.53	2.34
1:2:18:U:H2'	1:2:19:C:C6	0.53	2.39
1:1:9:C:H2'	1:1:10:C:C6	0.52	2.39
1:1:2:G:H2'	1:1:3:6MZ:C8	0.52	2.35
1:1:9:C:O2'	1:1:10:C:H5'	0.50	2.07
1:1:5:U:H2'	1:1:6:A:H8	0.49	1.68
1:2:19:C:H2'	1:2:20:C:C6	0.48	2.44
1:1:6:A:C6	1:1:7:G:C6	0.48	3.01
1:2:11:G:C6	1:2:12:G:C6	0.48	3.01
1:1:7:G:HO2'	1:1:8:U:H5'	0.48	1.67
1:2:18:U:O2'	1:2:19:C:H5'	0.47	2.10
1:1:8:U:H2'	1:1:9:C:C6	0.47	2.44
1:1:4:C:H2'	1:1:5:U:C6	0.44	2.46
1:1:3:6MZ:N7	1:1:3:6MZ:C9	0.44	2.80
1:1:1:G:H2'	1:1:2:G:C8	0.44	2.47
1:1:7:G:H2'	1:1:8:U:C6	0.44	2.48
1:2:13:6MZ:N7	1:2:13:6MZ:C9	0.44	2.81
1:2:19:C:O2'	1:2:20:C:H5'	0.44	2.13
1:2:16:A:C6	1:2:17:G:C6	0.44	3.06
1:2:11:G:N1	1:2:12:G:C6	0.44	2.86
1:2:12:G:C6	1:2:13:6MZ:C6	0.44	3.01
1:2:16:A:H2'	1:2:17:G:C8	0.43	2.48
1:2:11:G:C2	1:2:12:G:C4	0.41	3.09
1:2:15:U:H2'	1:2:16:A:O4'	0.40	2.16
1:2:17:G:H2'	1:2:18:U:C6	0.40	2.51
1:1:3:6MZ:O2'	1:1:4:C:C5'	0.40	2.68

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	1	8/10 (80%)	0 (0%)	0 (0%)	0.28
1	2	8/10 (80%)	0 (0%)	0 (0%)	0.25
All	All	16/20 (80%)	0 (0%)	0 (0%)	0.26

The overall RNA backbone suiteness is 0.26.

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	6MZ	2	13	1	18,25,26	1.24	3 (16%)
1	6MZ	1	3	1	18,25,26	1.25	3 (16%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	6MZ	2	13	1	16,36,39	1.32	2 (12%)
1	6MZ	1	3	1	16,36,39	1.31	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	6MZ	2	13	1	-	0,5,27,28	0,3,3,3
1	6MZ	1	3	1	-	0,5,27,28	0,3,3,3

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1	3	6MZ	C9-N6	3.72	1.38	1.45
1	2	13	6MZ	C9-N6	3.67	1.38	1.45
1	2	13	6MZ	O4'-C1'	2.22	1.44	1.41
1	1	3	6MZ	O4'-C1'	2.18	1.44	1.41
1	1	3	6MZ	C8-N7	2.13	1.30	1.34
1	2	13	6MZ	C8-N7	2.12	1.30	1.34

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	2	13	6MZ	C9-N6-C6	3.34	119.99	122.87
1	1	3	6MZ	C9-N6-C6	3.32	120.02	122.87
1	1	3	6MZ	C2-N1-C6	2.94	119.11	116.59
1	2	13	6MZ	C2-N1-C6	2.93	119.11	116.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 25% for the well-defined parts and 25% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	120
Number of shifts mapped to atoms	120
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 25%, i.e. 85 atoms were assigned a chemical shift out of a possible 340. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Sugar	44/198 (22%)	35/108 (32%)	9/90 (10%)	0/0 (—%)
Base	41/142 (29%)	28/88 (32%)	13/30 (43%)	0/24 (0%)
Overall	85/340 (25%)	63/196 (32%)	22/120 (18%)	0/24 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 25%, i.e. 85 atoms were assigned a chemical shift out of a possible 340. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Sugar	44/198 (22%)	35/108 (32%)	9/90 (10%)	0/0 (—%)
Base	41/142 (29%)	28/88 (32%)	13/30 (43%)	0/24 (0%)
Overall	85/340 (25%)	63/196 (32%)	22/120 (18%)	0/24 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	383
Intra-residue ($ i-j =0$)	178
Sequential ($ i-j =1$)	145
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range ($ i-j \geq 5$)	0
Inter-chain	60
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	172
Number of unmapped restraints	0
Number of restraints per residue	27.8
Number of long range restraints per residue ¹	0.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	9.0	0.19
0.2-0.5 (Medium)	8.0	0.45
>0.5 (Large)	3.0	0.93

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	178	46.5	3	1.7	0.8	3	1.7	0.8
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	178	46.5	3	1.7	0.8	3	1.7	0.8
Sequential ($i-j =1$)	145	37.9	6	4.1	1.6	6	4.1	1.6
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	145	37.9	6	4.1	1.6	6	4.1	1.6
Medium range ($i-j >1$ & $i-j <5$)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range ($i-j \geq 5$)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	60	15.7	11	18.3	2.9	11	18.3	2.9
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	60	15.7	11	18.3	2.9	11	18.3	2.9
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	383	100.0	20	5.2	5.2	20	5.2	5.2
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	383	100.0	20	5.2	5.2	20	5.2	5.2

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	3	6	0	0	11	20	0.3	0.93	0.2	0.22

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 363(IR:175, SQ:139, MR:0, LR:0, IC:49) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
3	6	0	0	11	20	1	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)




9.4 Most violated distance restraints in the ensemble [i](#)

No violations found

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,14)	1:2:14:C:H42	1:1:6:A:H8	1	0.93
(1,25)	1:1:7:G:O6	1:2:14:C:H41	1	0.62
(2,244)	1:2:14:C:H1'	1:2:15:U:H6	1	0.55
(2,252)	1:2:14:C:H3'	1:2:15:U:H5	1	0.45
(2,232)	1:2:13:6MZ:H2'	1:2:14:C:H6	1	0.4
(1,21)	1:1:6:A:H61	1:2:15:U:O4	1	0.36
(2,90)	1:1:5:U:H2'	1:1:6:A:H8	1	0.32
(2,185)	1:1:4:C:H42	1:2:16:A:H8	1	0.25
(1,15)	1:2:17:G:O6	1:1:4:C:H41	1	0.25
(1,20)	1:1:6:A:N1	1:2:15:U:H3	1	0.23
(2,287)	1:2:16:A:H5'	1:2:16:A:H1'	1	0.21
(2,69)	1:1:3:6MZ:H8	1:1:4:C:H6	1	0.19
(1,4)	1:1:1:G:O6	1:2:20:C:H41	1	0.19
(1,28)	1:2:13:6MZ:H6	1:1:8:U:O4	1	0.19
(1,23)	1:1:7:G:H21	1:2:14:C:O2	1	0.18
(1,34)	1:2:11:G:H21	1:1:10:C:O2	1	0.13

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,333)	1:2:19:C:H3'	1:2:19:C:H6	1	0.12
(2,161)	1:1:9:C:H3'	1:1:9:C:H6	1	0.12
(2,76)	1:1:4:C:H2'	1:1:5:U:H5	1	0.11
(1,11)	1:1:3:6MZ:H6	1:2:18:U:O4	1	0.11

10 Dihedral-angle violation analysis


10.1 Summary of dihedral-angle violations

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
NU1	20	11.6	0	0.0	0.0	0	0.0	0.0
BETA	18	10.5	0	0.0	0.0	0	0.0	0.0
NU0	20	11.6	0	0.0	0.0	0	0.0	0.0
NU3	20	11.6	0	0.0	0.0	0	0.0	0.0
EPSILON	18	10.5	0	0.0	0.0	0	0.0	0.0
NU2	20	11.6	0	0.0	0.0	0	0.0	0.0
GAMMA	20	11.6	0	0.0	0.0	0	0.0	0.0
CHI	20	11.6	0	0.0	0.0	0	0.0	0.0
.	16	9.3	0	0.0	0.0	0	0.0	0.0
Total	172	100.0	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

No violations found

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

No violations found

10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

No violations found

10.5 All violated dihedral-angle restraints [i](#)

No violations found