

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 4, 2023 – 03:14 AM EDT

PDB ID : 6MYQ

Title: Avian mitochondrial complex II with ferulenol bound

Authors: Berry, E.A.; Huang, L.-S.

Deposited on : 2018-11-02

Resolution : 1.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	621	83%	15% •
2	В	252	5% 81%	13% 5%
3	С	140	87%	12% •
4	D	103	<del>5%</del> 84%	14% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	MOH	A	1011	-	-	-	X
6	Y3P	A	1002	-	X	X	-
8	UNL	A	1027	-	-	-	X
8	UNL	С	1021	-	-	-	X
8	UNL	D	203	-	-	X	-
8	UNL	D	207	-	-	-	X
8	UNL	D	212	-	-	-	X



# 2 Entry composition (i)

There are 18 unique types of molecules in this entry. The entry contains 9426 atoms, of which 2 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	611	Total 4723	C 2956	N 843	O 895	S 29	0	1	0

• Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues		$\mathbf{At}$	oms			ZeroOcc	AltConf	Trace
2	В	239	Total 1921	C 1215	N 325	O 359	S 22	0	0	0

• Molecule 3 is a protein called Succinate dehydrogenase cytochrome b, large subunit.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3	С	139	Total 1076	C 707	N 178	O 186	S 5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
С	139	ALA	SER	conflict	UNP D0VWW3
С	140	MET	GLU	conflict	UNP D0VWW3

• Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
4	D	102	Total 771	C 508	N 122	O 138	S 3	0	0	0


• Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).





Mo	l Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 53	C 27		O 15	P 2	0	0

• Molecule 6 is ( $\{Z\}$ )-2-oxidanylbut-2-enedioic acid (three-letter code: Y3P) (formula:  $C_4H_4O_5$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total 11	C 4	H 2	O 5	0	0

• Molecule 7 is POTASSIUM ION (three-letter code: K) (formula: K).




$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total K 1 1	0	0
7	В	1	Total K 1 1	0	0

• Molecule 8 is UNKNOWN LIGAND (three-letter code: UNL) (formula: ).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	21	Total C Mn N O S 58 14 1 17 24 2	0	0
8	В	15	Total C N O S 42 9 12 20 1	0	1
8	С	19	Total C N O S 78 39 5 32 2	0	0
8	D	10	Total C O P 59 29 29 1	0	0


 $\bullet \ \, \text{Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$)}. \\$ 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total 5	C 3	O 2	0	0


 $\bullet$  Molecule 10 is METHANOL (three-letter code: MOH) (formula:  $\mathrm{CH_{4}O}).$ 





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total C C 2 1 1	)	0	0


 $\bullet \ \ Molecule \ 11 \ is \ FE2/S2 \ (INORGANIC) \ CLUSTER \ (three-letter \ code: \ FES) \ (formula: \ Fe_2S_2).$ 



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
11	В	1	Total 4	Fe 2	S 2	0	0


 $\bullet$  Molecule 12 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe\_4S\_4).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
12	В	1	Total 8	Fe 4	S 4	0	0


• Molecule 13 is FE3-S4 CLUSTER (three-letter code: F3S) (formula:  $Fe_3S_4$ ).



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
13	В	1	Total 7	Fe 3	S 4	0	0

• Molecule 14 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
14	C	1	Total	С	Fe	N	О	0	0
1.1		1	41	32	1	4	4		

• Molecule 15 is 4-oxidanyl-3-[(2 {E},6 {E})-3,7,11-trimethyldodeca-2,6,10-trienyl]chrom en-2-one (three-letter code: 9AU) (formula:  $C_{24}H_{30}O_3$ ) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
15	С	1	Total 19	C 16	O 3	0	0


 $\bullet$  Molecule 16 is UNDECYL-MALTOSIDE (three-letter code: UMQ) (formula:  $\mathrm{C}_{23}\mathrm{H}_{44}\mathrm{O}_{11}).$ 





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
16	С	1	Total 34	C 23	O 11	0	0

 $\bullet$  Molecule 17 is 1,2-Distear oyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula:  $\rm C_{41}H_{82}NO_8P).$ 



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
17	D	1	Total 42	C 32	N 1	O 8	P 1	0	0

• Molecule 18 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
18	A	1	Total O 2 2	0	1
18	A	234	Total O 237 237	0	3
18	В	145	Total O 146 146	0	1
18	С	1	Total O 2 2	0	1
18	С	1	Total O 1 1	0	0
18	С	51	Total O 51 51	0	0
18	D	31	Total O 31 31	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial



• Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial




• Molecule 3: Succinate dehydrogenase cytochrome b, large subunit



• Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	69.91Å 84.55Å 290.52Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.62 - 1.97	Depositor
rtesolution (A)	29.62 - 1.97	EDS
% Data completeness	93.7 (29.62-1.97)	Depositor
(in resolution range)	82.9 (29.62-1.97)	EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.48 (at 1.96Å)	Xtriage
Refinement program	PHENIX dev_3150	Depositor
P. P.	0.208 , 0.239	Depositor
$R, R_{free}$	0.209 , 0.240	DCC
$R_{free}$ test set	2334 reflections $(2.02\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.5	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.30 , 48.7	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.45, < L^2>=0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9426	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: Y3P, 3PE, SF4, UMQ, F3S, PEG, K, 9AU, MLZ, UNL, FAD, FES, HEM, MOH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.41	0/4824	0.60	0/6531	
2	В	0.43	0/1963	0.58	0/2646	
3	С	0.38	0/1094	0.54	0/1486	
4	D	0.38	0/794	0.50	0/1089	
All	All	0.41	0/8675	0.58	0/11752	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4723	0	4608	67	0
2	В	1921	0	1913	21	0
3	С	1076	0	1116	12	0
4	D	771	0	763	13	0
5	A	53	0	29	5	0
6	A	9	2	0	6	0
7	A	1	0	0	0	0
7	В	1	0	0	0	0
8	A	58	0	0	3	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	В	42	0	0	3	0
8	С	78	0	0	1	0
8	D	59	0	0	3	0
9	A	5	0	5	3	0
10	A	2	0	0	1	0
11	В	4	0	0	0	0
12	В	8	0	0	0	0
13	В	7	0	0	0	0
14	С	41	0	24	0	0
15	С	19	0	0	0	0
16	С	34	0	44	0	0
17	D	42	0	61	2	0
18	A	239	0	0	9	0
18	В	146	0	0	3	0
18	С	54	0	0	0	0
18	D	31	0	0	1	0
All	All	9424	2	8563	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
1:A:297:ARG:HH22	6:A:1002:Y3P:C3	1.67	1.07	
1:A:187:GLU:HG2	1:A:390:PRO:HB2	1.65	0.77	
3:C:118:GLN:OE1	8:C:1019:UNL:O	2.11	0.68	
4:D:85:TYR:OH	8:D:212:UNL:O	2.12	0.67	
1:A:101:GLU:OE2	18:A:1101:HOH:O	2.13	0.67	

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	A	610/621~(98%)	590 (97%)	20 (3%)	0	100	100	
2	В	$237/252\ (94\%)$	232 (98%)	5 (2%)	0	100	100	
3	$\mathbf{C}$	136/140~(97%)	135 (99%)	1 (1%)	0	100	100	
4	D	$100/103\ (97\%)$	98 (98%)	2 (2%)	0	100	100	
All	All	$1083/1116 \ (97\%)$	1055 (97%)	28 (3%)	0	100	100	

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	otameric Outliers		Percentiles		
1	A	496/506~(98%)	489 (99%)	7 (1%)	67	62		
2	В	$215/219 \ (98\%)$	211 (98%)	4 (2%)	57	50		
3	C	116/117 (99%)	116 (100%)	0	100	100		
4	D	78/79 (99%)	75 (96%)	3 (4%)	33	21		
All	All	905/921~(98%)	891 (98%)	14 (2%)	65	59		

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type	
2	В	63	ARG	
2	В	66	ARG	
4	D	101	TRP	
4	D	14	ARG	
4	D	19	LEU	

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:



Mol	Chain	$\operatorname{Res}$	Type
2	В	150	GLN
3	С	118	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
						Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
Ī	3	MLZ	С	15	3	8,9,10	0.61	0	4,9,11	1.74	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	$\mathbf{Type}$	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
3	MLZ	С	15	3	-	2/7/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$	
3	С	15	MLZ	CM-NZ-CE	3.29	121.45	111.95	

There are no chirality outliers.

All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	С	15	MLZ	CG-CD-CE-NZ
3	С	15	MLZ	CE-CD-CG-CB

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 79 ligands modelled in this entry, 2 are monoatomic and 66 are unknown - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
14	HEM	С	201	4,3	40,48,50	1.44	6 (15%)	46,80,82	1.73	10 (21%)
17	3PE	D	201	-	41,41,50	1.09	2 (4%)	44,46,55	1.09	4 (9%)
6	Y3P	A	1002	-	8,8,8	2.65	4 (50%)	9,10,10	2.46	4 (44%)
9	PEG	A	1005	-	4,4,6	0.59	0	3,3,5	0.12	0
10	MOH	A	1011	-	1,1,1	0.07	0	-		
11	FES	В	1001	2	0,4,4	-	-	-		
16	UMQ	С	1002	-	35,35,35	0.55	0	46,46,46	0.91	2 (4%)
13	F3S	В	1003	2	0,9,9	-	-	-		
12	SF4	В	1002	2	0,12,12	-	-	-		
5	FAD	A	1001	1	53,58,58	0.80	1 (1%)	68,89,89	1.33	7 (10%)
15	9AU	С	202	-	20,20,28	2.95	8 (40%)	24,27,37	2.08	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	HEM	С	201	4,3	-	3/10/50/54	-
17	3PE	D	201	-	-	16/45/45/54	-
6	Y3P	A	1002	-	-	4/8/8/8	-
9	PEG	A	1005	-	-	0/2/2/4	-
11	FES	В	1001	2	-	-	0/1/1/1
16	UMQ	С	1002	-	-	8/20/60/60	0/2/2/2
13	F3S	В	1003	2	-	-	0/3/3/3
12	SF4	В	1002	2	-	-	0/6/5/5
5	FAD	A	1001	1	-	5/30/50/50	0/6/6/6
15	9AU	С	202	-	-	3/8/8/17	0/2/2/2

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
15	С	202	9AU	OAS-CAZ	7.69	1.50	1.38
6	A	1002	Y3P	C2-C1	5.59	1.55	1.48
15	С	202	9AU	CAW-CAX	5.55	1.48	1.36
15	С	202	9AU	OAS-CAY	5.13	1.45	1.38
17	D	201	3PE	O21-C21	4.59	1.47	1.34

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
15	С	202	9AU	CAZ-OAS-CAY	-4.73	117.62	122.23
14	С	201	HEM	C4B-CHC-C1C	4.33	128.28	122.56
6	A	1002	Y3P	O2-C1-C2	-4.31	115.11	120.08
6	A	1002	Y3P	O1-C1-C2	4.09	119.85	115.91
5	A	1001	FAD	C4-N3-C2	-3.78	118.66	125.64

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1001	FAD	N10-C1'-C2'-O2'
5	A	1001	FAD	N10-C1'-C2'-C3'
6	A	1002	Y3P	O1-C1-C2-O3
6	A	1002	Y3P	O2-C1-C2-O3
17	D	201	3PE	C1-O11-P-O14

There are no ring outliers.


5 monomers are involved in 17 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
17	D	201	3PE	2	0
6	A	1002	Y3P	6	0
9	A	1005	PEG	3	0
10	A	1011	MOH	1	0
5	A	1001	FAD	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















# 5.7 Other polymers (i)

There are no such residues in this entry.



# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	611/621 (98%)	0.31	20 (3%) 46 49	32, 51, 75, 101	0
2	В	239/252 (94%)	0.34	13 (5%) 25 28	31, 42, 71, 89	0
3	С	138/140 (98%)	0.52	10 (7%) 15 17	35, 48, 74, 81	0
4	D	102/103 (99%)	0.44	5 (4%) 29 32	36, 48, 66, 86	0
All	All	1090/1116 (97%)	0.36	48 (4%) 34 36	31, 49, 73, 101	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	С	79	LEU	7.6
3	С	81	PRO	4.5
4	D	30	TYR	4.2
1	A	567	LEU	4.2
2	В	25	GLY	4.1

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	MLZ	С	15	10/11	0.94	0.12	50,59,65,65	0

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ m \AA}^2)$	Q<0.9
8	UNL	D	207	12/-	0.58	0.43	75,81,89,90	0
8	UNL	D	211	1/-	0.61	0.18	61,61,61,61	0
8	UNL	С	1021	1/-	0.64	0.47	60,60,60,60	0
8	UNL	С	1022	1/-	0.65	0.18	66,66,66,66	0
8	UNL	A	1023	1/-	0.70	0.20	72,72,72,72	0
8	UNL	В	1016	1/-	0.70	0.18	59,59,59,59	0
17	3PE	D	201	42/51	0.70	0.28	66,82,115,119	0
8	UNL	A	1012	2/-	0.72	0.23	70,70,70,71	0
8	UNL	С	1005	15/-	0.73	0.39	52,72,78,78	15
8	UNL	С	1018	1/-	0.74	0.17	57,57,57,57	0
8	UNL	A	1027	1/-	0.75	0.48	49,49,49,49	1
8	UNL	A	1025	1/-	0.76	0.20	73,73,73,73	0
8	UNL	В	1009	3/-	0.76	0.40	42,42,43,46	3
8	UNL	D	212	1/-	0.77	0.68	63,63,63,63	0
8	UNL	D	206	5/-	0.78	0.37	74,74,79,79	0
8	UNL	В	1017	1/-	0.79	0.18	47,47,47,47	0
8	UNL	С	1007	5/-	0.79	0.19	54,55,60,62	0
10	MOH	A	1011	2/2	0.80	0.41	79,79,79,81	0
8	UNL	В	1012	2/-	0.80	0.38	85,85,85,86	0
8	UNL	С	1006	12/-	0.81	0.16	63,73,75,75	0
8	UNL	A	1020	3/-	0.81	0.26	39,39,46,53	0
8	UNL	С	1010	15/-	0.81	0.41	74,76,93,94	15
8	UNL	В	1014	4/-	0.82	0.23	89,91,91,92	0
8	UNL	С	1009	4/-	0.82	0.24	75,75,76,76	0
9	PEG	A	1005	5/7	0.83	0.40	67,67,72,74	0
8	UNL	A	1021	3/-	0.83	0.21	44,44,60,64	0
8	UNL	A	1022	1/-	0.83	0.18	76,76,76,76	0
8	UNL	A	1017	2/-	0.84	0.57	80,80,80,82	0
8	UNL	В	1006	3/-	0.84	0.20	64,64,66,67	0
8	UNL	В	1007	3/-	0.84	0.13	58,58,65,65	0
8	UNL	A	1010	4/-	0.84	0.15	68,68,72,80	0
8	UNL	В	1011	4/-	0.84	0.16	69,73,74,78	0
15	9AU	С	202	19/27	0.84	0.26	38,57,67,69	0
8	UNL	A	1016	2/-	0.84	0.13	64,64,64,64	0
8	UNL	В	1013	4/-	0.85	0.21	45,46,47,50	4
8	UNL	С	1003	3/-	0.85	0.19	47,47,52,56	0
8	UNL	D	205	1/-	0.85	0.20	65,65,65,65	0

Continued on next page...




Continued from previous page...

Mol	Type	m previou Chain	$ hootnote{Res}$	Atoms	RSCC	RSR	$\operatorname{B-factors}(  ext{\AA}^2 )$	Q<0.9
8	UNL	A	1018	2/-	0.85	0.33	53,53,53,60	0
8	UNL	A	1013	3/-	0.85	0.26	37,37,37,49	3
8	UNL	D	208	12/-	0.85	0.23	72,76,79,80	0
8	UNL	С	1016	1/-	0.86	0.21	55,55,55,55	0
8	UNL	D	203	11/-	0.86	0.17	56,74,78,79	0
8	UNL	С	1017	1/-	0.86	0.15	54,54,54,54	0
8	UNL	D	210	1/-	0.87	0.10	45,45,45,45	0
8	UNL	С	1015	1/-	0.88	0.13	44,44,44,44	0
8	UNL	С	1019	1/-	0.88	0.19	53,53,53,53	0
8	UNL	С	1008	3/-	0.88	0.28	51,51,59,60	0
8	UNL	С	1011	4/-	0.88	0.28	67,67,71,80	0
8	UNL	В	1010	6/-	0.89	0.30	61,64,65,67	6
8	UNL	В	1018[A]	1/-	0.89	0.27	29,29,29,29	1
8	UNL	В	1018[B]	1/-	0.89	0.27	28,28,28,28	1
8	UNL	В	1019	1/-	0.90	0.37	56,56,56,56	0
8	UNL	В	1015	1/-	0.90	0.35	64,64,64,64	0
8	UNL	D	209	10/-	0.91	0.16	60,70,71,73	0
16	UMQ	С	1002	34/34	0.91	0.23	40,61,78,83	0
8	UNL	A	1024	1/-	0.91	0.19	48,48,48,48	0
8	UNL	В	1005	4/-	0.92	0.12	51,54,57,63	0
8	UNL	С	1012	3/-	0.92	0.30	73,73,74,76	0
8	UNL	С	1013	3/-	0.92	0.10	68,68,68,68	0
8	UNL	A	1007	4/-	0.92	0.13	69,71,73,76	0
8	UNL	D	204	5/-	0.93	0.29	45,48,49,51	5
8	UNL	A	1019	3/-	0.93	0.15	64,64,66,66	0
8	UNL	A	1026	1/-	0.93	0.25	56,56,56,56	0
8	UNL	A	1015	3/-	0.93	0.14	65,65,68,69	0
8	UNL	C	1014	1/-	0.93	0.13	49,49,49,49	0
8	UNL	C	1004	3/-	0.93	0.13	50,50,52,54	0
8	UNL	A	1006	7/-	0.93	0.16	68,70,71,72	0
8	UNL	A	1008	4/-	0.94	0.10	63,66,67,69	0
8	UNL	В	1008	3/-	0.94	0.12	40,40,50,53	0
8	UNL	A	1004	6/-	0.95	0.15	66,67,69,77	4
6	Y3P	A	1002	9/9	0.95	0.13	41,53,64,66	0
8	UNL	A	1009	4/-	0.95	0.09	65,65,66,73	0
7	K	В	1004	1/1	0.97	0.17	84,84,84,84	0
12	SF4	В	1002	8/8	0.97	0.11	30,36,41,41	0
14	HEM	C	201	41/43	0.97	0.18	34,42,51,53	0
5	FAD	A	1001	53/53	0.98	0.13	32,39,46,49	0
7	K	A	1003	$\frac{1/1}{7/7}$	0.99	0.09	53,53,53,53	0
13	F3S	В	1003	7/7	0.99	0.15	31,35,36,36	0
11	FES	В	1001	4/4	0.99	0.10	36,38,38,39	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








# Electron density around 9AU C 202: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)

#### Electron density around UMQ C 1002:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











# 









# 6.5 Other polymers (i)

There are no such residues in this entry.

