

wwPDB X-ray Structure Validation Summary Report (i)

Mar 2, 2022 – 12:06 PM EST

PDB ID : 7N0J

Title : Structure of YebY from E. coli K12 Authors : Hadley, R.C.; Rosenzweig, A.C.

Deposited on : 2021-05-25

Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.27

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)
oteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.27

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	101	68%	22%	• 7%
1	В	101	63%	28%	9%
1	С	101	57%	31%	10%
1	D	101	77%	15%	8%
1	Е	101	68%	23%	•• 6%



Continued from previous page...

Mol	Chain	Length	Quality of chain					
1	F	101	71%	20%	9%			
1	G	101	66%	25%	• 8%			
1	Н	101	72%	19%	9%			
1	I	101	70%	22%	8%			
1	J	101	61%	29%	• 9%			
1	K	101	76%	15%	• 8%			
1	L	101	74%	14%	• 9%			



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9024 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called YebY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	94	Total	С	N	О	S	Se	0	0	0
1	A	94	730	462	125	139	2	2	0	U	
1	С	91	Total	С	N	О	S	Se	0	0	0
1		91	711	449	122	136	2	2	U	U	U
1	D	93	Total	С	N	О	S	Se	0	0	0
1	D	90	723	457	124	138	2	2	U	U	U
1	E	95	Total	С	N	Ο	S	Se	0	0	0
1	ш	30	738	466	126	142	2	2	O	0	U
1	F	92	Total	С	N	Ο	S	Se	0	0	0
1	I.	32	718	454	123	137	2	2	O	O	U
1	Н	92	Total	\mathbf{C}	N	Ο	S	Se	0	0	0
1	11	92	718	454	123	137	2	2	U	O	0
1	I	93	Total	С	N	Ο	S	Se	0	0	0
1	1	30	725	459	124	138	2	2	O	U	U
1	J	92	Total	\mathbf{C}	N	O	S	Se	0	0	0
1	9	32	718	454	123	137	2	2	O	0	U
1	В	92	Total	С	N	O	S	Se	0	0	0
1	D	32	718	454	123	137	2	2	0		U
1	G	93	Total	С	N	О	S	Se	0	0	0
1	G G	30	723	457	124	138	2	2	0	0	U
1	K	93	Total	С	N	O	S	Se	0	0	0
1	17	30	723	457	124	138	2	2	0	0	U
1	L	92	Total	\mathbf{C}	N	O	S	Se	0	0	0
1	L	52	718	454	123	137	2	2			U

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	VAL	-	expression tag	UNP P64506
A	95	ASP	-	expression tag	UNP P64506
A	96	HIS	-	expression tag	UNP P64506
A	97	HIS	-	expression tag	UNP P64506
A	98	HIS	ı	expression tag	UNP P64506



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	99	HIS	-	expression tag	UNP P64506
A	100	HIS	-	expression tag	UNP P64506
A	101	HIS	-	expression tag	UNP P64506
С	94	VAL	-	expression tag	UNP P64506
С	95	ASP	-	expression tag	UNP P64506
С	96	HIS	-	expression tag	UNP P64506
С	97	HIS	-	expression tag	UNP P64506
С	98	HIS	-	expression tag	UNP P64506
С	99	HIS	-	expression tag	UNP P64506
С	100	HIS	-	expression tag	UNP P64506
С	101	HIS	-	expression tag	UNP P64506
D	94	VAL	-	expression tag	UNP P64506
D	95	ASP	_	expression tag	UNP P64506
D	96	HIS	-	expression tag	UNP P64506
D	97	HIS	-	expression tag	UNP P64506
D	98	HIS	-	expression tag	UNP P64506
D	99	HIS	-	expression tag	UNP P64506
D	100	HIS	-	expression tag	UNP P64506
D	101	HIS	-	expression tag	UNP P64506
Е	94	VAL	-	expression tag	UNP P64506
Е	95	ASP	-	expression tag	UNP P64506
Е	96	HIS	-	expression tag	UNP P64506
Е	97	HIS	-	expression tag	UNP P64506
Е	98	HIS	-	expression tag	UNP P64506
Е	99	HIS	-	expression tag	UNP P64506
Е	100	HIS	-	expression tag	UNP P64506
Е	101	HIS	-	expression tag	UNP P64506
F	94	VAL	-	expression tag	UNP P64506
F	95	ASP	-	expression tag	UNP P64506
F	96	HIS	-	expression tag	UNP P64506
F	97	HIS	-	expression tag	UNP P64506
F	98	HIS	-	expression tag	UNP P64506
F	99	HIS	-	expression tag	UNP P64506
F	100	HIS	-	expression tag	UNP P64506
F	101	HIS	-	expression tag	UNP P64506
Н	94	VAL	-	expression tag	UNP P64506
Н	95	ASP	_	expression tag	UNP P64506
Н	96	HIS	-	expression tag	UNP P64506
Н	97	HIS	-	expression tag	UNP P64506
Н	98	HIS	-	expression tag	UNP P64506
Н	99	HIS	-	expression tag	UNP P64506
Н	100	HIS	-	expression tag	UNP P64506



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Н	101	HIS	-	expression tag	UNP P64506
I	94	VAL	-	expression tag	UNP P64506
I	95	ASP	-	expression tag	UNP P64506
I	96	HIS	-	expression tag	UNP P64506
I	97	HIS	-	expression tag	UNP P64506
I	98	HIS	-	expression tag	UNP P64506
I	99	HIS	-	expression tag	UNP P64506
I	100	HIS	-	expression tag	UNP P64506
I	101	HIS	-	expression tag	UNP P64506
J	94	VAL	-	expression tag	UNP P64506
J	95	ASP	-	expression tag	UNP P64506
J	96	HIS	-	expression tag	UNP P64506
J	97	HIS	-	expression tag	UNP P64506
J	98	HIS	-	expression tag	UNP P64506
J	99	HIS	-	expression tag	UNP P64506
J	100	HIS	-	expression tag	UNP P64506
J	101	HIS	-	expression tag	UNP P64506
В	94	VAL	_	expression tag	UNP P64506
В	95	ASP	_	expression tag	UNP P64506
В	96	HIS	-	expression tag	UNP P64506
В	97	HIS	_	expression tag	UNP P64506
В	98	HIS	-	expression tag	UNP P64506
В	99	HIS	-	expression tag	UNP P64506
В	100	HIS	-	expression tag	UNP P64506
В	101	HIS	-	expression tag	UNP P64506
G	94	VAL	-	expression tag	UNP P64506
G	95	ASP	-	expression tag	UNP P64506
G	96	HIS	-	expression tag	UNP P64506
G	97	HIS	-	expression tag	UNP P64506
G	98	HIS	-	expression tag	UNP P64506
G	99	HIS	-	expression tag	UNP P64506
G	100	HIS	-	expression tag	UNP P64506
G	101	HIS	-	expression tag	UNP P64506
K	94	VAL	-	expression tag	UNP P64506
K	95	ASP	-	expression tag	UNP P64506
K	96	HIS	-	expression tag	UNP P64506
K	97	HIS	-	expression tag	UNP P64506
K	98	HIS	-	expression tag	UNP P64506
K	99	HIS	-	expression tag	UNP P64506
K	100	HIS	-	expression tag	UNP P64506
K	101	HIS	-	expression tag	UNP P64506
L	94	VAL	-	expression tag	UNP P64506



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	95	ASP	-	expression tag	UNP P64506
L	96	HIS	-	expression tag	UNP P64506
L	97	HIS	-	expression tag	UNP P64506
L	98	HIS	-	expression tag	UNP P64506
L	99	HIS	-	expression tag	UNP P64506
L	100	HIS	-	expression tag	UNP P64506
L	101	HIS	-	expression tag	UNP P64506


• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	A	52	Total O	0	0	
	Λ	52	52 52	0	U	
2	С	35	Total O	0	0	
	C	30	35 35	0	U	
2	D	33	Total O	0	0	
	D	99	33 33	U	Ü	
2	E	23	Total O	0	0	
	П	20	23 23	Ů,	Ü	
2	F	22	Total O	0	0	
	1	22	22 22	U	U	
2	Н	31	Total O	0	0	
	11	01	31 31	Ü		
2	I	17	17 Total O 0	0		
	1	11	17 17	Ů	Ü	
2	J	41	Total O	0	0	
			41 41		0	
2	В	17	Total O	0	0	
		11	17 17		0	
2	G	28	Total O	0	0	
			28 28		0	
2	K	19	Total O	0	0	
	1.	10	19 19		Ŭ	
2	L	43	Total O	0	0	
		10	43 43		Ŭ	




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	96.14Å 53.20Å 132.36Å	Donogitor
a, b, c, α , β , γ	90.00° 102.25° 90.00°	Depositor
Resolution (Å)	34.67 - 1.88	Depositor
rtesolution (A)	49.20 - 1.88	EDS
% Data completeness	90.3 (34.67-1.88)	Depositor
(in resolution range)	90.5 (49.20-1.88)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.50 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
P. P.	0.235 , 0.282	Depositor
R, R_{free}	0.235 , 0.283	DCC
R_{free} test set	2000 reflections (2.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.540	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 39.3	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9024	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Во	ond angles
IVIOI	Wioi Chain		# Z > 5	RMSZ	# Z > 5
1	A	0.93	1/742~(0.1%)	0.75	0/1006
1	В	0.70	$2/730 \ (0.3\%)$	0.72	0/988
1	С	0.62	$3/722 \ (0.4\%)$	0.68	0/977
1	D	0.89	3/735~(0.4%)	0.79	0/996
1	Ε	0.93	1/750~(0.1%)	0.80	1/1017 (0.1%)
1	F	0.87	1/730 (0.1%)	0.78	1/988 (0.1%)
1	G	1.10	4/735~(0.5%)	0.77	2/996~(0.2%)
1	Н	0.83	0/729	0.78	0/987
1	I	1.10	1/737 (0.1%)	0.98	1/998 (0.1%)
1	J	0.62	1/729~(0.1%)	0.73	1/987 (0.1%)
1	K	0.69	1/735~(0.1%)	0.67	0/996
1	L	0.78	0/730	0.74	0/988
All	All	0.85	18/8804 (0.2%)	0.77	6/11924 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
1	Е	2	PRO	N-CD	-19.13	1.21	1.47
1	F	2	PRO	N-CD	-15.79	1.25	1.47
1	G	76	PRO	N-CD	-15.70	1.25	1.47
1	G	73	PRO	N-CD	-13.64	1.28	1.47
1	В	93	CYS	CB-SG	9.58	1.98	1.82

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	Ε	2	PRO	N-CD-CG	8.27	115.61	103.20
1	J	27	THR	CB-CA-C	-7.79	90.57	111.60
1	G	73	PRO	N-CD-CG	6.35	112.72	103.20
1	I	80	MSE	CG-SE-CE	5.85	111.77	98.90
1	G	28	CYS	CB-CA-C	-5.57	99.26	110.40

There are no chirality outliers.



There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	730	0	738	25	0
1	В	718	0	722	14	0
1	С	711	0	713	27	0
1	D	723	0	729	17	0
1	Е	738	0	742	20	0
1	F	718	0	722	14	0
1	G	723	0	729	18	0
1	Н	718	0	723	27	0
1	I	725	0	731	12	0
1	J	718	0	723	21	0
1	K	723	0	729	10	0
1	L	718	0	722	24	0
2	A	52	0	0	1	0
2	В	17	0	0	0	0
2	С	35	0	0	0	0
2	D	33	0	0	1	0
2	${ m E}$	23	0	0	1	0
2	F	22	0	0	0	0
2	G	28	0	0	0	0
2	Н	31	0	0	0	0
2	I	17	0	0	0	0
2	J	41	0	0	0	0
2	K	19	0	0	0	0
2	L	43	0	0	1	0
All	All	9024	0	8723	205	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 205 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:C:21:ARG:HD2	1:C:22:GLU:H	1.13	1.07



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance } (ext{Å}) \end{aligned}$	Clash overlap (Å)
1:A:44:GLN:HG2	1:H:42:LEU:HD12	1.37	1.05
1:C:64:PRO:O	1:C:67:VAL:HG12	1.65	0.95
1:L:21:ARG:HH11	1:L:21:ARG:HG3	1.29	0.95
1:A:44:GLN:CG	1:H:42:LEU:HD12	1.96	0.94

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	92/101 (91%)	91 (99%)	1 (1%)	0	100	100
1	В	90/101 (89%)	87 (97%)	3 (3%)	0	100	100
1	С	89/101 (88%)	87 (98%)	2 (2%)	0	100	100
1	D	91/101 (90%)	89 (98%)	2 (2%)	0	100	100
1	E	93/101 (92%)	86 (92%)	6 (6%)	1 (1%)	14	5
1	F	90/101 (89%)	85 (94%)	5 (6%)	0	100	100
1	G	91/101 (90%)	90 (99%)	1 (1%)	0	100	100
1	Н	90/101 (89%)	89 (99%)	1 (1%)	0	100	100
1	I	91/101 (90%)	89 (98%)	2 (2%)	0	100	100
1	J	90/101 (89%)	87 (97%)	3 (3%)	0	100	100
1	K	91/101 (90%)	89 (98%)	2 (2%)	0	100	100
1	L	90/101 (89%)	89 (99%)	1 (1%)	0	100	100
All	All	1088/1212 (90%)	1058 (97%)	29 (3%)	1 (0%)	51	41

All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	Ε	59	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	82/87 (94%)	80 (98%)	2 (2%)	49	39
1	В	81/87 (93%)	77 (95%)	4 (5%)	25	13
1	C	80/87 (92%)	78 (98%)	2 (2%)	47	37
1	D	81/87 (93%)	80 (99%)	1 (1%)	71	67
1	E	83/87 (95%)	78 (94%)	5 (6%)	19	8
1	F	81/87 (93%)	81 (100%)	0	100	100
1	G	81/87 (93%)	77 (95%)	4 (5%)	25	13
1	Н	81/87 (93%)	80 (99%)	1 (1%)	71	67
1	I	82/87 (94%)	80 (98%)	2 (2%)	49	39
1	J	81/87 (93%)	79 (98%)	2 (2%)	47	37
1	K	81/87 (93%)	79 (98%)	2 (2%)	47	37
1	L	81/87 (93%)	78 (96%)	3 (4%)	34	22
All	All	975/1044 (93%)	947 (97%)	28 (3%)	42	32

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	45	TYR
1	L	78	GLU
1	В	45	TYR
1	K	67	VAL
1	В	40	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:



Mol	Chain	Res	Type
1	I	52	GLN
1	J	32	ASN
1	L	75	ASN
1	K	69	GLN
1	L	3	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	92/101 (91%)	0.33	1 (1%) 80 82	28, 34, 47, 52	0
1	В	90/101 (89%)	0.24	3 (3%) 46 47	35, 44, 61, 70	0
1	С	89/101 (88%)	0.16	2 (2%) 62 63	33, 40, 62, 69	0
1	D	91/101 (90%)	0.20	1 (1%) 80 82	33, 43, 56, 68	0
1	E	93/101 (92%)	0.51	5 (5%) 25 27	30, 39, 61, 70	0
1	F	90/101 (89%)	0.32	4 (4%) 34 35	34, 44, 65, 72	0
1	G	91/101 (90%)	0.27	3 (3%) 46 47	30, 42, 53, 60	0
1	Н	90/101 (89%)	0.49	2 (2%) 62 63	30, 37, 61, 70	0
1	I	91/101 (90%)	0.44	3 (3%) 46 47	35, 44, 69, 76	0
1	J	90/101 (89%)	0.42	1 (1%) 80 82	31, 38, 48, 62	0
1	K	91/101 (90%)	0.26	3 (3%) 46 47	29, 42, 55, 57	0
1	L	90/101 (89%)	0.08	0 100 100	29, 37, 51, 59	0
All	All	1088/1212 (89%)	0.31	28 (2%) 56 57	28, 40, 59, 76	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	1	ALA	7.3
1	K	76	PRO	5.3
1	F	76	PRO	4.9
1	D	1	ALA	4.8
1	I	2	PRO	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

