

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 1, 2023 – 10:29 PM EDT

PDB ID	:	6N1C
Title	:	Crystal structure of Inorganic pyrophosphatase from Legionella pneumophila
		Philadelphia 1
Authors	:	Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on		
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	175	Total	С	Ν	0	S	0	2	0
		175	1356	870	217	261	8	0	Δ	0
1	В	171	Total	С	Ν	0	S	0	3	0
	D	1/1	1312	844	211	249	8	0	5	0
1	С	167	Total	С	Ν	0	S	0	2	0
		107	1269	817	200	244	8	0		0
1	D	171	Total	С	Ν	Ο	S	0	3	0
	D	1/1	1317	849	211	249	8	0	5	U
1	Е	174	Total	С	Ν	Ο	S	0	4	0
		114	1348	867	215	258	8	0	4	0
1	F	171	Total	С	Ν	0	S	0	1	0
			1309	838	210	252	9			U

• Molecule 1 is a protein called Inorganic pyrophosphatase.

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP Q5ZRW2
А	-6	ALA	-	expression tag	UNP Q5ZRW2
А	-5	HIS	-	expression tag	UNP Q5ZRW2
А	-4	HIS	-	expression tag	UNP Q5ZRW2
А	-3	HIS	-	expression tag	UNP Q5ZRW2
A	-2	HIS	-	expression tag	UNP Q5ZRW2
А	-1	HIS	-	expression tag	UNP Q5ZRW2
А	0	HIS	-	expression tag	UNP Q5ZRW2
В	-7	MET	-	initiating methionine	UNP Q5ZRW2
В	-6	ALA	-	expression tag	UNP Q5ZRW2
В	-5	HIS	-	expression tag	UNP Q5ZRW2
В	-4	HIS	-	expression tag	UNP Q5ZRW2
В	-3	HIS	-	expression tag	UNP Q5ZRW2
В	-2	HIS	-	expression tag	UNP Q5ZRW2
В	-1	HIS	-	expression tag	UNP Q5ZRW2
В	0	HIS	-	expression tag	UNP Q5ZRW2
С	-7	MET	-	initiating methionine	UNP Q5ZRW2

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	-6	ALA	-	expression tag	UNP Q5ZRW2
С	-5	HIS	_	expression tag	UNP Q5ZRW2
С	-4	HIS	-	expression tag	UNP Q5ZRW2
С	-3	HIS	-	expression tag	UNP Q5ZRW2
С	-2	HIS	-	expression tag	UNP Q5ZRW2
С	-1	HIS	-	expression tag	UNP Q5ZRW2
С	0	HIS	-	expression tag	UNP Q5ZRW2
D	-7	MET	-	initiating methionine	UNP Q5ZRW2
D	-6	ALA	-	expression tag	UNP Q5ZRW2
D	-5	HIS	-	expression tag	UNP Q5ZRW2
D	-4	HIS	-	expression tag	UNP Q5ZRW2
D	-3	HIS	-	expression tag	UNP Q5ZRW2
D	-2	HIS	-	expression tag	UNP Q5ZRW2
D	-1	HIS	-	expression tag	UNP Q5ZRW2
D	0	HIS	-	expression tag	UNP Q5ZRW2
Е	-7	MET	-	initiating methionine	UNP Q5ZRW2
E	-6	ALA	-	expression tag	UNP Q5ZRW2
E	-5	HIS	-	expression tag	UNP Q5ZRW2
Е	-4	HIS	-	expression tag	UNP Q5ZRW2
E	-3	HIS	-	expression tag	UNP Q5ZRW2
Е	-2	HIS	-	expression tag	UNP Q5ZRW2
E	-1	HIS	-	expression tag	UNP Q5ZRW2
E	0	HIS	-	expression tag	UNP Q5ZRW2
F	-7	MET	-	initiating methionine	UNP Q5ZRW2
F	-6	ALA	-	expression tag	UNP Q5ZRW2
F	-5	HIS	-	expression tag	UNP Q5ZRW2
F	-4	HIS	-	expression tag	UNP Q5ZRW2
F	-3	HIS	-	expression tag	UNP Q5ZRW2
F	-2	HIS	-	expression tag	UNP Q5ZRW2
F	-1	HIS	-	expression tag	UNP Q5ZRW2
F	0	HIS	-	expression tag	UNP Q5ZRW2


Continued from previous page...

• Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Na 2 2	0	0
2	D	1	Total Na 1 1	0	0
2	F	1	Total Na 1 1	0	0




• Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	Ε	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
3	F	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0


• Molecule 4 is ALANINE (three-letter code: ALA) (formula:  $C_3H_7NO_2$ ).





Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
4	А	1	Total C 6 3			0	0
4	Е	1	Total C 6 3	C N B 1	O 2	0	0

• Molecule 5 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Ate	oms		ZeroOcc	AltConf
5	В	1	Total 8	C 6	O 2	0	0



• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	115	Total O 115 115	0	0
6	В	67	Total         O           67         67	0	0
6	С	50	$\begin{array}{cc} \text{Total} & \text{O} \\ 50 & 50 \end{array}$	0	0
6	D	86	Total O 86 86	0	0
6	Е	99	Total O 99 99	0	0
6	F	63	Total         O           63         63	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	64.08Å 119.94Å 74.90Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $109.59^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	40.97 - 2.00	Depositor
% Data completeness	99.7 (40.97-2.00)	Depositor
(in resolution range)		-
R <sub>merge</sub>	0.05	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.23 (at 2.00 \text{\AA})$	Xtriage
Refinement program	PHENIX $(dev_{3304})$	Depositor
$R, R_{free}$	0.193 , $0.232$	Depositor
Wilson B-factor $(Å^2)$	35.0	Xtriage
Anisotropy	0.323	Xtriage
L-test for twinning <sup>2</sup>	$ < L >=0.52, < L^2>=0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8471	wwPDB-VP
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



			1				0 (	0 /		
Mol	Turne	Chain	Res	Link	B	ond leng	$_{ m gths}$	Bond angles		
NIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
3	MPD	Е	302	-	7,7,7	0.26	0	$9,\!10,\!10$	0.55	0
3	MPD	А	203	-	7,7,7	0.24	0	9,10,10	0.63	0
3	MPD	С	201	-	7,7,7	0.23	0	9,10,10	0.27	0
3	MPD	Е	303	-	7,7,7	0.24	0	9,10,10	0.54	0
3	MPD	В	202	-	7,7,7	0.22	0	$9,\!10,\!10$	0.33	0
4	ALA	Е	301	-	$5,\!5,\!5$	0.95	0	$6,\!6,\!6$	1.36	1 (16%)
3	MPD	F	201	-	7,7,7	0.27	0	9,10,10	0.34	0
4	ALA	А	204	-	$5,\!5,\!5$	1.11	1 (20%)	$6,\!6,\!6$	1.35	1 (16%)
5	MRD	В	201	-	7,7,7	0.10	0	9,10,10	0.28	0
3	MPD	D	202	-	7,7,7	0.26	0	$9,\!10,\!10$	0.42	0

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	Е	302	-	-	3/5/5/5	-
3	MPD	А	203	-	-	2/5/5/5	-
3	MPD	С	201	-	-	0/5/5/5	-
3	MPD	Е	303	-	-	2/5/5/5	-
3	MPD	В	202	-	-	0/5/5/5	-
4	ALA	Е	301	-	-	2/4/4/4	-
3	MPD	F	201	-	-	1/5/5/5	-
4	ALA	А	204	-	-	3/4/4/4	-
5	MRD	В	201	-	-	0/5/5/5	-
3	MPD	D	202	-	-	2/5/5/5	_

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	А	204	ALA	OXT-C	-2.17	1.23	1.30

All (2) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	А	204	ALA	OXT-C-O	-2.58	118.24	124.09
4	Е	301	ALA	OXT-C-O	-2.53	118.34	124.09

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	203	MPD	C2-C3-C4-O4
3	F	201	MPD	C2-C3-C4-C5
4	А	204	ALA	O-C-CA-CB
4	А	204	ALA	OXT-C-CA-CB
4	Е	301	ALA	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 4.7 Other polymers (i)

There are no such residues in this entry.

### 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

## 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

