



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2023 – 08:24 PM EDT

PDB ID : 4N2G  
Title : Crystal structure of Protein Arginine Deiminase 2 (D169A, 10 mM Ca<sup>2+</sup>)  
Authors : Slade, D.J.; Zhang, X.; Fang, P.; Dreyton, C.J.; Zhang, Y.; Gross, M.L.; Guo, M.; Coonrod, S.A.; Thompson, P.R.  
Deposited on : 2013-10-04  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	690	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	708	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
5	GOL	A	710	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein-arginine deiminase type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	659	5164	3321	848	964	31	0	5	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q9Y2J8
A	-19	GLY	-	expression tag	UNP Q9Y2J8
A	-18	HIS	-	expression tag	UNP Q9Y2J8
A	-17	HIS	-	expression tag	UNP Q9Y2J8
A	-16	HIS	-	expression tag	UNP Q9Y2J8
A	-15	HIS	-	expression tag	UNP Q9Y2J8
A	-14	HIS	-	expression tag	UNP Q9Y2J8
A	-13	HIS	-	expression tag	UNP Q9Y2J8
A	-12	HIS	-	expression tag	UNP Q9Y2J8
A	-11	HIS	-	expression tag	UNP Q9Y2J8
A	-10	HIS	-	expression tag	UNP Q9Y2J8
A	-9	HIS	-	expression tag	UNP Q9Y2J8
A	-8	SER	-	expression tag	UNP Q9Y2J8
A	-7	SER	-	expression tag	UNP Q9Y2J8
A	-6	GLY	-	expression tag	UNP Q9Y2J8
A	-5	HIS	-	expression tag	UNP Q9Y2J8
A	-4	ILE	-	expression tag	UNP Q9Y2J8
A	-3	GLU	-	expression tag	UNP Q9Y2J8
A	-2	GLY	-	expression tag	UNP Q9Y2J8
A	-1	ARG	-	expression tag	UNP Q9Y2J8
A	0	HIS	-	expression tag	UNP Q9Y2J8
A	169	ALA	ASP	engineered mutation	UNP Q9Y2J8
A	666	SER	-	expression tag	UNP Q9Y2J8
A	667	ARG	-	expression tag	UNP Q9Y2J8
A	668	ARG	-	expression tag	UNP Q9Y2J8
A	669	SER	-	expression tag	UNP Q9Y2J8

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 6 2	0	0
2	A	1	Total C O 8 6 2	0	0
2	A	1	Total C O 8 6 2	0	0
2	A	1	Total C O 8 6 2	0	0
2	A	1	Total C O 8 6 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total Ca 4 4	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	425	Total 425	O 425	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein-arginine deiminase type-2





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	204.45Å 50.94Å 75.94Å 90.00° 105.98° 90.00°	Depositor
Resolution (Å)	37.78 – 1.85 38.05 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.78-1.85) 96.3 (38.05-1.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.21 (at 1.85Å)	Xtrriage
Refinement program	PHENIX 1.7.2_869	Depositor
R, $R_{free}$	0.178 , 0.216 0.176 , 0.212	Depositor DCC
$R_{free}$ test set	3270 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtrriage
Anisotropy	0.602	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5643	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, MPD, CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/5300	0.60	0/7201

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5164	0	5044	67	1
2	A	40	0	70	11	0
3	A	4	0	0	0	0
4	A	4	0	3	5	0
5	A	6	0	8	6	0
6	A	425	0	0	6	1
All	All	5643	0	5125	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:473:ASP:HB2	5:A:710:GOL:H31	1.51	0.90
1:A:257:PHE:H	4:A:708:ACT:H1	1.42	0.84
1:A:87:GLU:H	1:A:89:SER:HB2	1.41	0.83
1:A:16:GLU:OE1	6:A:803:HOH:O	2.05	0.75
1:A:169:ALA:O	6:A:886:HOH:O	2.09	0.70
1:A:311:VAL:HG22	1:A:649:THR:HG22	1.75	0.68
1:A:418:THR:HG22	1:A:423:THR:HG22	1.77	0.67
1:A:219:ASN:ND2	1:A:224:GLN:OE1	2.28	0.66
1:A:433:SER:HB3	2:A:702:MPD:HM3	1.78	0.65
1:A:217:VAL:HG22	1:A:226:TYR:HD1	1.63	0.62
1:A:550:TRP:HH2	2:A:702:MPD:H31	1.64	0.62
1:A:350:GLN:HE22	2:A:709:MPD:HM3	1.67	0.59
1:A:330:LYS:NZ	1:A:628:GLU:OE1	2.26	0.59
1:A:350:GLN:OE1	2:A:709:MPD:H13	2.03	0.59
1:A:87:GLU:N	1:A:89:SER:HB2	2.14	0.57
1:A:87:GLU:H	1:A:89:SER:CB	2.14	0.57
1:A:257:PHE:H	4:A:708:ACT:CH3	2.14	0.56
1:A:217:VAL:HG22	1:A:226:TYR:CD1	2.41	0.55
1:A:347:ARG:HH22	1:A:590:ASN:HD21	1.53	0.55
1:A:240:LYS:H	2:A:701:MPD:H13	1.72	0.54
1:A:87:GLU:HA	1:A:88:ALA:C	2.28	0.54
1:A:87:GLU:HB2	1:A:191:PRO:HB3	1.90	0.54
1:A:240:LYS:H	2:A:701:MPD:C1	2.21	0.54
1:A:350:GLN:HG3	6:A:1097:HOH:O	2.08	0.53
1:A:373:ARG:HG2	1:A:378:LYS:HG2	1.91	0.51
1:A:347:ARG:HH11	1:A:645:VAL:HB	1.76	0.51
2:A:709:MPD:O2	6:A:1011:HOH:O	2.19	0.50
1:A:218:GLU:OE2	1:A:225:ARG:NE	2.45	0.50
1:A:592:ILE:HG13	2:A:709:MPD:H32	1.94	0.50
1:A:395:ARG:HH22	4:A:708:ACT:CH3	2.25	0.49
1:A:577:GLU:H	1:A:577:GLU:CD	2.15	0.49
1:A:27:THR:HB	1:A:77:LEU:HB2	1.93	0.49
1:A:473:ASP:CB	5:A:710:GOL:H31	2.35	0.49
1:A:353:ILE:HG23	1:A:368:VAL:HG13	1.94	0.48
1:A:11:TYR:HE1	1:A:34:PRO:HA	1.78	0.48
1:A:485:LYS:HE3	1:A:485:LYS:H	1.79	0.47
1:A:473:ASP:HB2	5:A:710:GOL:C3	2.35	0.47
1:A:199:GLU:OE1	2:A:711:MPD:H53	2.14	0.47
1:A:349:ILE:HD12	1:A:353:ILE:HD12	1.97	0.46
1:A:530:LYS:NZ	6:A:1186:HOH:O	2.48	0.46
1:A:189:LYS:HA	1:A:244:GLY:O	2.16	0.46
1:A:312[B]:CYS:SG	1:A:349:ILE:HG22	2.55	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:HIS:O	1:A:363:LYS:HB3	2.16	0.46
1:A:591:MET:H	5:A:710:GOL:C1	2.29	0.45
1:A:395:ARG:HH22	4:A:708:ACT:H3	1.80	0.45
1:A:370:ASP:OD1	1:A:378:LYS:NZ	2.45	0.45
1:A:319:LEU:HD12	1:A:319:LEU:HA	1.76	0.44
1:A:6:THR:OG1	1:A:26:TRP:NE1	2.49	0.44
1:A:485:LYS:H	1:A:485:LYS:CD	2.31	0.44
1:A:305:PRO:HB2	1:A:654:LYS:HE2	2.00	0.43
1:A:514:MET:HG3	1:A:526:ILE:HG22	2.00	0.43
1:A:590:ASN:OD1	5:A:710:GOL:H32	2.19	0.43
1:A:485:LYS:H	1:A:485:LYS:CE	2.32	0.43
1:A:550:TRP:CH2	2:A:702:MPD:H31	2.50	0.43
1:A:96:ASN:OD1	1:A:108:GLN:NE2	2.52	0.43
1:A:164:LYS:HD2	1:A:168:ARG:HH11	1.84	0.43
1:A:431:ILE:HG13	1:A:461:GLU:HG2	2.01	0.43
1:A:257:PHE:N	4:A:708:ACT:H1	2.22	0.42
1:A:530:LYS:NZ	6:A:1041:HOH:O	2.38	0.42
1:A:38:GLN:HG3	1:A:100:GLU:HG2	2.02	0.42
1:A:204:ILE:HD12	1:A:209:SER:HA	2.03	0.41
1:A:220:PRO:HD2	1:A:223:GLY:O	2.21	0.41
1:A:461:GLU:HB3	2:A:702:MPD:CM	2.50	0.41
1:A:175:LYS:HZ1	1:A:179:LYS:HG3	1.86	0.41
1:A:176:GLU:OE2	1:A:179:LYS:NZ	2.31	0.41
1:A:604:PHE:CD1	1:A:635:ILE:HD11	2.56	0.40
1:A:591:MET:H	5:A:710:GOL:H11	1.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:GLU:OE2	6:A:1173:HOH:O[3_455]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	660/690 (96%)	643 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	563/611 (92%)	548 (97%)	15 (3%)	44	29

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	GLU
1	A	28	ASP
1	A	100	GLU
1	A	145	GLU
1	A	175	LYS
1	A	176	GLU
1	A	206	MET
1	A	234	LYS
1	A	314	MET
1	A	319	LEU
1	A	347	ARG
1	A	382	VAL
1	A	485	LYS
1	A	530	LYS
1	A	642	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	96	ASN
1	A	108	GLN
1	A	219	ASN
1	A	224	GLN
1	A	590	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MPD	A	707	-	7,7,7	0.24	0	9,10,10	0.22	0
2	MPD	A	702	-	7,7,7	0.27	0	9,10,10	0.52	0
2	MPD	A	711	-	7,7,7	0.26	0	9,10,10	0.31	0
2	MPD	A	709	-	7,7,7	0.32	0	9,10,10	0.56	0
2	MPD	A	701	-	7,7,7	0.26	0	9,10,10	0.34	0
5	GOL	A	710	-	5,5,5	0.23	0	5,5,5	1.01	0
4	ACT	A	708	-	3,3,3	0.66	0	3,3,3	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	A	707	-	-	4/5/5/5	-
2	MPD	A	702	-	-	2/5/5/5	-
2	MPD	A	711	-	-	1/5/5/5	-
2	MPD	A	709	-	-	2/5/5/5	-
2	MPD	A	701	-	-	4/5/5/5	-
5	GOL	A	710	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	MPD	C1-C2-C3-C4
2	A	701	MPD	O2-C2-C3-C4
2	A	707	MPD	C2-C3-C4-O4
2	A	709	MPD	C2-C3-C4-O4
2	A	711	MPD	C2-C3-C4-O4
5	A	710	GOL	O1-C1-C2-O2
2	A	707	MPD	O2-C2-C3-C4
5	A	710	GOL	O1-C1-C2-C3
2	A	701	MPD	CM-C2-C3-C4
2	A	707	MPD	C1-C2-C3-C4
2	A	707	MPD	CM-C2-C3-C4
2	A	709	MPD	C1-C2-C3-C4
2	A	702	MPD	O2-C2-C3-C4
2	A	701	MPD	C2-C3-C4-C5
2	A	702	MPD	C2-C3-C4-C5

There are no ring outliers.

6 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	MPD	4	0
2	A	711	MPD	1	0
2	A	709	MPD	4	0
2	A	701	MPD	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	710	GOL	6	0
4	A	708	ACT	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	659/690 (95%)	-0.09	23 (3%) 44 41	16, 33, 67, 99	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	377	LEU	5.7
1	A	399	PHE	5.2
1	A	380	PHE	4.5
1	A	161	TRP	3.7
1	A	374	ASP	3.6
1	A	381	PRO	3.6
1	A	89	SER	3.1
1	A	88	ALA	3.0
1	A	173	TYR	2.9
1	A	375	GLY	2.8
1	A	664	VAL	2.8
1	A	373	ARG	2.7
1	A	646	HIS	2.5
1	A	376	ASN	2.4
1	A	99	ASP	2.3
1	A	642	LEU	2.2
1	A	346	ASP	2.2
1	A	90	SER	2.1
1	A	645	VAL	2.1
1	A	174	SER	2.1
1	A	169	ALA	2.1
1	A	666	SER	2.0
1	A	101	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ACT	A	708	4/4	0.67	0.19	21,40,42,48	0
2	MPD	A	711	8/8	0.83	0.19	55,60,65,70	0
2	MPD	A	709	8/8	0.84	0.24	47,56,59,66	0
2	MPD	A	707	8/8	0.85	0.15	42,67,75,75	0
5	GOL	A	710	6/6	0.85	0.20	25,43,45,45	0
2	MPD	A	702	8/8	0.88	0.23	35,47,55,57	0
2	MPD	A	701	8/8	0.90	0.22	35,58,66,73	0
3	CA	A	704	1/1	0.95	0.06	30,30,30,30	0
3	CA	A	705	1/1	0.96	0.06	28,28,28,28	0
3	CA	A	706	1/1	0.99	0.07	36,36,36,36	0
3	CA	A	703	1/1	1.00	0.08	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.