



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 04:38 am BST

PDB ID : 5N2E
Title : Structure of the E9 DNA polymerase from vaccinia virus
Authors : Tarbouriech, N.; Burmeister, W.P.; Iseni, F.
Deposited on : 2017-02-07
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1010	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	999	8188	5239	1364	1533	52	0	2	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P20509
A	-2	ALA	-	expression tag	UNP P20509
A	-1	MET	-	expression tag	UNP P20509
A	0	ASP	-	expression tag	UNP P20509
A	1	PRO	-	expression tag	UNP P20509

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	15	8	2	4	1	0	0

- Molecule 5 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
5	A	1	8	4	2	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	219	Total 219	O 219	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	133.41Å 133.41Å 230.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	115.81 – 2.74 46.18 – 2.74	Depositor EDS
% Data completeness (in resolution range)	99.8 (115.81-2.74) 99.9 (46.18-2.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.73Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.186 , 0.236 0.191 , 0.236	Depositor DCC
R_{free} test set	3075 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	64.5	Xtrriage
Anisotropy	0.070	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8478	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EPE, DTT, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/8372	0.93	16/11309 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	A	23	ARG	NE-CZ-NH1	6.37	123.48	120.30
1	A	23	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	A	577	ARG	NE-CZ-NH2	-6.19	117.20	120.30
1	A	365	VAL	CB-CA-C	5.95	122.71	111.40
1	A	246	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	A	625	ARG	NE-CZ-NH2	-5.76	117.42	120.30
1	A	617	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	A	727	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	81	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	727	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	577	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	A	205	GLU	CB-CA-C	-5.33	99.73	110.40
1	A	146	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	234	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	A	244	LEU	CA-CB-CG	5.10	127.03	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	281	GLY	Peptide
1	A	424	LEU	Peptide
1	A	859	MET	Peptide
1	A	860	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8188	0	8140	68	0
2	A	36	0	39	4	0
3	A	12	0	16	0	0
4	A	15	0	18	0	0
5	A	8	0	10	0	0
6	A	219	0	0	5	1
All	All	8478	0	8223	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:833:ARG:HG3	1:A:893:SER:OG	1.49	1.08
1:A:681:TYR:CZ	1:A:685:LYS:HD2	2.25	0.71
1:A:946:ASN:C	1:A:948:LYS:H	1.92	0.71
1:A:266:ASN:OD1	1:A:270:ARG:NH1	2.29	0.66
1:A:753:ASP:HB2	6:A:1317:HOH:O	1.98	0.63
1:A:528[B]:TYR:CD1	1:A:685:LYS:HG2	2.33	0.62
1:A:946:ASN:C	1:A:948:LYS:N	2.53	0.62
1:A:833:ARG:CG	1:A:893:SER:OG	2.39	0.61
1:A:656:MET:HE3	6:A:1275:HOH:O	2.01	0.61
1:A:833:ARG:HB3	1:A:893:SER:H	1.66	0.59
1:A:495:GLU:OE1	2:A:1103:MES:H52	2.03	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528[B]:TYR:CE2	1:A:688:THR:HG21	2.39	0.58
1:A:23:ARG:NH2	1:A:257:ASP:OD2	2.37	0.58
1:A:266:ASN:O	1:A:270:ARG:HD2	2.04	0.57
1:A:304:GLN:O	1:A:314:ALA:HB3	2.04	0.57
1:A:681:TYR:CE1	1:A:685:LYS:HD2	2.39	0.56
1:A:497:ARG:HG2	2:A:1102:MES:O1S	2.06	0.55
1:A:764:VAL:CG2	1:A:817:SER:HA	2.36	0.55
1:A:528[A]:TYR:CD2	1:A:684:ALA:HB1	2.42	0.55
1:A:403:CYS:SG	1:A:421:CYS:HB3	2.48	0.53
1:A:833:ARG:HG3	1:A:893:SER:CB	2.36	0.52
1:A:205:GLU:OE2	1:A:223:ARG:NE	2.43	0.52
1:A:546:LEU:HD22	1:A:771:ALA:HB2	1.90	0.52
1:A:26:ASN:N	1:A:26:ASN:OD1	2.43	0.52
1:A:800:MET:HB2	1:A:806:TYR:HB2	1.93	0.51
1:A:862:ASN:N	1:A:862:ASN:OD1	2.44	0.51
1:A:528[A]:TYR:CE2	1:A:672:GLY:HA3	2.44	0.51
1:A:712:LEU:HD23	1:A:714:PHE:CZ	2.46	0.50
1:A:495:GLU:OE1	2:A:1103:MES:H61	2.12	0.50
1:A:751:ASP:O	1:A:753:ASP:N	2.44	0.50
1:A:751:ASP:O	1:A:752:THR:C	2.50	0.49
1:A:528[A]:TYR:CE2	1:A:684:ALA:HB1	2.48	0.49
1:A:587:LEU:HD21	1:A:614:ILE:HD11	1.95	0.48
1:A:833:ARG:NH1	1:A:835:VAL:O	2.47	0.48
1:A:528[B]:TYR:HE2	1:A:688:THR:HG21	1.77	0.48
1:A:890:PHE:HB3	1:A:971:TYR:CZ	2.49	0.47
1:A:528[B]:TYR:CD1	1:A:685:LYS:CG	2.98	0.47
1:A:297:LEU:C	1:A:297:LEU:HD23	2.35	0.47
1:A:491:SER:HB2	2:A:1103:MES:H21	1.97	0.46
1:A:633:GLU:OE2	1:A:636:ARG:NH1	2.49	0.45
1:A:833:ARG:HB3	1:A:893:SER:N	2.30	0.45
1:A:442:GLN:HG2	1:A:924:LEU:CD2	2.47	0.45
1:A:835:VAL:HG13	1:A:839:HIS:HB3	1.98	0.45
1:A:853:MET:HE3	1:A:864:VAL:HG22	1.99	0.44
1:A:831:THR:O	1:A:832:ARG:C	2.56	0.44
1:A:133:THR:HG22	6:A:1396:HOH:O	2.16	0.44
1:A:946:ASN:O	1:A:948:LYS:N	2.51	0.44
1:A:746:ARG:NH1	1:A:758:GLU:OE1	2.47	0.44
1:A:170:HIS:HA	1:A:444:TYR:CZ	2.52	0.44
1:A:872:LEU:HD21	1:A:976:THR:HG22	1.99	0.44
1:A:266:ASN:HB2	1:A:318:PHE:CD2	2.53	0.43
1:A:170:HIS:HA	1:A:444:TYR:CE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:ILE:HD11	1:A:420:LEU:HB2	2.01	0.43
1:A:523:LYS:HE2	1:A:674:ARG:O	2.19	0.42
1:A:800:MET:HB2	1:A:806:TYR:CB	2.49	0.42
1:A:319:HIS:CE1	1:A:495:GLU:HG3	2.55	0.42
1:A:209:THR:HG21	6:A:1259:HOH:O	2.19	0.41
1:A:681:TYR:CZ	1:A:685:LYS:CD	2.98	0.41
1:A:830:GLU:HG3	1:A:831:THR:N	2.35	0.41
1:A:76:LEU:HD21	1:A:587:LEU:HA	2.03	0.41
1:A:277:GLU:O	1:A:280:THR:O	2.38	0.41
1:A:773:GLU:OE2	1:A:776:ARG:NH1	2.54	0.41
1:A:945:VAL:HB	1:A:946:ASN:H	1.66	0.41
1:A:918:ASN:HB3	6:A:1285:HOH:O	2.21	0.41
1:A:116:SER:O	1:A:146:ARG:NH1	2.53	0.40
1:A:23:ARG:HH22	1:A:257:ASP:CG	2.24	0.40
1:A:370:PHE:CD1	1:A:370:PHE:N	2.90	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1265:HOH:O	6:A:1265:HOH:O[6_555]	0.71	1.49

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	995/1010 (98%)	921 (93%)	61 (6%)	13 (1%)	12 21

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	422	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	527	PRO
1	A	528[A]	TYR
1	A	528[B]	TYR
1	A	752	THR
1	A	821	PRO
1	A	51	SER
1	A	365	VAL
1	A	831	THR
1	A	945	VAL
1	A	832	ARG
1	A	425	PRO
1	A	947	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	924/928 (100%)	852 (92%)	72 (8%)	12 22

All (72) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	GLU
1	A	25	ARG
1	A	26	ASN
1	A	43	THR
1	A	79	LYS
1	A	96	LYS
1	A	98	SER
1	A	103	THR
1	A	106	GLU
1	A	129	GLU
1	A	174	LYS
1	A	196	LYS
1	A	277	GLU
1	A	291	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	292	LYS
1	A	304	GLN
1	A	306	SER
1	A	345	LYS
1	A	361	LEU
1	A	376	THR
1	A	381	LYS
1	A	403	CYS
1	A	418	VAL
1	A	420	LEU
1	A	424	LEU
1	A	428	THR
1	A	436	ASP
1	A	467	GLN
1	A	484	SER
1	A	499	SER
1	A	519	ARG
1	A	520	SER
1	A	521	GLU
1	A	526[A]	PHE
1	A	526[B]	PHE
1	A	529	GLU
1	A	532	LYS
1	A	540	MET
1	A	602	GLU
1	A	643	GLN
1	A	648	THR
1	A	674	ARG
1	A	692	ARG
1	A	735	THR
1	A	738	PRO
1	A	746	ARG
1	A	748	VAL
1	A	752	THR
1	A	754	SER
1	A	809	MET
1	A	814	SER
1	A	819	SER
1	A	831	THR
1	A	833	ARG
1	A	835	VAL
1	A	849	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	862	ASN
1	A	868	ILE
1	A	873	GLU
1	A	886	PRO
1	A	891	MET
1	A	905	ASN
1	A	924	LEU
1	A	927	ARG
1	A	941	THR
1	A	949	THR
1	A	962	SER
1	A	967	PHE
1	A	974	ARG
1	A	988	LEU
1	A	1002	THR
1	A	1005	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	816	ASN
1	A	860	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	DTT	A	1107	-	7,7,7	1.13	0	4,8,8	1.60	1 (25%)
4	EPE	A	1106	-	15,15,15	1.90	1 (6%)	18,20,20	1.99	4 (22%)
2	MES	A	1101	-	12,12,12	1.80	1 (8%)	14,16,16	2.37	5 (35%)
3	GOL	A	1104	-	5,5,5	0.68	0	5,5,5	0.58	0
2	MES	A	1102	-	12,12,12	2.02	2 (16%)	14,16,16	2.33	5 (35%)
2	MES	A	1103	-	12,12,12	2.03	1 (8%)	14,16,16	3.22	4 (28%)
3	GOL	A	1105	-	5,5,5	0.45	0	5,5,5	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DTT	A	1107	-	-	2/8/8/8	-
4	EPE	A	1106	-	-	6/9/19/19	0/1/1/1
2	MES	A	1101	-	-	0/6/14/14	0/1/1/1
3	GOL	A	1104	-	-	4/4/4/4	-
2	MES	A	1102	-	-	3/6/14/14	0/1/1/1
2	MES	A	1103	-	-	1/6/14/14	0/1/1/1
3	GOL	A	1105	-	-	2/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1106	EPE	C10-S	-6.62	1.68	1.77
2	A	1103	MES	C8-S	-6.00	1.69	1.77
2	A	1102	MES	C8-S	-5.97	1.69	1.77
2	A	1101	MES	C8-S	-5.56	1.69	1.77
2	A	1102	MES	O1S-S	2.14	1.51	1.45

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1103	MES	O2S-S-C8	9.07	117.83	106.92
2	A	1102	MES	O2S-S-C8	6.74	115.03	106.92
2	A	1103	MES	C6-C5-N4	5.82	118.93	110.10
2	A	1101	MES	O3S-S-C8	4.99	113.84	105.77
4	A	1106	EPE	O2S-S-C10	4.56	112.40	106.92
4	A	1106	EPE	O3S-S-C10	4.46	112.98	105.77
2	A	1101	MES	C6-C5-N4	-4.09	103.91	110.10
2	A	1103	MES	C2-C3-N4	3.73	115.76	110.10
2	A	1101	MES	O2S-S-C8	-3.66	102.50	106.92
4	A	1106	EPE	O3S-S-O2S	-3.28	103.26	111.27
2	A	1101	MES	O1S-S-C8	3.12	110.68	106.92
5	A	1107	DTT	O3-C3-C2	3.11	116.11	109.72
2	A	1102	MES	C6-O1-C2	2.96	119.77	109.89
2	A	1103	MES	O3S-S-O1S	-2.87	104.26	111.27
2	A	1102	MES	C6-C5-N4	-2.41	106.45	110.10
2	A	1101	MES	C7-N4-C3	-2.23	105.53	111.23
2	A	1102	MES	O2S-S-O1S	-2.20	106.35	113.95
4	A	1106	EPE	C6-C5-N4	2.11	114.96	110.64
2	A	1102	MES	O3S-S-O2S	-2.09	106.16	111.27

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1102	MES	C7-C8-S-O1S
2	A	1102	MES	C7-C8-S-O2S
2	A	1102	MES	C7-C8-S-O3S
2	A	1103	MES	N4-C7-C8-S
3	A	1104	GOL	O1-C1-C2-C3
4	A	1106	EPE	N4-C7-C8-O8
4	A	1106	EPE	C9-C10-S-O2S
4	A	1106	EPE	C9-C10-S-O3S
5	A	1107	DTT	C2-C3-C4-S4
5	A	1107	DTT	O3-C3-C4-S4
3	A	1104	GOL	C1-C2-C3-O3
4	A	1106	EPE	C10-C9-N1-C2
4	A	1106	EPE	C10-C9-N1-C6
3	A	1104	GOL	O1-C1-C2-O2
3	A	1104	GOL	O2-C2-C3-O3
3	A	1105	GOL	O2-C2-C3-O3
4	A	1106	EPE	C9-C10-S-O1S
3	A	1105	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1102	MES	1	0
2	A	1103	MES	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	999/1010 (98%)	-0.07	37 (3%) 41 46	37, 66, 132, 167	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	989	CYS	5.9
1	A	528[A]	TYR	4.9
1	A	531	GLY	4.7
1	A	909	TYR	4.7
1	A	1004	TYR	4.6
1	A	900	TYR	4.4
1	A	365	VAL	4.0
1	A	527	PRO	3.5
1	A	990	ILE	3.2
1	A	437	ASP	3.2
1	A	423	THR	3.1
1	A	366	ARG	3.1
1	A	364	GLY	2.9
1	A	359	LYS	2.8
1	A	1000	LYS	2.7
1	A	306	SER	2.7
1	A	1002	THR	2.7
1	A	988	LEU	2.5
1	A	368	MET	2.5
1	A	370	PHE	2.4
1	A	943	LYS	2.4
1	A	371	ILE	2.4
1	A	908	MET	2.3
1	A	991	SER	2.3
1	A	864	VAL	2.3
1	A	363	ARG	2.3
1	A	522	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	401	ILE	2.2
1	A	530	GLY	2.2
1	A	987	VAL	2.2
1	A	361	LEU	2.2
1	A	999	SER	2.1
1	A	832	ARG	2.1
1	A	992	PHE	2.1
1	A	906	PRO	2.1
1	A	829	SER	2.1
1	A	823	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1104	6/6	0.75	0.19	86,106,110,114	0
5	DTT	A	1107	8/8	0.84	0.24	99,114,124,132	0
4	EPE	A	1106	15/15	0.85	0.27	101,117,134,136	0
2	MES	A	1103	12/12	0.91	0.34	93,132,145,149	0
3	GOL	A	1105	6/6	0.94	0.29	74,76,83,92	0
2	MES	A	1102	12/12	0.96	0.14	73,87,95,97	0
2	MES	A	1101	12/12	0.97	0.18	48,64,72,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.