

Nov 28, 2022 – 08:04 AM EST

PDB ID	:	7N32
EMDB ID	:	EMD-22677
Title	:	protofilaments of microtubule doublets bound to outer-arm dynein
Authors	:	Rao, Q.; Zhang, K.
Deposited on	:	2021-05-31
Resolution	:	4.50 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1. dev 43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			47%	
1	a	449	96%	• •
			61%	
1	с	449	95%	• •
			45%	
1	е	449	97%	••
			35%	
1	g	449	97%	••
			58%	
1	i	449	96%	• •
			44%	
1	k	449	97%	•••
			41%	
1	m	449	97%	•••
			60%	
1	0	449	96%	• •



Mol	Chain	Length	Quality of chain	
			48%	
1	q	449	96%	••
			43%	
1	S	449	95%	••
			50%	
1	u	449	97%	••
- 1		1.10	67%	
1	W	449	96%	••
0	1.	449	39%	
	D	443	95%	••
9	d	442	2000/	_
2	u	440	98%	•
2	f	1/13	060/	
2	1	110	32%	••
2	h	443	06%	
-	11	110	59%	••
2	i	443	96%	•••
	J		40%	
2	1	443	97%	•
			36%	
2	n	443	99%	••
			52%	
2	р	443	95%	• •
			42%	
2	r	443	97%	·
-			36%	
2	t	443	96%	• •
		4.4.9	45%	
2	V	443	97%	•
0		449	54%	
2	Х	443	97%	•



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 82021 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			AltConf	Trace
1		440	Total	С	Ν	0	S	0	0
	W	440	3405	2152	578	653	22	0	0
1		440	Total	С	Ν	0	S	0	0
	u	440	3409	2155	579	653	22	0	0
1	G	440	Total	С	Ν	0	S	0	0
	5	440	3409	2155	579	653	22	0	0
1	a	440	Total	С	Ν	0	S	0	0
	q	440	3403	2151	578	652	22	0	0
1	0	440	Total	С	Ν	0	S	0	0
1	0	440	3397	2146	577	652	22	0	0
1	m	440	Total	С	Ν	0	S	0	0
	111	440	3409	2155	579	653	22	0	0
1	k	440	Total	С	Ν	0	S	0	0
1	ň	440	3405	2152	578	653	22	0	0
1	ſſ	440	Total	С	Ν	0	S	0	0
	8	440	3413	2158	580	653	22	0	0
1	0	440	Total	С	Ν	0	\mathbf{S}	0	0
	С	440	3402	2149	579	652	22	0	0
1	C	440	Total	С	Ν	0	\mathbf{S}	0	0
L	C	440	3402	2149	579	652	22	0	0
1	9	440	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
	a	440	3406	2152	580	652	22	0	0
1	i	440	Total	С	Ν	0	S	0	0
	1	440	3405	2152	578	653	22	U	U

• Molecule 1 is a protein called Tubulin alpha chain.

• Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	x	430	Total 3366	C 2115	N 577	O 646	S 28	0	0
2	V	430	Total 3366	C 2115	N 577	0 646	S 28	0	0
2	t	430	Total 3366	C 2115	N 577	0 646	S 28	0	0



Mol	Chain	Residues	_	At	oms			AltConf	Trace
9	r	430	Total	С	Ν	0	\mathbf{S}	0	0
	1	430	3359	2109	577	645	28	0	
9	n	428	Total	С	Ν	0	\mathbf{S}	0	0
	р	420	3345	2101	574	642	28	0	
9	n	440	Total	С	Ν	0	\mathbf{S}	0	0
	11	440	3412	2141	587	656	28	0	
2	1	430	Total	С	Ν	0	\mathbf{S}	0	0
2	Ţ	400	3366	2115	577	646	28	0	0
2	h	430	Total	С	Ν	Ο	\mathbf{S}	0	0
2	11	400	3366	2115	577	646	28	0	0
2	f	430	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
2	L	450	3362	2113	576	645	28	0	0
2	h	430	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
	U U	400	3362	2113	576	645	28	0	0
2	d	442	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0
	u	442	3416	2145	588	655	28	0	0
2	i	430	Total	\mathbf{C}	Ν	Ο	S	0	0
2	J	-100	3366	2115	577	646	28	0	0

• Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues		Ate	oms			AltConf
9		1	Total	С	Ν	Ο	Р	0
0	3 W		32	10	5	14	3	0
9		1	Total	С	Ν	0	Р	0
3	u	1	32	10	5	14	3	0



Mol	Chain	Residues		Ate	oms			AltConf
2	G	1	Total	С	Ν	0	Р	0
0	5	L	32	10	5	14	3	0
2		1	Total	С	Ν	Ο	Р	0
0	q	L	32	10	5	14	3	0
2	0	1	Total	С	Ν	0	Р	0
0	0	L	32	10	5	14	3	0
2	m	1	Total	С	Ν	0	Р	0
0	111	L	32	10	5	14	3	0
2	l,	1	Total	С	Ν	0	Р	0
0	K	I	32	10	5	14	3	0
3	ď	1	Total	С	Ν	Ο	Р	0
0	8	T	32	10	5	14	3	0
3	0	1	Total	С	Ν	Ο	Р	0
0	C	T	32	10	5	14	3	0
3	C	1	Total	С	Ν	Ο	Р	0
0	C	T	32	10	5	14	3	0
3	9	1	Total	С	Ν	Ο	Р	0
5	a	I	32	10	5	14	3	U
3	i	1	Total	С	Ν	Ο	Р	0
0			32	10	5	14	3	U

Continued from previous page...

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
4	х	1	Total Mg 1 1	0
4	u	1	Total Mg 1 1	0
4	t	1	Total Mg 1 1	0
4	q	1	Total Mg 1 1	0
4	0	1	Total Mg 1 1	0
4	m	1	Total Mg 1 1	0
4	k	1	Total Mg 1 1	0
4	g	1	Total Mg 1 1	0
4	е	1	Total Mg 1 1	0



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
4	с	1	Total Mg 1 1	0
4	a	1	Total Mg 1 1	0
4	i	1	Total Mg 1 1	0

• Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues		Atoms				AltConf
5		1	Total	С	Ν	0	Р	0
0	X	L	28	10	5	11	2	0
5	37	1	Total	С	Ν	Ο	Р	0
0	v	T	28	10	5	11	2	0
5	+	1	Total	С	Ν	Ο	Р	0
0	U	T	28	10	5	11	2	0
5	r	1	Total	С	Ν	Ο	Р	0
0	1	L	28	10	5	11	2	0
5	n	1	Total	С	Ν	Ο	Р	0
0	р	L	28	10	5	11	2	0
5	n	1	Total	С	Ν	Ο	Р	0
0	11	L	28	10	5	11	2	0
5	1	1	Total	С	Ν	Ο	Р	0
5			28	10	5	11	2	0
5	h	1	Total	С	Ν	Ο	Р	0
5	11		28	10	5	11	2	0



Continued from previous page...

Mol	Chain	Residues		Atoms				AltConf
F	f	1	Total	С	Ν	0	Р	0
0	1	1	28	10	5	11	2	0
5	h	1	Total	С	Ν	Ο	Р	0
5	D	1	28	10	5	11	2	0
5	;	1	Total	С	Ν	0	Р	0
- 0	J	1	28	10	5	11	2	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tubulin alpha chain



















 \bullet Molecule 1: Tubulin alpha chain

35%

Chain g:

97%



••



























G12 C12 D12 C12

G93 Q94 G96 G96 A97 G98 G98 G98 N99











4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	191776	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	53.3	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	11.012	Depositor
Minimum map value	0.000	Depositor
Average map value	0.020	Depositor
Map value standard deviation	0.134	Depositor
Recommended contour level	0.8	Depositor
Map size (Å)	530.5338, 477.21387, 449.22086	wwPDB
Map dimensions	398, 358, 337	wwPDB
Map angles $(^{\circ})$	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.3329996, 1.3329996, 1.3329996	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	a	0.67	0/3477	0.83	0/4712	
1	с	0.69	0/3473	0.82	0/4708	
1	е	0.67	0/3473	0.81	0/4708	
1	g	0.67	0/3485	0.79	0/4723	
1	i	0.68	0/3477	0.79	0/4715	
1	k	0.67	0/3477	0.79	0/4715	
1	m	0.66	0/3481	0.81	0/4719	
1	0	0.68	0/3467	0.79	0/4700	
1	q	0.67	0/3475	0.80	0/4712	
1	s	0.66	0/3481	0.80	0/4719	
1	u	0.67	0/3481	0.80	0/4719	
1	W	0.68	0/3477	0.81	0/4715	
2	b	0.66	0/3435	0.80	0/4650	
2	d	0.69	0/3488	0.82	0/4721	
2	f	0.67	0/3435	0.80	0/4650	
2	h	0.66	0/3439	0.78	0/4655	
2	j	0.68	0/3439	0.78	0/4655	
2	1	0.67	0/3439	0.78	0/4655	
2	n	0.66	0/3485	0.79	0/4719	
2	р	0.68	0/3416	0.78	0/4622	
2	r	0.66	0/3431	0.77	0/4644	
2	t	0.66	0/3439	0.79	0/4655	
2	V	0.66	0/3439	0.78	0/4655	
2	Х	0.67	0/3439	0.77	0/4655	
All	All	0.67	0/83048	0.80	0/112501	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.



Mol	Chain	#Chirality outliers	#Planarity outliers
1	а	0	1
1	с	0	1
1	i	0	1
1	q	0	1
1	s	0	1
1	W	0	1
All	All	0	6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	а	399	TYR	Peptide
1	с	402	ARG	Peptide
1	i	349	THR	Peptide
1	q	402	ARG	Peptide
1	s	402	ARG	Peptide
1	W	402	ARG	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	a	3406	0	3345	0	0
1	с	3402	0	3334	0	0
1	е	3402	0	3334	0	0
1	g	3413	0	3352	0	0
1	i	3405	0	3330	0	0
1	k	3405	0	3330	0	0
1	m	3409	0	3341	0	0
1	0	3397	0	3329	0	0
1	q	3403	0	3325	0	0
1	S	3409	0	3341	0	0
1	u	3409	0	3341	0	0
1	W	3405	0	3330	0	0



	nuea fron	<i>i previous</i>		TT(11 1)		
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	b	3362	0	3251	0	0
2	d	3416	0	3273	0	0
2	f	3362	0	3251	0	0
2	h	3366	0	3257	0	0
2	j	3366	0	3257	0	0
2	1	3366	0	3257	0	0
2	n	3412	0	3269	0	0
2	р	3345	0	3239	0	0
2	r	3359	0	3250	0	0
2	\mathbf{t}	3366	0	3257	0	0
2	V	3366	0	3257	0	0
2	X	3366	0	3257	0	0
3	a	32	0	12	0	0
3	с	32	0	12	0	0
3	е	32	0	12	0	0
3	g	32	0	12	0	0
3	i	32	0	12	0	0
3	k	32	0	12	0	0
3	m	32	0	12	0	0
3	0	32	0	12	0	0
3	q	32	0	12	0	0
3	S	32	0	12	0	0
3	u	32	0	12	0	0
3	W	32	0	12	0	0
4	a	1	0	0	0	0
4	с	1	0	0	0	0
4	е	1	0	0	0	0
4	g	1	0	0	0	0
4	i	1	0	0	0	0
4	k	1	0	0	0	0
4	m	1	0	0	0	0
4	0	1	0	0	0	0
4	q	1	0	0	0	0
4	t	1	0	0	0	0
4	u	1	0	0	0	0
4	X	1	0	0	0	0
5	b	28	0	12	0	0
5	f	28	0	12	0	0
5	h	28	0	12	0	0
5	j	28	0	12	0	0
5	1	28	0	12	0	0
5	n	28	0	12	0	0

 α ntia d fa .



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	р	28	0	12	0	0
5	r	28	0	12	0	0
5	t	28	0	12	0	0
5	V	28	0	12	0	0
5	Х	28	0	12	0	0
All	All	82021	0	79383	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	a	438/449~(98%)	390~(89%)	45 (10%)	3~(1%)	22	62
1	с	438/449~(98%)	405 (92%)	30 (7%)	3(1%)	22	62
1	е	438/449~(98%)	403 (92%)	34 (8%)	1 (0%)	47	81
1	g	438/449~(98%)	413 (94%)	25~(6%)	0	100	100
1	i	438/449~(98%)	410 (94%)	24 (6%)	4 (1%)	17	56
1	k	438/449~(98%)	413 (94%)	25 (6%)	0	100	100
1	m	438/449~(98%)	393 (90%)	44 (10%)	1 (0%)	47	81
1	О	438/449~(98%)	405 (92%)	31 (7%)	2~(0%)	29	68
1	q	438/449~(98%)	414 (94%)	24 (6%)	0	100	100
1	S	438/449~(98%)	374 (85%)	60 (14%)	4 (1%)	17	56
1	u	438/449~(98%)	407 (93%)	31 (7%)	0	100	100
1	W	438/449~(98%)	418 (95%)	20 (5%)	0	100	100
2	b	428/443~(97%)	393~(92%)	35 (8%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	d	438/443~(99%)	407~(93%)	30 (7%)	1 (0%)	47	81
2	f	428/443~(97%)	398~(93%)	28~(6%)	2~(0%)	29	68
2	h	428/443~(97%)	405 (95%)	23~(5%)	0	100	100
2	j	428/443~(97%)	401 (94%)	27~(6%)	0	100	100
2	1	428/443~(97%)	410 (96%)	18 (4%)	0	100	100
2	n	438/443~(99%)	403 (92%)	34 (8%)	1 (0%)	47	81
2	р	424/443~(96%)	397~(94%)	27~(6%)	0	100	100
2	r	428/443~(97%)	409 (96%)	19 (4%)	0	100	100
2	t	428/443~(97%)	387~(90%)	39~(9%)	2~(0%)	29	68
2	v	428/443~(97%)	406 (95%)	22~(5%)	0	100	100
2	x	428/443~(97%)	407 (95%)	21 (5%)	0	100	100
All	All	10408/10704~(97%)	9668 (93%)	716 (7%)	24 (0%)	50	81

All (24) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	n	432	GLU
1	a	403	ALA
1	s	322	ASP
1	с	266	HIS
1	с	291	ILE
1	0	38	SER
1	m	403	ALA
1	е	266	HIS
1	a	240	ALA
1	i	38	SER
1	i	282	TYR
1	i	403	ALA
1	s	206	ASN
2	t	143	THR
1	0	399	TYR
1	a	399	TYR
1	i	351	PHE
2	f	275	SER
2	f	280	GLN
1	с	399	TYR
1	s	265	ILE
1	s	261	PRO



Continued from previous page...

Mol	Chain	Res	Type
2	d	434	GLY
2	t	175	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	a	369/376~(98%)	364~(99%)	5 (1%)	67	81
1	с	368/376~(98%)	360~(98%)	8 (2%)	52	71
1	е	368/376~(98%)	363~(99%)	5 (1%)	67	81
1	g	370/376~(98%)	366~(99%)	4 (1%)	73	85
1	i	368/376~(98%)	363~(99%)	5 (1%)	67	81
1	k	368/376~(98%)	363~(99%)	5 (1%)	67	81
1	m	369/376~(98%)	365~(99%)	4 (1%)	73	85
1	О	367/376~(98%)	361 (98%)	6 (2%)	62	79
1	q	367/376~(98%)	361~(98%)	6(2%)	62	79
1	s	369/376~(98%)	359~(97%)	10 (3%)	44	66
1	u	369/376~(98%)	364 (99%)	5 (1%)	67	81
1	W	368/376~(98%)	361 (98%)	7 (2%)	57	75
2	b	364/376~(97%)	357~(98%)	7 (2%)	57	75
2	d	363/376~(96%)	358~(99%)	5 (1%)	67	81
2	f	364/376~(97%)	361 (99%)	3 (1%)	81	89
2	h	365/376~(97%)	362 (99%)	3 (1%)	81	89
2	j	365/376~(97%)	359~(98%)	6 (2%)	62	79
2	1	365/376~(97%)	363 (100%)	2 (0%)	88	93
2	n	364/376~(97%)	362 (100%)	2 (0%)	88	93
2	р	363/376~(96%)	358~(99%)	5 (1%)	67	81
2	r	364/376~(97%)	362 (100%)	2 (0%)	88	93
2	t	$\overline{365/376}\ (97\%)$	361 (99%)	4 (1%)	73	85



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	V	365/376~(97%)	364 (100%)	1 (0%)	92 95
2	х	365/376~(97%)	364 (100%)	1 (0%)	92 95
All	All	8792/9024~(97%)	8681 (99%)	111 (1%)	70 82

All (111) residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	Res	Type
1	W	137	VAL
1	W	140	SER
1	W	199	ASP
1	W	254	GLU
1	W	260	VAL
1	W	305	CYS
1	W	329	ASN
2	х	344	TRP
1	u	140	SER
1	u	199	ASP
1	u	254	GLU
1	u	260	VAL
1	u	305	CYS
2	V	344	TRP
1	s	102	ASN
1	s	137	VAL
1	s	140	SER
1	s	174	SER
1	s	199	ASP
1	s	254	GLU
1	s	305	CYS
1	s	373	ARG
1	S	381	SER
1	s	413	MET
2	t	128	ASP
2	t	249	ASP
2	t	344	TRP
2	t	420	SER
1	q	137	VAL
1	q	140	SER
1	q	199	ASP
1	q	254	GLU
1	q	260	VAL
1	q	305	CYS



Mol	Chain	Res	Type
2	r	31	ASP
2	r	344	TRP
1	0	137	VAL
1	0	250	VAL
1	0	254	GLU
1	0	260	VAL
1	0	305	CYS
1	0	337	THR
2	р	31	ASP
2	р	128	ASP
2	р	197	ASP
2	р	328	GLU
2	р	344	TRP
1	m	137	VAL
1	m	140	SER
1	m	254	GLU
1	m	305	CYS
2	n	31	ASP
2	n	344	TRP
1	k	140	SER
1	k	199	ASP
1	k	254	GLU
1	k	260	VAL
1	k	305	CYS
2	1	31	ASP
2	1	344	TRP
1	g	140	SER
1	g	199	ASP
1	g	254	GLU
1	g	305	CYS
2	h	31	ASP
2	h	274	THR
2	h	344	TRP
1	е	140	SER
1	е	254	GLU
1	е	260	VAL
1	е	305	CYS
1	e	438	GLU
2	f	274	THR
2	f	282	ARG
2	f	344	TRP
1	с	140	SER



Mol	Chain	Res	Type
1	С	174	SER
1	с	260	VAL
1	с	277	SER
1	с	300	ASN
1	с	305	CYS
1	с	331	SER
1	с	417	GLU
1	a	140	SER
1	a	199	ASP
1	a	254	GLU
1	a	305	CYS
1	a	432	TYR
2	b	145	SER
2	b	203	ASP
2	b	222	TYR
2	b	249	ASP
2	b	344	TRP
2	b	355	ASP
2	b	368	VAL
2	d	12	CYS
2	d	222	TYR
2	d	249	ASP
2	d	344	TRP
2	d	427	ASP
1	i	140	SER
1	i	254	GLU
1	i	260	VAL
1	i	305	CYS
1	i	438	GLU
2	j	31	ASP
2	j	128	ASP
2	j	197	ASP
2	j	249	ASP
2	j	282	ARG
2	j	344	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (158) such sidechains are listed below:

Mol	Chain	Res	Type
1	W	85	GLN
1	W	206	ASN
1	W	216	ASN



Mol	Chain	Res	Type
1	W	258	ASN
1	W	293	ASN
1	W	342	GLN
1	W	356	ASN
2	х	226	ASN
2	х	247	ASN
2	х	256	ASN
2	Х	292	GLN
2	х	348	ASN
2	х	375	GLN
2	х	423	GLN
1	u	85	GLN
1	u	206	ASN
1	u	258	ASN
1	u	293	ASN
1	u	342	GLN
1	u	356	ASN
2	v	15	GLN
2	v	99	ASN
2	V	105	HIS
2	v	191	GLN
2	V	226	ASN
2	v	247	ASN
2	v	256	ASN
2	V	348	ASN
2	V	375	GLN
2	V	414	ASN
1	S	216	ASN
1	S	293	ASN
1	S	356	ASN
2	t	99	ASN
2	t	204	ASN
2	t	226	ASN
2	t	247	ASN
2	t	292	GLN
2	t	348	ASN
2	t	375	GLN
2	t	414	ASN
1	q	85	GLN
1	q	133	GLN
1	q	206	ASN
1	q	216	ASN



Mol	Chain	Res	Type
1	q	258	ASN
1	q	283	HIS
1	q	293	ASN
1	q	356	ASN
2	r	15	GLN
2	r	134	GLN
2	r	226	ASN
2	r	247	ASN
2	r	256	ASN
2	r	348	ASN
1	0	85	GLN
1	0	102	ASN
1	0	258	ASN
1	0	293	ASN
1	0	356	ASN
2	р	99	ASN
2	p	204	ASN
2	р	226	ASN
2	р	247	ASN
2	р	375	GLN
1	m	85	GLN
1	m	206	ASN
1	m	258	ASN
1	m	293	ASN
1	m	342	GLN
1	m	356	ASN
2	n	99	ASN
2	n	134	GLN
2	n	191	GLN
2	n	226	ASN
2	n	245	GLN
2	n	247	ASN
2	n	256	ASN
2	n	292	GLN
2	n	348	ASN
2	n	414	ASN
1	k	85	GLN
1	k	206	ASN
1	k	258	ASN
1	k	293	ASN
1	k	342	GLN
2	1	134	GLN



Mol	Chain	Res	Type
2	1	226	ASN
2	1	247	ASN
2	1	256	ASN
2	1	348	ASN
1	g	85	GLN
1	g	206	ASN
1	g	258	ASN
1	g	293	ASN
1	g	342	GLN
1	g	356	ASN
2	h	99	ASN
2	h	134	GLN
2	h	226	ASN
2	h	247	ASN
2	h	256	ASN
2	h	348	ASN
1	е	61	HIS
1	е	85	GLN
1	е	206	ASN
1	е	216	ASN
1	е	258	ASN
1	е	293	ASN
1	е	309	HIS
1	е	342	GLN
2	f	8	GLN
2	f	15	GLN
2	f	105	HIS
2	f	134	GLN
2	f	226	ASN
2	f	256	ASN
2	f	292	GLN
2	f	348	ASN
1	с	31	GLN
1	с	85	GLN
1	с	258	ASN
1	с	293	ASN
1	с	342	GLN
1	a	85	GLN
1	a	206	ASN
1	a	258	ASN
1	a	283	HIS
1	a	293	ASN



\mathbf{Mol}	Chain	Res	Type
1	a	342	GLN
2	b	15	GLN
2	b	99	ASN
2	b	105	HIS
2	b	134	GLN
2	b	226	ASN
2	b	247	ASN
2	b	256	ASN
2	b	292	GLN
2	b	348	ASN
2	d	99	ASN
2	d	134	GLN
2	d	226	ASN
2	d	256	ASN
2	d	348	ASN
1	i	85	GLN
1	i	258	ASN
1	i	293	ASN
1	i	380	ASN
2	j	14	ASN
2	j	99	ASN
2	j	105	HIS
2	j	191	GLN
2	j	226	ASN
2	j	256	ASN
2	j	292	GLN
2	j	298	ASN
2	j	348	ASN
2	j	414	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 35 ligands modelled in this entry, 12 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	Bond angles		
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	GTP	u	501	4	26,34,34	0.97	2 (7%)	$32,\!54,\!54$	0.82	0
5	GDP	n	501	-	24,30,30	0.93	1 (4%)	30,47,47	1.48	5 (16%)
5	GDP	f	501	-	24,30,30	0.95	2 (8%)	30,47,47	1.34	5 (16%)
5	GDP	b	501	-	24,30,30	0.95	1 (4%)	30,47,47	1.19	3 (10%)
5	GDP	j	501	-	24,30,30	0.92	1 (4%)	30,47,47	1.29	4 (13%)
3	GTP	q	501	4	26,34,34	0.95	2 (7%)	32,54,54	0.78	0
5	GDP	h	501	-	24,30,30	0.92	1 (4%)	30,47,47	1.34	4 (13%)
3	GTP	0	501	4	26,34,34	0.95	2 (7%)	32,54,54	0.76	0
3	GTP	е	501	4	26,34,34	0.94	2 (7%)	32,54,54	0.82	1 (3%)
3	GTP	a	501	4	26,34,34	0.93	2 (7%)	32,54,54	0.72	0
3	GTP	g	501	4	26,34,34	0.93	2 (7%)	32,54,54	0.77	0
3	GTP	i	501	4	26,34,34	0.93	2 (7%)	32,54,54	0.79	0
5	GDP	r	501	-	24,30,30	0.95	1 (4%)	30,47,47	1.29	4 (13%)
3	GTP	k	501	4	26,34,34	0.95	2 (7%)	32,54,54	0.77	0
3	GTP	С	501	4	26,34,34	0.93	1 (3%)	32,54,54	0.82	1 (3%)
3	GTP	W	501	4	26,34,34	0.96	2 (7%)	32,54,54	0.76	0
5	GDP	1	501	-	24,30,30	0.91	1 (4%)	30,47,47	1.29	5 (16%)
5	GDP	р	501	-	24,30,30	0.93	0	30,47,47	1.36	4 (13%)
3	GTP	s	501	4	26,34,34	0.96	2 (7%)	32,54,54	0.81	0
5	GDP	t	502	-	24,30,30	0.96	2 (8%)	30,47,47	1.56	5 (16%)
5	GDP	v	501	-	24,30,30	0.91	1 (4%)	30,47,47	1.35	5 (16%)
3	GTP	m	501	4	26,34,34	0.97	2 (7%)	32,54,54	0.81	1 (3%)
5	GDP	х	502	-	24,30,30	0.89	0	30,47,47	1.31	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	u	501	4	-	3/18/38/38	0/3/3/3
5	GDP	n	501	-	-	3/12/32/32	0/3/3/3
5	GDP	f	501	-	-	4/12/32/32	0/3/3/3
5	GDP	b	501	-	-	3/12/32/32	0/3/3/3
5	GDP	j	501	-	-	5/12/32/32	0/3/3/3
3	GTP	q	501	4	-	5/18/38/38	0/3/3/3
5	GDP	h	501	-	-	4/12/32/32	0/3/3/3
3	GTP	0	501	4	-	4/18/38/38	0/3/3/3
3	GTP	е	501	4	-	3/18/38/38	0/3/3/3
3	GTP	a	501	4	-	3/18/38/38	0/3/3/3
3	GTP	g	501	4	-	3/18/38/38	0/3/3/3
3	GTP	i	501	4	-	5/18/38/38	0/3/3/3
5	GDP	r	501	-	-	4/12/32/32	0/3/3/3
3	GTP	k	501	4	-	3/18/38/38	0/3/3/3
3	GTP	С	501	4	-	5/18/38/38	0/3/3/3
3	GTP	W	501	4	-	5/18/38/38	0/3/3/3
5	GDP	1	501	-	-	4/12/32/32	0/3/3/3
5	GDP	р	501	-	-	1/12/32/32	0/3/3/3
3	GTP	s	501	4	-	3/18/38/38	0/3/3/3
5	GDP	t	502	-	-	3/12/32/32	0/3/3/3
5	GDP	v	501	-	-	2/12/32/32	0/3/3/3
3	GTP	m	501	4	-	4/18/38/38	0/3/3/3
5	GDP	Х	502	-	-	6/12/32/32	0/3/3/3

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	m	501	GTP	C5-C6	-2.75	1.41	1.47
3	u	501	GTP	C5-C6	-2.68	1.42	1.47
3	W	501	GTP	C5-C6	-2.66	1.42	1.47
3	s	501	GTP	C5-C6	-2.61	1.42	1.47
3	q	501	GTP	C5-C6	-2.58	1.42	1.47
3	a	501	GTP	C5-C6	-2.57	1.42	1.47
3	k	501	GTP	C5-C6	-2.54	1.42	1.47
3	0	501	GTP	C5-C6	-2.53	1.42	1.47
3	i	501	GTP	C5-C6	-2.51	1.42	1.47



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	g	501	GTP	C5-C6	-2.47	1.42	1.47
5	b	501	GDP	C6-N1	-2.44	1.34	1.37
3	с	501	GTP	C5-C6	-2.42	1.42	1.47
5	\mathbf{t}	502	GDP	C6-N1	-2.41	1.34	1.37
3	е	501	GTP	C5-C6	-2.41	1.42	1.47
3	е	501	GTP	C8-N7	-2.23	1.31	1.35
5	f	501	GDP	C6-N1	-2.20	1.34	1.37
5	1	501	GDP	C6-N1	-2.20	1.34	1.37
5	n	501	GDP	C6-N1	-2.19	1.34	1.37
3	a	501	GTP	C8-N7	-2.16	1.31	1.35
3	m	501	GTP	C8-N7	-2.16	1.31	1.35
3	k	501	GTP	C8-N7	-2.15	1.31	1.35
3	s	501	GTP	C8-N7	-2.14	1.31	1.35
3	u	501	GTP	C8-N7	-2.13	1.31	1.35
3	W	501	GTP	C8-N7	-2.12	1.31	1.35
5	V	501	GDP	C6-N1	-2.12	1.34	1.37
5	r	501	GDP	C6-N1	-2.10	1.34	1.37
3	i	501	GTP	C8-N7	-2.09	1.31	1.35
3	0	501	GTP	C8-N7	-2.09	1.31	1.35
3	g	501	GTP	C8-N7	-2.09	1.31	1.35
3	q	501	GTP	C8-N7	-2.08	1.31	1.35
5	\mathbf{t}	502	GDP	O4'-C1'	2.07	1.44	1.41
5	j	501	GDP	C6-N1	-2.06	1.34	1.37
5	h	501	GDP	C6-N1	-2.05	1.34	1.37
5	f	501	GDP	C2-N3	2.03	1.38	1.33

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	t	502	GDP	PA-O3A-PB	-3.91	119.42	132.83
5	n	501	GDP	PA-O3A-PB	-3.90	119.43	132.83
5	t	502	GDP	C3'-C2'-C1'	3.62	106.43	100.98
5	V	501	GDP	C3'-C2'-C1'	3.46	106.19	100.98
5	h	501	GDP	C3'-C2'-C1'	3.35	106.02	100.98
5	р	501	GDP	C3'-C2'-C1'	3.34	106.00	100.98
5	n	501	GDP	C3'-C2'-C1'	3.26	105.88	100.98
5	р	501	GDP	PA-O3A-PB	-3.21	121.80	132.83
5	l	501	GDP	PA-O3A-PB	-3.16	121.99	132.83
5	r	501	GDP	PA-O3A-PB	-3.14	122.04	132.83
5	t	502	GDP	C5-C6-N1	3.13	119.47	113.95
5	Х	502	GDP	PA-O3A-PB	-3.10	122.20	132.83
5	V	501	GDP	PA-O3A-PB	-3.08	122.26	132.83



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	h	501	GDP	PA-O3A-PB	-3.03	122.44	132.83
5	j	501	GDP	PA-O3A-PB	-2.94	122.72	132.83
5	f	501	GDP	C5-C6-N1	2.92	119.11	113.95
5	b	501	GDP	C8-N7-C5	2.89	108.50	102.99
5	f	501	GDP	C3'-C2'-C1'	2.81	105.22	100.98
5	f	501	GDP	PA-O3A-PB	-2.80	123.22	132.83
5	b	501	GDP	C3'-C2'-C1'	2.80	105.19	100.98
5	f	501	GDP	C8-N7-C5	2.75	108.23	102.99
5	t	502	GDP	O6-C6-C5	-2.70	119.09	124.37
5	Х	502	GDP	C5-C6-N1	2.66	118.64	113.95
5	r	501	GDP	C3'-C2'-C1'	2.65	104.97	100.98
5	n	501	GDP	O6-C6-C5	-2.55	119.40	124.37
5	р	501	GDP	C8-N7-C5	2.54	107.83	102.99
5	b	501	GDP	C5-C6-N1	2.53	118.42	113.95
5	l	501	GDP	C5-C6-N1	2.52	118.40	113.95
5	V	501	GDP	C5-C6-N1	2.48	118.34	113.95
5	n	501	GDP	C5-C6-N1	2.48	118.32	113.95
5	r	501	GDP	C5-C6-N1	2.47	118.31	113.95
5	h	501	GDP	C8-N7-C5	2.47	107.69	102.99
5	j	501	GDP	C8-N7-C5	2.47	107.69	102.99
5	r	501	GDP	C8-N7-C5	2.41	107.58	102.99
5	р	501	GDP	C5-C6-N1	2.38	118.16	113.95
5	l	501	GDP	C3'-C2'-C1'	2.38	104.56	100.98
5	1	501	GDP	C8-N7-C5	2.37	107.50	102.99
5	Х	502	GDP	C3'-C2'-C1'	2.34	104.50	100.98
5	Х	502	GDP	O6-C6-C5	-2.31	119.85	124.37
5	Х	502	GDP	C8-N7-C5	2.29	107.36	102.99
5	j	501	GDP	C3'-C2'-C1'	2.29	104.42	100.98
5	h	501	GDP	C5-C6-N1	2.28	117.98	113.95
5	V	501	GDP	C8-N7-C5	2.28	107.33	102.99
5	j	501	GDP	C5-C6-N1	2.26	117.95	113.95
5	t	502	GDP	C8-N7-C5	2.24	107.27	102.99
5	V	501	GDP	O6-C6-C5	-2.10	120.26	124.37
5	n	501	GDP	C8-N7-C5	$2.1\overline{0}$	107.00	102.99
3	с	501	GTP	O6-C6-C5	2.06	128.40	124.37
5	1	501	GDP	O6-C6-C5	-2.05	120.37	124.37
3	m	501	GTP	O6-C6-C5	2.04	128.35	124.37
3	е	501	GTP	O6-C6-C5	2.02	128.31	124.37
5	f	501	GDP	06-C6-C5	-2.01	120.45	124.37

There are no chirality outliers.

All (85) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	W	501	GTP	C5'-O5'-PA-O3A
3	u	501	GTP	C5'-O5'-PA-O3A
3	s	501	GTP	C5'-O5'-PA-O3A
3	s	501	GTP	C5'-O5'-PA-O2A
3	q	501	GTP	C5'-O5'-PA-O3A
3	q	501	GTP	C5'-O5'-PA-O2A
3	0	501	GTP	C5'-O5'-PA-O1A
3	0	501	GTP	C5'-O5'-PA-O2A
3	m	501	GTP	C5'-O5'-PA-O3A
3	m	501	GTP	C5'-O5'-PA-O2A
3	k	501	GTP	C5'-O5'-PA-O3A
3	g	501	GTP	C5'-O5'-PA-O3A
3	e	501	GTP	C5'-O5'-PA-O3A
3	с	501	GTP	PB-O3B-PG-O2G
3	с	501	GTP	C5'-O5'-PA-O3A
3	с	501	GTP	C5'-O5'-PA-O2A
3	a	501	GTP	C5'-O5'-PA-O3A
3	a	501	GTP	C5'-O5'-PA-O2A
3	i	501	GTP	C5'-O5'-PA-O1A
3	i	501	GTP	C5'-O5'-PA-O2A
5	X	502	GDP	C5'-O5'-PA-O2A
5	t	502	GDP	C5'-O5'-PA-O1A
5	t	502	GDP	C5'-O5'-PA-O2A
5	j	501	GDP	C5'-O5'-PA-O2A
5	h	501	GDP	O4'-C4'-C5'-O5'
5	h	501	GDP	C3'-C4'-C5'-O5'
5	r	501	GDP	O4'-C4'-C5'-O5'
5	n	501	GDP	O4'-C4'-C5'-O5'
5	f	501	GDP	O4'-C4'-C5'-O5'
5	r	501	GDP	C3'-C4'-C5'-O5'
5	n	501	GDP	C3'-C4'-C5'-O5'
5	V	501	GDP	O4'-C4'-C5'-O5'
5	1	501	GDP	O4'-C4'-C5'-O5'
5	f	501	GDP	C3'-C4'-C5'-O5'
5	1	501	GDP	C3'-C4'-C5'-O5'
5	Х	502	GDP	O4'-C4'-C5'-O5'
5	X	502	GDP	C3'-C4'-C5'-O5'
5	V	501	GDP	C3'-C4'-C5'-O5'
3	е	501	GTP	C4'-C5'-O5'-PA
3	a	501	GTP	C4'-C5'-O5'-PA
3	0	501	GTP	C5'-O5'-PA-O3A
5	X	502	GDP	C5'-O5'-PA-O3A
5	j	501	GDP	C5'-O5'-PA-O3A



Mol	Chain	Res		Atoms
5	x	502	GDP	PB-O3A-PA-O1A
5	r	501	GDP	PB-O3A-PA-O1A
5	p	501	GDP	PB-O3A-PA-O1A
5	f	501	GDP	PB-O3A-PA-O1A
5	b	501	GDP	PB-O3A-PA-O1A
5	i	501	GDP	PB-O3A-PA-O1A
3	W	501	GTP	C5'-O5'-PA-O2A
3	u	501	GTP	C5'-O5'-PA-O2A
3	k	501	GTP	C5'-O5'-PA-O2A
3	g	501	GTP	C5'-O5'-PA-O2A
3	e	501	GTP	C5'-O5'-PA-O2A
5	j	501	GDP	C5'-O5'-PA-O1A
3	q	501	GTP	C4'-C5'-O5'-PA
3	s	501	GTP	PB-O3A-PA-O2A
3	0	501	GTP	PB-O3A-PA-O2A
3	с	501	GTP	PB-O3A-PA-O2A
5	h	501	GDP	PB-O3A-PA-O1A
3	g	501	GTP	C4'-C5'-O5'-PA
3	u	501	GTP	C4'-C5'-O5'-PA
3	k	501	GTP	C4'-C5'-O5'-PA
3	q	501	GTP	PB-O3A-PA-O2A
5	n	501	GDP	PB-O3A-PA-O1A
3	i	501	GTP	C5'-O5'-PA-O3A
5	t	502	GDP	C5'-O5'-PA-O3A
5	1	501	GDP	C5'-O5'-PA-O3A
5	j	501	GDP	C3'-C4'-C5'-O5'
3	W	501	GTP	PB-O3A-PA-O1A
3	W	501	GTP	PB-O3A-PA-O2A
3	q	501	GTP	PB-O3A-PA-O1A
3	m	501	GTP	PB-O3A-PA-O1A
3	m	501	GTP	PB-O3A-PA-O2A
3	i	501	GTP	PB-O3A-PA-O2A
5	r	501	GDP	PB-O3A-PA-O2A
5	h	501	GDP	PB-O3A-PA-O2A
5	f	501	GDP	PB-O3A-PA-O2A
5	b	501	GDP	PB-O3A-PA-O2A
5	X	502	GDP	C5'-O5'-PA-O1A
5		501	GDP	C5'-O5'-PA-O1A
5	b	501	GDP	<u>U4'-C4'-C5'-O5'</u>
3	С	501	GTP	PB-03B-PG-01G
3	W .	501	GTP	C4'-C5'-O5'-PA
3	i	501	GTP	C4'-C5'-O5'-PA

Continued from previous page...



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-22677. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 199



Y Index: 179



Z Index: 168



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1Primary map



X Index: 222

Y Index: 223

Z Index: 191

The images above show the largest variance slices of the map in three orthogonal directions.

Orthogonal surface views (i) 6.4

6.4.1**Primary** map



The images above show the 3D surface view of the map at the recommended contour level 0.8. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 1003 $\rm nm^3;$ this corresponds to an approximate mass of 906 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-22677 and PDB model 7N32. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.8 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.8).



9.4 Atom inclusion (i)



At the recommended contour level, 64% of all backbone atoms, 41% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4085	0.2820
a	0.4075	0.1760
b	0.4491	0.2160
С	0.3283	0.1470
d	0.3561	0.1580
е	0.4222	0.2070
f	0.4241	0.2020
g	0.4558	0.3130
h	0.4752	0.3370
i	0.3610	0.3110
j	0.3554	0.3090
k	0.4202	0.3150
1	0.4404	0.3140
m	0.4437	0.3000
n	0.4744	0.3080
0	0.3415	0.3320
р	0.3821	0.3570
q	0.4163	0.3220
r	0.4346	0.3120
S	0.4305	0.3040
t	0.4700	0.3130
u	0.3993	0.2990
V	0.4089	0.3090
W	0.3238	0.2990
Х	0.3839	0.3030

