



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 09:26 PM EDT

PDB ID : 5N4K  
Title : N-terminal domain of a human Coronavirus NL63 nucleocapsid protein  
Authors : Zdzalik, M.; Szelazek, B.; Kabala, W.; Golik, P.; Burmistrz, M.; Florek, D.;  
Kus, K.; Pyrc, K.; Dubin, G.  
Deposited on : 2017-02-10  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	
1	B	150	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	406	-	-	X	-

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 2645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	130	1114	727	197	186	4	0	13	0
1	B	129	1091	706	196	186	3	0	12	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	HIS	-	expression tag	UNP Q6Q1R8
A	-8	HIS	-	expression tag	UNP Q6Q1R8
A	-7	HIS	-	expression tag	UNP Q6Q1R8
A	-6	HIS	-	expression tag	UNP Q6Q1R8
A	-5	SER	-	expression tag	UNP Q6Q1R8
A	-4	GLY	-	expression tag	UNP Q6Q1R8
A	-3	ASP	-	expression tag	UNP Q6Q1R8
A	-2	ASP	-	expression tag	UNP Q6Q1R8
A	-1	ASP	-	expression tag	UNP Q6Q1R8
A	0	ASP	-	expression tag	UNP Q6Q1R8
A	1	LYS	-	expression tag	UNP Q6Q1R8
B	-9	HIS	-	expression tag	UNP Q6Q1R8
B	-8	HIS	-	expression tag	UNP Q6Q1R8
B	-7	HIS	-	expression tag	UNP Q6Q1R8
B	-6	HIS	-	expression tag	UNP Q6Q1R8
B	-5	SER	-	expression tag	UNP Q6Q1R8
B	-4	GLY	-	expression tag	UNP Q6Q1R8
B	-3	ASP	-	expression tag	UNP Q6Q1R8
B	-2	ASP	-	expression tag	UNP Q6Q1R8
B	-1	ASP	-	expression tag	UNP Q6Q1R8
B	0	ASP	-	expression tag	UNP Q6Q1R8
B	1	LYS	-	expression tag	UNP Q6Q1R8

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 10 8 2	0	1
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	187	Total 192	O 192	0	5
3	B	169	Total 173	O 173	0	4


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoprotein



- Molecule 1: Nucleoprotein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.09Å 48.09Å 101.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.65 – 1.49 41.65 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.65-1.49) 99.1 (41.65-1.49)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.78 (at 1.49Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.151 , 0.197 0.152 , 0.197	Depositor DCC
$R_{free}$ test set	2132 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.0	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l 0.479 for h,-h-k,-l 0.023 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2645	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.17	2/1184 (0.2%)	1.32	10/1599 (0.6%)
1	B	1.23	4/1149 (0.3%)	1.36	12/1552 (0.8%)
All	All	1.20	6/2333 (0.3%)	1.34	22/3151 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	79	TYR	CZ-OH	6.46	1.48	1.37
1	B	120[A]	GLN	CG-CD	5.68	1.64	1.51
1	B	120[B]	GLN	CG-CD	5.68	1.64	1.51
1	A	79	TYR	CE2-CZ	-5.64	1.31	1.38
1	A	128	SER	CB-OG	-5.58	1.34	1.42
1	B	79	TYR	CE2-CZ	-5.14	1.31	1.38

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	61	ARG	NE-CZ-NH1	15.18	127.89	120.30
1	A	92	ARG	NE-CZ-NH2	-15.04	112.78	120.30
1	B	61	ARG	NE-CZ-NH1	14.53	127.57	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	92	ARG	NE-CZ-NH2	-12.88	113.86	120.30
1	B	29[A]	SER	C-N-CA	8.57	143.13	121.70
1	B	29[B]	SER	C-N-CA	8.57	143.13	121.70
1	B	61	ARG	NE-CZ-NH2	-7.43	116.59	120.30
1	B	116	ARG	NE-CZ-NH2	7.25	123.93	120.30
1	A	61	ARG	NE-CZ-NH2	-7.09	116.75	120.30
1	B	23[A]	MET	CG-SD-CE	6.21	110.13	100.20
1	B	23[B]	MET	CG-SD-CE	6.21	110.13	100.20
1	B	30	ASP	N-CA-CB	5.89	121.20	110.60
1	A	120[A]	GLN	CA-CB-CG	5.63	125.79	113.40
1	A	120[B]	GLN	CA-CB-CG	5.63	125.79	113.40
1	A	88[A]	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	88[B]	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	30	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	79	TYR	CD1-CE1-CZ	-5.26	115.07	119.80
1	A	126	LYS	CD-CE-NZ	5.13	123.50	111.70
1	B	92	ARG	NH1-CZ-NH2	5.05	124.95	119.40
1	B	120[A]	GLN	CA-CB-CG	5.02	124.44	113.40
1	B	120[B]	GLN	CA-CB-CG	5.02	124.44	113.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	9	ASP	Peptide
1	B	29[B]	SER	Mainchain,Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1114	0	1158	27	0
1	B	1091	0	1125	37	0
2	A	45	0	0	1	0
2	B	30	0	0	7	0
3	A	192	0	0	12	0
3	B	173	0	0	7	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2645	0	2283	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35[A]:ARG:HB2	1:B:35[A]:ARG:NH2	1.29	1.47
1:B:35[A]:ARG:HH21	1:B:35[A]:ARG:CB	1.35	1.35
1:B:29[B]:SER:HB2	1:B:31[B]:LYS:H	0.98	1.13
1:B:29[A]:SER:HB3	1:B:30:ASP:CB	1.80	1.12
1:B:28[A]:SER:O	1:B:29[A]:SER:OG	1.76	1.04
1:B:29[A]:SER:CB	1:B:30:ASP:HB3	1.90	1.00
2:B:406:SO4:O2	3:B:502[A]:HOH:O	1.76	1.00
1:B:104[B]:GLU:OE1	2:B:406:SO4:O4	1.85	0.95
1:B:29[A]:SER:HB3	1:B:30:ASP:HB2	1.47	0.92
1:B:29[A]:SER:CB	1:B:30:ASP:CB	2.47	0.91
1:A:104[A]:GLU:HG2	3:A:657:HOH:O	1.73	0.87
1:B:29[A]:SER:HB2	1:B:30:ASP:HB3	1.56	0.85
1:A:104[A]:GLU:CG	3:A:657:HOH:O	2.26	0.83
1:B:29[A]:SER:HB3	1:B:30:ASP:HB3	1.52	0.81
1:B:29[B]:SER:CB	1:B:31[B]:LYS:H	1.88	0.81
1:A:77:HIS:HD2	3:A:540:HOH:O	1.68	0.77
1:B:103:LYS:HA	2:B:406:SO4:O1	1.85	0.74
1:A:104[B]:GLU:HG2	3:A:502:HOH:O	1.88	0.73
1:B:77:HIS:HD2	3:B:542:HOH:O	1.72	0.72
1:A:40:ASN:ND2	2:A:408:SO4:O1	2.25	0.69
1:B:117:LYS:HB2	1:B:120[B]:GLN:HG3	1.74	0.69
1:B:40:ASN:ND2	2:B:406:SO4:O3	2.18	0.68
1:A:117:LYS:HB2	1:A:120[A]:GLN:HG2	1.76	0.66
1:A:23[B]:MET:HE2	1:A:124:GLU:HA	1.77	0.66
1:B:29[A]:SER:N	3:B:501:HOH:O	2.28	0.66
1:A:23[B]:MET:HE3	1:A:137:VAL:CG1	2.26	0.66
2:B:406:SO4:O4	3:B:504:HOH:O	2.13	0.63
1:B:23[A]:MET:HE2	1:B:124:GLU:HA	1.80	0.62
1:B:35[A]:ARG:HH21	1:B:35[A]:ARG:CG	2.12	0.61
1:A:139:GLU:OE2	1:A:140:PHE:N	2.34	0.60
1:B:104[B]:GLU:OE1	2:B:406:SO4:S	2.60	0.58
1:B:35[A]:ARG:HB2	1:B:35[A]:ARG:HH21	0.47	0.55
1:B:23[A]:MET:HE3	1:B:137:VAL:CG1	2.38	0.54

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:ARG:NH1	3:A:507:HOH:O	2.33	0.53
1:B:35[A]:ARG:NH2	1:B:35[A]:ARG:CB	2.19	0.53
1:A:23[B]:MET:CE	1:A:137:VAL:CG1	2.87	0.53
1:B:117:LYS:HB2	1:B:120[B]:GLN:CG	2.39	0.53
1:A:23[B]:MET:HE3	1:A:138:VAL:O	2.10	0.52
1:A:23[B]:MET:HE3	1:A:137:VAL:HG12	1.91	0.51
1:B:77:HIS:CD2	3:B:542:HOH:O	2.54	0.51
1:A:23[B]:MET:CE	1:A:137:VAL:HG11	2.42	0.49
1:A:116:ARG:HG2	1:A:120[A]:GLN:HG3	1.94	0.49
1:B:35[A]:ARG:NH2	1:B:35[A]:ARG:CG	2.73	0.48
1:B:29[A]:SER:OG	3:B:503:HOH:O	1.98	0.47
1:B:48[A]:ASN:OD1	1:B:50[A]:ASP:OD1	2.31	0.47
1:A:87[B]:LYS:HB3	1:A:87[B]:LYS:HE3	1.31	0.46
1:A:77:HIS:CD2	3:A:540:HOH:O	2.51	0.45
1:A:13:ARG:HB2	3:A:615:HOH:O	2.17	0.44
1:A:48[A]:ASN:OD1	1:A:50[A]:ASP:OD1	2.35	0.44
1:B:23[A]:MET:CE	1:B:124:GLU:HA	2.47	0.44
1:B:116:ARG:HG2	1:B:120[A]:GLN:HB3	2.00	0.44
1:B:107[A]:LYS:HB3	1:B:107[A]:LYS:HE2	1.81	0.44
1:A:31[B]:LYS:HD2	3:A:510:HOH:O	2.18	0.43
1:A:82:GLY:O	1:A:87[B]:LYS:HG3	2.18	0.43
1:B:23[A]:MET:HE3	1:B:137:VAL:HG12	1.99	0.43
1:B:23[A]:MET:CE	1:B:137:VAL:CG1	2.97	0.43
1:A:29:SER:HB3	3:A:510:HOH:O	2.19	0.42
1:A:50[A]:ASP:HB2	1:A:107[A]:LYS:HG3	2.02	0.42
1:A:140:PHE:CD2	1:A:140:PHE:C	2.92	0.42
1:B:23[A]:MET:CE	1:B:137:VAL:HG11	2.49	0.41
1:A:23[B]:MET:CE	1:A:124:GLU:HA	2.49	0.41
1:A:30:ASP:N	3:A:510:HOH:O	2.53	0.40
1:A:115:ASN:HA	3:A:533:HOH:O	2.21	0.40
1:B:103:LYS:CA	2:B:406:SO4:O1	2.64	0.40
1:B:29[A]:SER:CB	3:B:503:HOH:O	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/150 (93%)	133 (96%)	5 (4%)	1 (1%)	22	6
1	B	137/150 (91%)	132 (96%)	2 (2%)	3 (2%)	6	1
All	All	276/300 (92%)	265 (96%)	7 (2%)	4 (1%)	13	1

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	30	ASP
1	A	29	SER
1	B	29[A]	SER
1	B	29[B]	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/132 (93%)	115 (94%)	8 (6%)	17	1
1	B	119/132 (90%)	116 (98%)	3 (2%)	47	18
All	All	242/264 (92%)	231 (96%)	11 (4%)	39	5

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35[A]	ARG
1	A	35[B]	ARG
1	A	107[A]	LYS
1	A	107[B]	LYS
1	A	120[A]	GLN
1	A	120[B]	GLN
1	A	139	GLU
1	A	140	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	35[A]	ARG
1	B	35[B]	ARG
1	B	63	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	B	77	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	404	-	4,4,4	0.28	0	6,6,6	1.04	0
2	SO4	B	402	-	4,4,4	0.25	0	6,6,6	0.46	0
2	SO4	B	403	-	4,4,4	0.56	0	6,6,6	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	406	-	4,4,4	1.08	1 (25%)	6,6,6	2.64	2 (33%)
2	SO4	A	407[B]	-	4,4,4	0.21	0	6,6,6	1.21	1 (16%)
2	SO4	A	408	-	4,4,4	0.63	0	6,6,6	1.39	1 (16%)
2	SO4	B	404	-	4,4,4	0.43	0	6,6,6	0.38	0
2	SO4	A	405	-	4,4,4	0.73	0	6,6,6	1.48	1 (16%)
2	SO4	A	401	-	4,4,4	0.14	0	6,6,6	0.66	0
2	SO4	A	403	-	4,4,4	0.36	0	6,6,6	0.80	0
2	SO4	A	407[A]	-	4,4,4	0.18	0	6,6,6	1.05	0
2	SO4	B	405	-	4,4,4	0.69	0	6,6,6	1.50	1 (16%)
2	SO4	A	406	-	4,4,4	0.37	0	6,6,6	0.66	0
2	SO4	B	401	-	4,4,4	0.26	0	6,6,6	0.56	0
2	SO4	A	402	-	4,4,4	0.26	0	6,6,6	0.61	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	406	SO4	O4-S	-2.05	1.31	1.47

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	406	SO4	O4-S-O3	-5.85	84.10	109.06
2	A	408	SO4	O4-S-O3	-2.79	97.16	109.06
2	A	405	SO4	O4-S-O3	-2.70	97.54	109.06
2	B	405	SO4	O4-S-O3	2.65	120.37	109.06
2	A	407[B]	SO4	O4-S-O1	2.31	121.38	109.31
2	B	406	SO4	O2-S-O1	2.18	125.51	109.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	406	SO4	7	0
2	A	408	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	130/150 (86%)	-0.47	2 (1%) 73 78	9, 16, 50, 75	3 (2%)
1	B	129/150 (86%)	-0.50	2 (1%) 72 77	9, 16, 49, 71	0
All	All	259/300 (86%)	-0.49	4 (1%) 73 78	9, 16, 50, 75	3 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	ALA	6.9
1	B	140	PHE	4.1
1	B	11	ALA	3.5
1	A	64[A]	MET	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	402	5/5	0.91	0.19	29,31,35,36	5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	403	5/5	0.91	0.20	34,37,44,46	5
2	SO4	A	406	5/5	0.92	0.15	48,49,50,53	5
2	SO4	A	404	5/5	0.92	0.11	35,36,37,38	5
2	SO4	A	407[B]	5/5	0.93	0.12	21,21,24,25	5
2	SO4	A	407[A]	5/5	0.93	0.12	21,22,23,25	5
2	SO4	B	402	5/5	0.94	0.12	32,32,38,39	5
2	SO4	B	401	5/5	0.94	0.12	34,35,37,39	5
2	SO4	A	403	5/5	0.95	0.09	30,32,35,36	5
2	SO4	A	408	5/5	0.96	0.10	15,20,26,31	5
2	SO4	A	401	5/5	0.96	0.11	45,46,52,54	5
2	SO4	B	405	5/5	0.96	0.07	19,26,32,34	5
2	SO4	B	404	5/5	0.97	0.05	40,41,44,46	5
2	SO4	A	405	5/5	0.97	0.08	19,26,32,36	5
2	SO4	B	406	5/5	0.98	0.09	20,27,35,38	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.