

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 08:50 AM EDT

PDB ID : 6N45

Title: Crystal structure of the cryptic polo box domain of human activated Plk4

variant 1

Authors: Zhang, L.; Park, J.-E.; Meng, L.; Lee, K.S.

Deposited on : 2018-11-17

Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Chimera protein of Serine/threonine-protein kinase PLK4 and DDB1- and CUL4-associated factor 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	207	Total 1486	_	N 247	0	S	0	0	0
			1400	945	241	292	4			
1	В	200	Total	С	Ν	Ο	S	0	0	0
			1447	927	237	278	5			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	577	GLY	-	expression tag	UNP O00444
A	578	ALA	-	expression tag	UNP O00444
A	579	HIS	-	expression tag	UNP O00444
A	580	MET	-	expression tag	UNP O00444
A	698	GLU	SER	conflict	UNP O00444
A	700	GLU	SER	conflict	UNP O00444
A	704	GLU	THR	conflict	UNP O00444
A	707	ASP	THR	conflict	UNP O00444
A	809	VAL	-	linker	UNP O00444
В	577	GLY	-	expression tag	UNP O00444
В	578	ALA	-	expression tag	UNP O00444
В	579	HIS	-	expression tag	UNP O00444
В	580	MET	-	expression tag	UNP O00444
В	698	GLU	SER	conflict	UNP O00444
В	700	GLU	SER	conflict	UNP O00444
В	704	GLU	THR	conflict	UNP O00444
В	707	ASP	THR	conflict	UNP O00444
В	809	VAL	_	linker	UNP O00444

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	P 32	Depositor	
Cell constants	61.66Å 61.66Å 137.28Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor	
Resolution (Å)	34.77 - 2.64	Depositor	
% Data completeness	99.9 (34.77-2.64)	Depositor	
(in resolution range)	,	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.55 (at 2.65Å)	Xtriage	
Refinement program	REFMAC 5.8.0238	Depositor	
R, R_{free}	0.234 , 0.314	Depositor	
Wilson B-factor (Å ²)	81.6	Xtriage	
Anisotropy	0.045	Xtriage	
L-test for twinning ²	$< L >=0.52, < L^2>=0.36$	Xtriage	
	0.015 for -h,-k,l		
Estimated twinning fraction	0.487 for h,-h-k,-l	Xtriage	
	0.018 for -k,-h,-l		
Total number of atoms	2933	wwPDB-VP	
Average B, all atoms (Å ²)	82.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $^{^1 {\}rm Intensities}$ estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

