



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 3, 2024 – 09:29 PM EST

PDB ID : 1N7V  
Title : THE RECEPTOR-BINDING PROTEIN P2 OF BACTERIOPHAGE PRD1:  
CRYSTAL FORM III  
Authors : Xu, L.; Benson, S.D.; Butcher, S.J.; Bamford, D.H.; Burnett, R.M.  
Deposited on : 2002-11-18  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adsorption protein P2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	533	3990	2560	652	768	10	0	0	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0
2	A	1	4	2	2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	234	Total 234	O 234	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adsorption protein P2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.50Å 270.20Å 137.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.11 – 2.20 45.83 – 2.20	Depositor EDS
% Data completeness (in resolution range)	78.4 (38.11-2.20) 78.5 (45.83-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.20Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.229 , 0.262 0.225 , 0.262	Depositor DCC
$R_{free}$ test set	1937 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtrriage
Anisotropy	1.175	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4233	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/4106	0.66	5/5631 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	TRP	N-CA-C	-6.88	92.41	111.00
1	A	272	PRO	N-CA-CB	5.90	110.38	103.30
1	A	244	PRO	N-CA-CB	5.57	109.98	103.30
1	A	276	PRO	N-CA-CB	5.24	109.59	103.30
1	A	475	TYR	N-CA-C	-5.09	97.27	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3990	0	3746	98	0
2	A	8	0	6	0	0
3	A	1	0	0	0	0
4	A	234	0	0	2	0
All	All	4233	0	3752	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:512:ILE:HG13	1:A:513:ALA:H	1.24	0.99
1:A:465:SER:O	1:A:479:ALA:HB3	1.76	0.84
1:A:464:SER:HB3	1:A:481:THR:HG22	1.63	0.78
1:A:435:ARG:HB2	1:A:448:GLN:HG3	1.66	0.77
1:A:201:ASN:HD22	1:A:204:TRP:HA	1.51	0.76
1:A:204:TRP:HB3	1:A:205:PRO:HD3	1.70	0.73
1:A:291:ASN:HD21	1:A:293:MET:HB2	1.56	0.71
1:A:512:ILE:HG13	1:A:513:ALA:N	2.03	0.71
1:A:182:VAL:H	1:A:234:GLN:NE2	1.89	0.69
1:A:102:ARG:HG3	1:A:102:ARG:HH11	1.57	0.69
1:A:331:PHE:CZ	1:A:334:ASP:HA	2.28	0.69
1:A:238:TYR:CE2	1:A:302:PRO:HB3	2.28	0.69
1:A:313:LEU:HD12	1:A:313:LEU:H	1.58	0.69
1:A:426:LYS:HB2	1:A:430:LEU:HD23	1.75	0.68
1:A:116:PRO:HG3	1:A:131:GLN:HE22	1.59	0.67
1:A:161:THR:HG22	1:A:171:GLY:CA	2.25	0.67
1:A:92:ASN:HD22	1:A:92:ASN:C	1.99	0.65
1:A:227:ALA:HB1	4:A:629:HOH:O	1.97	0.65
1:A:465:SER:O	1:A:479:ALA:CB	2.46	0.64
1:A:117:ASN:HD22	1:A:120:SER:H	1.47	0.63
1:A:242:GLU:CB	1:A:298:GLY:HA2	2.28	0.63
1:A:469:TYR:CZ	1:A:524:ALA:HA	2.34	0.62
1:A:52:LEU:HD13	1:A:328:ASN:HA	1.80	0.62
1:A:478:GLY:O	1:A:479:ALA:HB3	1.99	0.62
1:A:102:ARG:HG3	1:A:110:ASP:O	2.00	0.62
1:A:92:ASN:ND2	1:A:95:SER:H	1.99	0.60
1:A:204:TRP:CB	1:A:205:PRO:HD3	2.32	0.58
1:A:164:ASP:HB3	1:A:167:THR:HB	1.85	0.58
1:A:313:LEU:HD12	1:A:313:LEU:N	2.18	0.58
1:A:335:GLU:O	1:A:336:VAL:HB	2.04	0.57
1:A:478:GLY:O	1:A:479:ALA:CB	2.53	0.56
1:A:331:PHE:O	1:A:526:ARG:NH2	2.39	0.56
1:A:57:SER:HB2	1:A:346:TYR:CD1	2.41	0.55
1:A:468:GLY:HA2	1:A:477:ALA:HA	1.89	0.55
1:A:283:ILE:HG22	1:A:284:ILE:N	2.22	0.54
1:A:204:TRP:O	1:A:207:LEU:N	2.40	0.54
1:A:454:LEU:N	1:A:454:LEU:HD23	2.22	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:431:GLU:OE2	1:A:451:LEU:HD21	2.07	0.54
1:A:163:PRO:HG2	1:A:286:VAL:HG21	1.89	0.54
1:A:111:GLU:OE2	1:A:136:HIS:HE1	1.91	0.53
1:A:469:TYR:CE1	1:A:524:ALA:HA	2.44	0.53
1:A:102:ARG:HG3	1:A:102:ARG:NH1	2.23	0.53
1:A:38:ILE:HG21	1:A:479:ALA:HB2	1.93	0.50
1:A:92:ASN:HD21	1:A:95:SER:H	1.60	0.50
1:A:57:SER:HB2	1:A:346:TYR:CE1	2.46	0.50
1:A:530:MET:HE2	1:A:553:PRO:HB3	1.93	0.50
1:A:482:PHE:CD1	1:A:487:ASN:HB3	2.47	0.49
1:A:203:ASN:HD22	1:A:207:LEU:HD12	1.77	0.49
1:A:518:ASP:OD2	1:A:549:ARG:NH2	2.44	0.49
1:A:283:ILE:CG2	1:A:284:ILE:N	2.75	0.49
1:A:133:THR:O	1:A:142:ASN:HB3	2.13	0.49
1:A:291:ASN:C	1:A:291:ASN:HD22	2.16	0.49
1:A:51:ALA:C	1:A:52:LEU:HD22	2.34	0.48
1:A:204:TRP:C	1:A:204:TRP:CE3	2.87	0.48
1:A:31:GLY:HA2	1:A:522:TYR:OH	2.14	0.47
1:A:464:SER:CB	1:A:481:THR:HG22	2.38	0.47
1:A:454:LEU:HD13	1:A:501:TRP:CG	2.50	0.47
1:A:161:THR:HG22	1:A:171:GLY:N	2.29	0.46
1:A:135:TYR:CD1	1:A:316:LYS:HD2	2.50	0.46
1:A:50:GLN:HG3	1:A:330:TYR:HB3	1.97	0.46
1:A:512:ILE:CG1	1:A:513:ALA:H	2.07	0.46
1:A:204:TRP:CG	1:A:205:PRO:HD3	2.51	0.46
1:A:467:ILE:O	1:A:467:ILE:HG23	2.17	0.45
1:A:241:GLU:O	1:A:242:GLU:O	2.34	0.45
1:A:331:PHE:CE2	1:A:334:ASP:HA	2.52	0.45
1:A:444:ASP:OD1	1:A:445:GLY:N	2.48	0.45
1:A:117:ASN:ND2	1:A:120:SER:H	2.13	0.45
1:A:205:PRO:HA	1:A:206:PRO:HA	1.75	0.45
1:A:288:GLN:HB3	1:A:291:ASN:O	2.16	0.44
1:A:291:ASN:ND2	1:A:293:MET:H	2.15	0.44
1:A:204:TRP:CB	1:A:205:PRO:CD	2.95	0.44
1:A:83:PHE:CD1	1:A:129:MET:HG3	2.52	0.43
1:A:335:GLU:O	1:A:336:VAL:CB	2.66	0.43
1:A:438:GLU:O	1:A:440:ILE:HG23	2.18	0.43
1:A:448:GLN:HA	1:A:448:GLN:OE1	2.18	0.43
1:A:161:THR:HG22	1:A:171:GLY:C	2.38	0.43
1:A:167:THR:O	1:A:168:LEU:HB2	2.16	0.43
1:A:195:ARG:HH11	1:A:195:ARG:HG2	1.83	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:ILE:HG21	1:A:312:PHE:CD2	2.54	0.43
1:A:354:TYR:CD2	1:A:374:GLY:HA2	2.53	0.43
1:A:358:TYR:CD2	1:A:368:CYS:HB3	2.54	0.43
1:A:291:ASN:ND2	1:A:293:MET:HB2	2.30	0.42
1:A:427:THR:OG1	1:A:428:THR:N	2.52	0.42
1:A:291:ASN:C	1:A:291:ASN:ND2	2.72	0.42
1:A:422:GLY:O	1:A:436:GLN:HG2	2.19	0.42
1:A:476:VAL:HG22	1:A:493:PHE:CD2	2.54	0.42
1:A:431:GLU:HB2	1:A:451:LEU:HD11	2.01	0.42
1:A:78:ASN:HD22	1:A:78:ASN:H	1.67	0.41
1:A:126:ASN:HB2	4:A:761:HOH:O	2.21	0.41
1:A:203:ASN:ND2	1:A:207:LEU:HD12	2.34	0.41
1:A:92:ASN:C	1:A:92:ASN:ND2	2.70	0.41
1:A:418:PHE:CG	1:A:419:PRO:HD2	2.55	0.41
1:A:468:GLY:HA3	1:A:475:TYR:CE2	2.56	0.41
1:A:117:ASN:HD22	1:A:120:SER:N	2.17	0.41
1:A:204:TRP:HB3	1:A:205:PRO:CD	2.46	0.41
1:A:313:LEU:N	1:A:313:LEU:CD1	2.84	0.41
1:A:57:SER:HB3	1:A:323:PRO:HB2	2.02	0.40
1:A:431:GLU:HA	1:A:452:PHE:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	529/555 (95%)	493 (93%)	25 (5%)	11 (2%)	<b>7</b> <b>4</b>

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	242	GLU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	243	ILE
1	A	244	PRO
1	A	272	PRO
1	A	275	SER
1	A	276	PRO
1	A	279	TYR
1	A	336	VAL
1	A	479	ALA
1	A	204	TRP
1	A	245	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	407/451 (90%)	394 (97%)	13 (3%)	39 50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LEU
1	A	78	ASN
1	A	92	ASN
1	A	102	ARG
1	A	204	TRP
1	A	206	PRO
1	A	291	ASN
1	A	299	ARG
1	A	309	VAL
1	A	313	LEU
1	A	345	ASN
1	A	368	CYS
1	A	435	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	92	ASN
1	A	100	HIS
1	A	117	ASN
1	A	123	ASN
1	A	131	GLN
1	A	136	HIS
1	A	172	GLN
1	A	201	ASN
1	A	203	ASN
1	A	234	GLN
1	A	288	GLN
1	A	291	ASN
1	A	294	ASN
1	A	321	ASN
1	A	345	ASN
1	A	382	GLN
1	A	414	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACT	A	560	-	3,3,3	1.40	0	3,3,3	1.23	0
2	ACT	A	561	-	3,3,3	1.38	0	3,3,3	1.28	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	533/555 (96%)	-0.09	20 (3%) 40 38	21, 36, 72, 98	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	PRO	6.9
1	A	247	PHE	6.6
1	A	279	TYR	5.4
1	A	243	ILE	4.8
1	A	301	VAL	3.3
1	A	514	ASP	3.2
1	A	281	ALA	3.1
1	A	513	ALA	3.1
1	A	515	ASP	3.0
1	A	277	CYS	2.8
1	A	283	ILE	2.7
1	A	278	ALA	2.7
1	A	270	LEU	2.5
1	A	556	PRO	2.5
1	A	245	VAL	2.4
1	A	242	GLU	2.4
1	A	280	GLU	2.2
1	A	303	TYR	2.1
1	A	204	TRP	2.1
1	A	22	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ACT	A	560	4/4	0.90	0.19	56,56,56,57	0
2	ACT	A	561	4/4	0.93	0.09	69,70,70,70	0
3	CA	A	601	1/1	0.96	0.06	55,55,55,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.