



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 07:52 am GMT

PDB ID : 5N7J
Title : Crystal structure of Neisseria polysaccharea amylosucrase mutant efficient for the synthesis of controlled size maltooligosaccharides
Authors : Verges, A.; Tranier, S.
Deposited on : 2017-02-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	628	
2	B	2	
2	C	2	
2	D	2	
3	E	3	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amylosucrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	628	5134	3260	890	963	21	0	14	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q9ZEU2
A	2	PRO	-	expression tag	UNP Q9ZEU2
A	3	ASN	-	expression tag	UNP Q9ZEU2
A	4	SER	-	expression tag	UNP Q9ZEU2
A	226	LYS	ARG	engineered mutation	UNP Q9ZEU2
A	228	VAL	ILE	engineered mutation	UNP Q9ZEU2
A	289	ILE	ALA	engineered mutation	UNP Q9ZEU2
A	290	TYR	PHE	engineered mutation	UNP Q9ZEU2
A	300	ILE	GLU	engineered mutation	UNP Q9ZEU2
A	331	THR	VAL	engineered mutation	UNP Q9ZEU2
A	437	SER	GLN	engineered mutation	UNP Q9ZEU2
A	439	ASP	ASN	engineered mutation	UNP Q9ZEU2
A	445	ALA	CYS	engineered mutation	UNP Q9ZEU2
A	537	ASP	GLY	engineered mutation	UNP Q9ZEU2

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0
2	C	2	23	12	11	0	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	E	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C O	0	0
			16	10 6		
5	A	1	Total	C O	0	0
			16	10 6		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C O	0	0
			13	8 5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	941	Total 941	O 941	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amylosucrase




- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose




- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.99Å 97.57Å 115.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.78 – 2.00 48.78 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (48.78-2.00) 98.2 (48.78-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.00Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.142 , 0.190 0.144 , 0.190	Depositor DCC
R_{free} test set	2305 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtrriage
Anisotropy	0.578	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6231	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, GLC, PG4, 1PE, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/5304	0.50	0/7215

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5134	0	4897	20	0
2	B	23	0	19	2	0
2	C	23	0	19	5	0
2	D	23	0	19	4	0
3	E	34	0	30	0	0
4	A	8	0	12	0	0
5	A	32	0	44	2	0
6	A	13	0	18	0	0
7	A	941	0	0	12	0
All	All	6231	0	5058	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:2:FRU:C2	2:D:2:FRU:O5	1.64	1.24
2:C:2:FRU:O5	2:C:2:FRU:C2	1.63	1.21
1:A:560[B]:ASN:OD1	7:A:801:HOH:O	1.95	0.84
7:A:810:HOH:O	2:B:2:FRU:H3	1.83	0.78
7:A:802:HOH:O	2:C:2:FRU:O6	2.00	0.78
2:D:1:GLC:C1	2:D:2:FRU:O5	2.41	0.68
1:A:460:ASP:OD2	7:A:803:HOH:O	2.12	0.67
1:A:381:GLU:OE1	7:A:804:HOH:O	2.13	0.66
7:A:805:HOH:O	2:D:1:GLC:O2	2.13	0.66
2:C:1:GLC:C1	2:C:2:FRU:O5	2.48	0.62
1:A:439:ASP:CG	7:A:816:HOH:O	2.41	0.59
1:A:495[A]:ASP:O	1:A:498[A]:TRP:HD1	1.89	0.56
1:A:441:SER:HB3	7:A:816:HOH:O	2.06	0.55
2:D:2:FRU:O5	2:D:2:FRU:O2	2.22	0.53
1:A:180:VAL:HG22	1:A:282:ILE:HB	1.92	0.51
1:A:357:LEU:HB3	1:A:471:LEU:HD22	1.92	0.51
1:A:580:ASN:O	1:A:620:TYR:HA	2.11	0.50
1:A:100:GLY:HA2	1:A:129:TYR:HB3	1.95	0.48
1:A:442:THR:N	7:A:816:HOH:O	2.48	0.47
2:C:2:FRU:O5	2:C:2:FRU:O2	2.28	0.46
1:A:5:GLN:O	1:A:9:THR:HG23	2.17	0.44
1:A:210:TYR:HB2	1:A:247:TRP:HB2	2.00	0.44
7:A:1505:HOH:O	2:C:2:FRU:H61	2.17	0.43
1:A:539:ARG:HA	1:A:539:ARG:HD2	1.86	0.43
1:A:182:ASP:OD1	1:A:284:ARG:HD3	2.20	0.42
1:A:498[A]:TRP:CG	1:A:499:SER:N	2.89	0.41
1:A:67[B]:GLN:HG3	7:A:995:HOH:O	2.20	0.41
1:A:356:LEU:HD23	1:A:356:LEU:HA	1.93	0.41
5:A:706:1PE:H152	7:A:888:HOH:O	2.19	0.41
1:A:594:GLN:HB2	2:B:2:FRU:H62	2.02	0.41
5:A:707:1PE:H251	5:A:707:1PE:H242	1.90	0.41
1:A:61:LEU:HD12	1:A:61:LEU:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	640/628 (102%)	627 (98%)	13 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	547/533 (103%)	540 (99%)	7 (1%)	69 74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67[A]	GLN
1	A	67[B]	GLN
1	A	129	TYR
1	A	208	PHE
1	A	496[A]	ASP
1	A	496[B]	ASP
1	A	622	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	11,11,12	2.27	2 (18%)	15,15,17	1.92	3 (20%)
2	FRU	B	2	2	11,12,12	5.10	6 (54%)	10,18,18	3.24	1 (10%)
2	GLC	C	1	2	11,11,12	2.02	2 (18%)	15,15,17	1.05	1 (6%)
2	FRU	C	2	2	11,12,12	5.21	7 (63%)	10,18,18	2.83	2 (20%)
2	GLC	D	1	2	11,11,12	2.05	2 (18%)	15,15,17	0.93	1 (6%)
2	FRU	D	2	2	11,12,12	5.18	6 (54%)	10,18,18	3.38	1 (10%)
3	GLC	E	1	3	12,12,12	1.49	2 (16%)	17,17,17	1.03	1 (5%)
3	GLC	E	2	3	11,11,12	1.52	2 (18%)	15,15,17	0.86	0
3	GLC	E	3	3	11,11,12	1.70	3 (27%)	15,15,17	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/19/22	0/1/1/1
2	FRU	B	2	2	-	3/5/24/24	0/1/1/1
2	GLC	C	1	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	C	2	2	-	4/5/24/24	0/1/1/1
2	GLC	D	1	2	-	0/2/19/22	0/1/1/1
2	FRU	D	2	2	-	4/5/24/24	0/1/1/1
3	GLC	E	1	3	-	0/2/22/22	0/1/1/1
3	GLC	E	2	3	-	0/2/19/22	0/1/1/1
3	GLC	E	3	3	-	0/2/19/22	0/1/1/1

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	FRU	O5-C2	13.41	1.64	1.43
2	C	2	FRU	O5-C2	13.18	1.63	1.43
2	B	2	FRU	O5-C2	12.53	1.62	1.43
2	B	2	FRU	O2-C2	6.61	1.52	1.40
2	C	2	FRU	O2-C2	6.38	1.51	1.40
2	D	2	FRU	O2-C2	6.37	1.51	1.40
2	C	2	FRU	O5-C5	-5.88	1.30	1.43
2	B	2	FRU	O5-C5	-5.75	1.31	1.43
2	B	1	GLC	O5-C1	5.61	1.52	1.43
2	D	2	FRU	O5-C5	-5.37	1.32	1.43
2	B	2	FRU	O3-C3	5.17	1.53	1.42
2	C	1	GLC	O5-C1	4.90	1.51	1.43
2	D	2	FRU	O3-C3	4.87	1.52	1.42
2	C	2	FRU	O3-C3	4.81	1.52	1.42
2	D	1	GLC	O5-C1	4.67	1.51	1.43
2	B	1	GLC	O5-C5	4.00	1.51	1.43
2	D	1	GLC	O5-C5	3.95	1.51	1.43
2	C	2	FRU	O4-C4	-3.75	1.34	1.43
3	E	2	GLC	O5-C1	3.71	1.49	1.43
3	E	3	GLC	O5-C1	3.63	1.49	1.43
2	D	2	FRU	O4-C4	-3.61	1.34	1.43
2	B	2	FRU	O4-C4	-3.55	1.34	1.43
2	C	1	GLC	O5-C5	3.47	1.50	1.43
3	E	1	GLC	O5-C1	3.44	1.51	1.42
2	B	2	FRU	C4-C5	2.56	1.59	1.53
2	C	2	FRU	C4-C5	2.41	1.59	1.53
3	E	1	GLC	O4-C4	2.40	1.48	1.43
2	D	2	FRU	C4-C5	2.33	1.58	1.53
3	E	3	GLC	C2-C3	-2.25	1.49	1.52
3	E	2	GLC	C2-C3	-2.23	1.49	1.52
3	E	3	GLC	O5-C5	2.11	1.47	1.43
2	C	2	FRU	C1-C2	-2.05	1.49	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	FRU	O2-C2-O5	-10.34	89.54	109.50
2	B	2	FRU	O2-C2-O5	-9.94	90.31	109.50
2	C	2	FRU	O2-C2-O5	-8.51	93.07	109.50
2	B	1	GLC	C1-O5-C5	5.81	120.06	112.19
2	C	1	GLC	C1-C2-C3	3.17	113.56	109.67
3	E	1	GLC	O5-C5-C4	2.51	114.25	109.69
2	B	1	GLC	C1-C2-C3	2.48	112.71	109.67
2	C	2	FRU	O1-C1-C2	-2.39	106.78	111.86
2	B	1	GLC	O5-C5-C4	2.34	116.51	110.83
2	D	1	GLC	O5-C1-C2	-2.10	107.52	110.77

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	FRU	O1-C1-C2-C3
2	C	2	FRU	C4-C5-C6-O6
2	C	2	FRU	O5-C5-C6-O6
2	D	2	FRU	O1-C1-C2-O2
2	D	2	FRU	O1-C1-C2-O5
2	D	2	FRU	O5-C5-C6-O6
2	D	2	FRU	C4-C5-C6-O6
2	B	1	GLC	O5-C5-C6-O6
2	B	1	GLC	C4-C5-C6-O6
2	B	2	FRU	O5-C5-C6-O6
2	C	2	FRU	O1-C1-C2-C3
2	C	2	FRU	O1-C1-C2-O5
2	B	2	FRU	C4-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	GLC	2	0
2	C	1	GLC	1	0
2	D	2	FRU	3	0
2	C	2	FRU	5	0
2	B	2	FRU	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	1PE	A	707	-	15,15,15	0.77	0	14,14,14	0.31	0
5	1PE	A	706	-	15,15,15	0.79	0	14,14,14	0.30	0
4	TRS	A	701	-	7,7,7	0.57	0	9,9,9	0.84	0
6	PG4	A	708	-	12,12,12	0.77	0	11,11,11	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1PE	A	707	-	-	7/13/13/13	-
5	1PE	A	706	-	-	8/13/13/13	-
4	TRS	A	701	-	-	0/9/9/9	-
6	PG4	A	708	-	-	3/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	706	1PE	OH4-C13-C23-OH3
5	A	707	1PE	OH6-C15-C25-OH5
5	A	706	1PE	C15-C25-OH5-C14
5	A	707	1PE	OH4-C13-C23-OH3
5	A	706	1PE	OH6-C15-C25-OH5
5	A	706	1PE	C13-C23-OH3-C22
6	A	708	PG4	O3-C5-C6-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	706	1PE	OH5-C14-C24-OH4
5	A	707	1PE	C24-C14-OH5-C25
5	A	706	1PE	C14-C24-OH4-C13
5	A	706	1PE	OH7-C16-C26-OH6
6	A	708	PG4	O4-C7-C8-O5
6	A	708	PG4	C5-C6-O4-C7
5	A	707	1PE	C23-C13-OH4-C24
5	A	706	1PE	C23-C13-OH4-C24
5	A	707	1PE	OH5-C14-C24-OH4
5	A	707	1PE	OH2-C12-C22-OH3
5	A	707	1PE	C13-C23-OH3-C22

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	707	1PE	1	0
5	A	706	1PE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	628/628 (100%)	-0.71	1 (0%) 95 94	8, 15, 30, 55	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	441	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	1	11/12	0.75	0.29	28,38,45,49	0
2	FRU	B	2	12/12	0.80	0.24	22,45,51,51	0
3	GLC	E	1	12/12	0.80	0.29	32,54,65,69	0
2	FRU	C	2	12/12	0.81	0.34	31,46,54,61	0
2	FRU	D	2	12/12	0.83	0.31	14,47,50,51	0
2	GLC	C	1	11/12	0.84	0.24	28,43,50,52	0
3	GLC	E	3	11/12	0.86	0.18	24,38,49,50	0
2	GLC	B	1	11/12	0.90	0.14	20,30,41,43	0
3	GLC	E	2	11/12	0.92	0.13	21,33,42,44	0




The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	1PE	A	707	16/16	0.85	0.17	27,45,60,62	0
6	PG4	A	708	13/13	0.87	0.26	37,45,59,61	0
5	1PE	A	706	16/16	0.90	0.17	30,38,53,55	0
4	TRS	A	701	8/8	0.94	0.11	10,15,17,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.