



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 28, 2023 – 09:55 PM EDT

PDB ID : 3N9U
Title : Crystal Structure of the Complex between the 25 kDa Subunit and the 59 kDa Subunit (RRM domain) of Human Cleavage Factor Im
Authors : Tresaugues, L.; Welin, M.; Arrowsmith, C.H.; Berglund, H.; Bountra, C.; Collins, R.; Edwards, A.M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Johansson, I.; Karlberg, T.; Kol, S.; Kotenyova, T.; Moche, M.; Nielsen, T.N.; Nyman, T.; Persson, C.; Schuler, H.; Schutz, P.; Siponen, M.I.; Thorsell, A.G.; van der Berg, S.; Wahlberg, E.; Weigelt, J.; Wisniewska, M.; Nordlund, P.; Structural Genomics Consortium (SGC)
Deposited on : 2010-05-31
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 6% 83% 6% • 10%
1	B	230	 2% 82% 6% 13%
2	C	156	 3% 54% 8% 38%

Continued on next page...

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.35

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	156	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a red segment (37%), a green segment (44%), a yellow segment (12%), and a grey segment (44%). The percentages are labeled above and below the segments.</p>

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5214 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cleavage and polyadenylation specificity factor subunit 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	1695	1107	283	301	4	0	1	0
1	B	201	1647	1076	274	293	4	0	2	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP O43809
A	-1	HIS	-	expression tag	UNP O43809
A	0	HIS	-	expression tag	UNP O43809
A	1	HIS	-	expression tag	UNP O43809
A	2	HIS	-	expression tag	UNP O43809
A	3	HIS	-	expression tag	UNP O43809
A	4	HIS	-	expression tag	UNP O43809
A	5	SER	-	expression tag	UNP O43809
A	6	SER	-	expression tag	UNP O43809
A	7	GLY	-	expression tag	UNP O43809
A	8	VAL	-	expression tag	UNP O43809
A	9	ASP	-	expression tag	UNP O43809
A	10	LEU	-	expression tag	UNP O43809
A	11	GLY	-	expression tag	UNP O43809
A	12	THR	-	expression tag	UNP O43809
A	13	GLU	-	expression tag	UNP O43809
A	14	ASN	-	expression tag	UNP O43809
A	15	LEU	-	expression tag	UNP O43809
A	16	TYR	-	expression tag	UNP O43809
A	17	PHE	-	expression tag	UNP O43809
A	18	GLN	-	expression tag	UNP O43809
A	19	SER	-	expression tag	UNP O43809
A	20	MET	-	expression tag	UNP O43809
B	-2	MET	-	expression tag	UNP O43809
B	-1	HIS	-	expression tag	UNP O43809

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP O43809
B	1	HIS	-	expression tag	UNP O43809
B	2	HIS	-	expression tag	UNP O43809
B	3	HIS	-	expression tag	UNP O43809
B	4	HIS	-	expression tag	UNP O43809
B	5	SER	-	expression tag	UNP O43809
B	6	SER	-	expression tag	UNP O43809
B	7	GLY	-	expression tag	UNP O43809
B	8	VAL	-	expression tag	UNP O43809
B	9	ASP	-	expression tag	UNP O43809
B	10	LEU	-	expression tag	UNP O43809
B	11	GLY	-	expression tag	UNP O43809
B	12	THR	-	expression tag	UNP O43809
B	13	GLU	-	expression tag	UNP O43809
B	14	ASN	-	expression tag	UNP O43809
B	15	LEU	-	expression tag	UNP O43809
B	16	TYR	-	expression tag	UNP O43809
B	17	PHE	-	expression tag	UNP O43809
B	18	GLN	-	expression tag	UNP O43809
B	19	SER	-	expression tag	UNP O43809
B	20	MET	-	expression tag	UNP O43809

- Molecule 2 is a protein called Cleavage and polyadenylation specificity factor subunit 7.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	C	96	Total	C	N	O	S	0	1	0
			749	477	130	141	1			
2	I	87	Total	C	N	O		0	0	0
			688	438	123	127				

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	27	MET	-	expression tag	UNP Q8N684
C	28	HIS	-	expression tag	UNP Q8N684
C	29	HIS	-	expression tag	UNP Q8N684
C	30	HIS	-	expression tag	UNP Q8N684
C	31	HIS	-	expression tag	UNP Q8N684
C	32	HIS	-	expression tag	UNP Q8N684
C	33	HIS	-	expression tag	UNP Q8N684
C	34	SER	-	expression tag	UNP Q8N684
C	35	SER	-	expression tag	UNP Q8N684

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	36	GLY	-	expression tag	UNP Q8N684
C	37	VAL	-	expression tag	UNP Q8N684
C	38	ASP	-	expression tag	UNP Q8N684
C	39	LEU	-	expression tag	UNP Q8N684
C	40	GLY	-	expression tag	UNP Q8N684
C	41	THR	-	expression tag	UNP Q8N684
C	42	GLU	-	expression tag	UNP Q8N684
C	43	ASN	-	expression tag	UNP Q8N684
C	44	LEU	-	expression tag	UNP Q8N684
C	45	TYR	-	expression tag	UNP Q8N684
C	46	PHE	-	expression tag	UNP Q8N684
C	47	GLN	-	expression tag	UNP Q8N684
C	48	SER	-	expression tag	UNP Q8N684
C	49	MET	-	expression tag	UNP Q8N684
I	27	MET	-	expression tag	UNP Q8N684
I	28	HIS	-	expression tag	UNP Q8N684
I	29	HIS	-	expression tag	UNP Q8N684
I	30	HIS	-	expression tag	UNP Q8N684
I	31	HIS	-	expression tag	UNP Q8N684
I	32	HIS	-	expression tag	UNP Q8N684
I	33	HIS	-	expression tag	UNP Q8N684
I	34	SER	-	expression tag	UNP Q8N684
I	35	SER	-	expression tag	UNP Q8N684
I	36	GLY	-	expression tag	UNP Q8N684
I	37	VAL	-	expression tag	UNP Q8N684
I	38	ASP	-	expression tag	UNP Q8N684
I	39	LEU	-	expression tag	UNP Q8N684
I	40	GLY	-	expression tag	UNP Q8N684
I	41	THR	-	expression tag	UNP Q8N684
I	42	GLU	-	expression tag	UNP Q8N684
I	43	ASN	-	expression tag	UNP Q8N684
I	44	LEU	-	expression tag	UNP Q8N684
I	45	TYR	-	expression tag	UNP Q8N684
I	46	PHE	-	expression tag	UNP Q8N684
I	47	GLN	-	expression tag	UNP Q8N684
I	48	SER	-	expression tag	UNP Q8N684
I	49	MET	-	expression tag	UNP Q8N684

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	133	Total O 133 133	0	0
4	B	194	Total O 194 194	0	0
4	C	70	Total O 70 70	0	0
4	I	20	Total O 20 20	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cleavage and polyadenylation specificity factor subunit 5




- Molecule 1: Cleavage and polyadenylation specificity factor subunit 5




- Molecule 2: Cleavage and polyadenylation specificity factor subunit 7



- Molecule 2: Cleavage and polyadenylation specificity factor subunit 7





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	165.95Å 124.73Å 40.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.50 – 1.92 41.38 – 1.92	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.50-1.92) 97.6 (41.38-1.92)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.196 , 0.231 0.197 , 0.228	Depositor DCC
R_{free} test set	3216 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.505	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5214	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	0/1746	0.71	2/2371 (0.1%)
1	B	0.68	0/1700	0.74	2/2310 (0.1%)
2	C	0.71	0/764	0.73	0/1034
2	I	0.40	0/698	0.51	0/943
All	All	0.66	0/4908	0.70	4/6658 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH1	6.69	123.64	120.30
1	A	125	MET	CG-SD-CE	-6.18	90.31	100.20
1	B	78	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	78	ARG	NE-CZ-NH1	5.09	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1695	0	1703	15	0
1	B	1647	0	1650	9	0
2	C	749	0	749	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	I	688	0	695	14	0
3	A	6	0	8	1	0
3	B	12	0	16	3	0
4	A	133	0	0	1	1
4	B	194	0	0	0	0
4	C	70	0	0	1	1
4	I	20	0	0	2	0
All	All	5214	0	4821	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:MET:CE	1:A:177:VAL:HG21	2.10	0.82
1:A:125:MET:HE1	1:A:177:VAL:HG21	1.63	0.81
2:C:82:ALA:HB1	2:C:138:VAL:HG21	1.61	0.80
2:I:95:ASP:OD2	4:I:509:HOH:O	2.11	0.68
1:B:211:ILE:HD12	3:B:228:GOL:H31	1.76	0.68
1:A:99:LEU:HD22	1:A:191:TYR:CE1	2.31	0.66
2:I:101:VAL:HG21	2:I:150:LEU:HD13	1.77	0.66
2:I:158:ARG:HB3	2:I:164:ASN:HD22	1.61	0.65
2:I:85:TYR:CZ	2:I:165:LEU:HD13	2.34	0.62
1:A:83:VAL:HG12	1:A:85[B]:ILE:HD11	1.87	0.57
2:C:99:ILE:HG12	2:C:113:LEU:HD11	1.88	0.55
2:I:145:LEU:HB3	2:I:157:VAL:HG21	1.88	0.55
1:A:125:MET:HE2	1:A:177:VAL:HG21	1.86	0.54
2:I:111:VAL:HG23	2:I:132:VAL:C	2.28	0.54
1:A:99:LEU:HD22	1:A:191:TYR:CD1	2.43	0.53
1:A:145:ILE:HG22	1:A:218:LEU:HD13	1.91	0.53
2:I:113:LEU:CD2	2:I:130:VAL:HG22	2.40	0.51
1:A:85[B]:ILE:CD1	1:A:94:VAL:HG22	2.40	0.51
2:I:113:LEU:HD21	2:I:130:VAL:HG22	1.95	0.49
1:A:137:GLN:HG2	1:A:139:TRP:CZ2	2.48	0.48
2:I:158:ARG:HB3	2:I:164:ASN:ND2	2.27	0.48
2:I:114:LYS:NZ	4:I:245:HOH:O	2.46	0.48
2:C:111[B]:VAL:CG1	2:C:131:VAL:HG12	2.46	0.46
1:B:101:THR:HB	1:B:102:THR:HG23	1.98	0.46
1:B:121:LEU:O	1:B:125:MET:HG2	2.16	0.46
2:C:111[B]:VAL:HG13	2:C:131:VAL:HG12	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:102:ILE:HG23	2:I:141:LEU:HD21	1.97	0.46
1:B:156:PRO:HA	3:B:229:GOL:H12	1.97	0.45
1:A:105:LYS:HA	3:A:228:GOL:H12	1.99	0.44
1:B:105:LYS:HA	3:B:228:GOL:H11	1.97	0.44
2:C:175:ARG:NE	4:C:3:HOH:O	2.44	0.43
1:A:105:LYS:NZ	4:A:296:HOH:O	2.51	0.43
2:I:85:TYR:CE1	2:I:165:LEU:HD13	2.52	0.43
2:C:145:LEU:HB3	2:C:157:VAL:HG21	2.00	0.42
2:C:177:CYS:SG	2:C:177:CYS:O	2.77	0.42
2:I:107:VAL:HG21	2:I:141:LEU:HG	2.02	0.42
1:A:83:VAL:CG1	1:A:85[B]:ILE:HD11	2.49	0.41
1:A:145:ILE:CG2	1:A:218:LEU:HD13	2.50	0.41
2:C:93:THR:HG21	2:C:150:LEU:HD13	2.03	0.41
1:A:35:ARG:HD3	1:B:32:THR:HG22	2.03	0.41
1:B:89:HIS:HB3	2:I:92:TRP:HB3	2.03	0.40
1:B:85:ILE:HD12	1:B:176:LEU:HD11	2.03	0.40
1:A:145:ILE:HG23	1:A:218:LEU:HB3	2.03	0.40
1:B:193:LEU:HD23	1:B:193:LEU:C	2.42	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:338:HOH:O	4:C:200:HOH:O[3_658]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/230 (89%)	200 (98%)	5 (2%)	0	100	100
1	B	201/230 (87%)	197 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	95/156 (61%)	94 (99%)	1 (1%)	0	100	100
2	I	83/156 (53%)	83 (100%)	0	0	100	100
All	All	584/772 (76%)	574 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/206 (89%)	181 (98%)	3 (2%)	62	58
1	B	178/206 (86%)	178 (100%)	0	100	100
2	C	78/137 (57%)	78 (100%)	0	100	100
2	I	73/137 (53%)	73 (100%)	0	100	100
All	All	513/686 (75%)	510 (99%)	3 (1%)	86	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	57	ASP
1	A	137	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	228	-	5,5,5	0.42	0	5,5,5	0.86	0
3	GOL	B	229	-	5,5,5	0.19	0	5,5,5	0.99	0
3	GOL	A	228	-	5,5,5	0.36	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	228	-	-	2/4/4/4	-
3	GOL	B	229	-	-	2/4/4/4	-
3	GOL	A	228	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	228	GOL	O1-C1-C2-C3
3	B	228	GOL	O2-C2-C3-O3
3	A	228	GOL	C1-C2-C3-O3
3	B	228	GOL	C1-C2-C3-O3
3	B	229	GOL	O1-C1-C2-C3
3	A	228	GOL	O1-C1-C2-O2
3	A	228	GOL	O2-C2-C3-O3
3	B	229	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	228	GOL	2	0
3	B	229	GOL	1	0
3	A	228	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/230 (89%)	0.40	13 (6%) 20 22	13, 25, 48, 55	0
1	B	201/230 (87%)	0.12	5 (2%) 57 60	13, 22, 42, 48	0
2	C	96/156 (61%)	0.14	5 (5%) 27 30	16, 23, 40, 65	0
2	I	87/156 (55%)	3.46	57 (65%) 0 0	30, 80, 96, 99	0
All	All	590/772 (76%)	0.72	80 (13%) 3 3	13, 25, 83, 99	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	105	ILE	9.4
2	I	141	LEU	8.5
2	I	139	HIS	8.1
2	I	132	VAL	8.0
2	I	144	LEU	7.9
2	I	110	VAL	7.9
2	I	142	LEU	7.4
2	I	149	VAL	7.4
2	I	107	VAL	7.4
2	I	104	SER	7.3
2	I	101	VAL	7.3
2	I	102	ILE	7.2
2	I	108	TYR	7.1
2	I	138	VAL	7.0
2	I	145	LEU	6.6
2	I	84	VAL	6.5
2	I	130	VAL	6.4
2	I	159	PRO	5.8
2	I	161	THR	5.7
1	A	136	LEU	5.7
2	I	140	LYS	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	111	VAL	4.9
2	I	113	LEU	4.8
2	I	157	VAL	4.7
2	I	109	ASP	4.7
2	I	150	LEU	4.5
2	I	103	ARG	4.5
2	I	143	GLU	4.4
2	I	112	GLU	4.3
2	I	147	GLY	4.3
2	I	156	ASP	4.3
2	I	175	ARG	4.2
2	I	167	GLN	4.2
2	I	162	ARG	4.2
2	I	158	ARG	4.1
2	C	177	CYS	4.1
2	I	98	LEU	4.0
2	I	160	ALA	4.0
2	I	100	GLN	3.9
1	A	134	GLY	3.9
2	I	173	ARG	3.9
2	I	163	GLN	3.9
2	I	131	VAL	3.9
2	I	87	GLY	3.8
2	I	154	LYS	3.8
2	I	155	VAL	3.6
2	I	148	LYS	3.6
2	I	152	GLY	3.5
1	A	135	VAL	3.5
2	I	85	TYR	3.4
1	B	28	THR	3.4
2	I	146	PRO	3.2
2	I	86	VAL	3.2
2	I	164	ASN	3.1
2	I	106	GLY	3.0
1	B	101	THR	2.9
2	C	176	GLU	2.9
1	A	133	ASP	2.9
1	A	74	ILE	2.9
1	A	132	GLN	2.8
2	I	153	GLU	2.8
2	I	99	ILE	2.8
1	A	101	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	129	GLU	2.6
1	A	33	LEU	2.5
1	B	134	GLY	2.5
1	B	132	GLN	2.5
2	I	96	GLN	2.4
1	A	58	SER	2.4
2	I	128	ALA	2.3
1	B	27	GLN	2.3
1	A	60	VAL	2.1
1	A	73	LYS	2.1
2	I	166	SER	2.1
1	A	24	TYR	2.1
2	I	168	PHE	2.1
2	C	147	GLY	2.1
2	C	149	VAL	2.0
1	A	65	GLN	2.0
2	C	152	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	229	6/6	0.81	0.17	37,41,43,46	0
3	GOL	B	228	6/6	0.92	0.21	31,39,41,42	0
3	GOL	A	228	6/6	0.93	0.15	31,39,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.