

Full wwPDB NMR Structure Validation Report (i)

Jun 4, 2023 – 09:42 AM EDT

PDB ID : 2NCO BMRB ID : 26723

Title : NMR solution structure for the C-terminal domain of Tetrahymena Tcb2 in

the absence of calcium

Authors: Fowler, A.; Kilpatrick, A.M.

Deposited on : 2016-04-11

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	102	62%	19%	•	18%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model						
1	A:120-A:179, A:191-A:214	0.51	5			
	(84)					

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 5, 7, 11, 14
2	8, 17, 18, 20
3	10, 12, 16, 19
4	1, 3, 15
5	6, 9
6	4, 13



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1619 atoms, of which 808 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called 25 kDa calcium-binding protein.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	100	Total	С	Н	N	О	S	0
	1 A 102	1619	509	808	134	165	3	0	




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: 25 kDa calcium-binding protein




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: 25 kDa calcium-binding protein




4.2.2 Score per residue for model 2






4.2.3 Score per residue for model 3

• Molecule 1: 25 kDa calcium-binding protein




4.2.4 Score per residue for model 4

• Molecule 1: 25 kDa calcium-binding protein




4.2.5 Score per residue for model 5 (medoid)

• Molecule 1: 25 kDa calcium-binding protein




4.2.6 Score per residue for model 6






4.2.7 Score per residue for model 7

• Molecule 1: 25 kDa calcium-binding protein




4.2.8 Score per residue for model 8

• Molecule 1: 25 kDa calcium-binding protein




4.2.9 Score per residue for model 9

• Molecule 1: 25 kDa calcium-binding protein




4.2.10 Score per residue for model 10






4.2.11 Score per residue for model 11

• Molecule 1: 25 kDa calcium-binding protein




4.2.12 Score per residue for model 12

• Molecule 1: 25 kDa calcium-binding protein




4.2.13 Score per residue for model 13

• Molecule 1: 25 kDa calcium-binding protein




4.2.14 Score per residue for model 14






4.2.15 Score per residue for model 15

• Molecule 1: 25 kDa calcium-binding protein




4.2.16 Score per residue for model 16

• Molecule 1: 25 kDa calcium-binding protein




4.2.17 Score per residue for model 17

• Molecule 1: 25 kDa calcium-binding protein




4.2.18 Score per residue for model 18






4.2.19 Score per residue for model 19

• Molecule 1: 25 kDa calcium-binding protein



4.2.20 Score per residue for model 20





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1228
Number of shifts mapped to atoms	1228
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	681	684	684	13±3
All	All	13620	13680	13680	253

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$\operatorname{Distance}(\mathring{\mathrm{A}})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:130:LEU:HD11	1:A:201:ILE:HD11	0.88	1.45	3	5
1:A:130:LEU:HD21	1:A:201:ILE:HD11	0.82	1.51	14	12
1:A:133:ALA:HB1	1:A:196:TYR:CE2	0.67	2.24	20	5
1:A:137:PHE:CE2	1:A:148:LEU:HD12	0.67	2.25	15	11
1:A:137:PHE:CD2	1:A:148:LEU:HD12	0.65	2.26	2	9
1:A:172:THR:O	1:A:176:VAL:HG23	0.65	1.92	17	20
1:A:204:LEU:HD22	1:A:209:ILE:HD13	0.65	1.67	12	9
1:A:207:ALA:HB3	1:A:209:ILE:HD12	0.64	1.69	16	6
1:A:140:TYR:HB3	1:A:148:LEU:HD21	0.61	1.72	16	1
1:A:166:MET:CE	1:A:209:ILE:HD11	0.61	2.26	4	3
1:A:128:ALA:O	1:A:132:VAL:HG23	0.60	1.97	3	14
1:A:204:LEU:HG	1:A:209:ILE:HD13	0.60	1.74	2	5
1:A:196:TYR:CE1	1:A:199:LEU:HD22	0.60	2.32	19	3
1:A:136:LEU:HD21	1:A:159:ASP:HB3	0.59	1.74	8	4

Continued on next page...



 $Continued\ from\ previous\ page...$

Continuea from previ		Cladb(Å)	Distance (Å)	Models	
Atom-1	Atom-2	Clash(Å)	$\operatorname{Distance}(\mathrm{\AA})$	Worst	Total
1:A:157:LEU:HD13	1:A:171:PRO:CG	0.59	2.27	17	12
1:A:130:LEU:HD21	1:A:201:ILE:CD1	0.59	2.27	11	6
1:A:134:ARG:CG	1:A:193:LEU:HD21	0.58	2.26	19	1
1:A:148:LEU:CD2	1:A:156:LEU:HD22	0.58	2.29	3	1
1:A:136:LEU:HD22	1:A:156:LEU:CD1	0.58	2.29	12	2
1:A:134:ARG:HG3	1:A:193:LEU:HD21	0.57	1.74	19	4
1:A:123:ASN:OD1	1:A:126:VAL:HG23	0.57	2.00	20	4
1:A:130:LEU:CD1	1:A:201:ILE:HD11	0.56	2.27	3	4
1:A:204:LEU:O	1:A:209:ILE:HD12	0.56	2.00	4	2
1:A:196:TYR:CE1	1:A:200:ILE:HD11	0.56	2.35	8	1
1:A:137:PHE:HA	1:A:156:LEU:HD11	0.56	1.77	17	1
1:A:126:VAL:HG13	1:A:204:LEU:CD2	0.56	2.30	3	1
1:A:207:ALA:HB3	1:A:209:ILE:CD1	0.56	2.31	16	5
1:A:191:VAL:HG23	1:A:195:GLU:OE2	0.56	2.00	1	2
1:A:126:VAL:HG13	1:A:204:LEU:HD23	0.55	1.79	3	1
1:A:205:GLN:NE2	1:A:211:VAL:HG12	0.55	2.17	19	4
1:A:153:ILE:HD11	1:A:191:VAL:HG11	0.54	1.78	13	2
1:A:157:LEU:HD13	1:A:171:PRO:HG3	0.53	1.79	1	13
1:A:204:LEU:CD2	1:A:209:ILE:HD13	0.53	2.34	13	2
1:A:196:TYR:CD1	1:A:199:LEU:HD22	0.52	2.39	19	1
1:A:148:LEU:HD13	1:A:156:LEU:HD12	0.52	1.80	2	2
1:A:148:LEU:HD23	1:A:153:ILE:HD13	0.52	1.80	17	1
1:A:191:VAL:HG23	1:A:195:GLU:OE1	0.51	2.06	10	1
1:A:134:ARG:HA	1:A:193:LEU:HD11	0.51	1.83	8	2
1:A:133:ALA:HB2	1:A:200:ILE:HG13	0.51	1.82	18	1
1:A:129:LYS:HB3	1:A:204:LEU:HD11	0.51	1.82	18	1
1:A:191:VAL:HG23	1:A:195:GLU:CD	0.50	2.26	18	1
1:A:199:LEU:O	1:A:199:LEU:HD13	0.50	2.07	13	2
1:A:126:VAL:HG21	1:A:211:VAL:HG22	0.50	1.84	8	3
1:A:140:TYR:CD1	1:A:156:LEU:HD13	0.49	2.42	8	3
1:A:148:LEU:HD23	1:A:156:LEU:HD22	0.49	1.85	3	1
1:A:136:LEU:HD21	1:A:159:ASP:CB	0.49	2.37	14	3
1:A:136:LEU:HD21	1:A:159:ASP:HB2	0.49	1.85	17	1
1:A:160:THR:HG21	1:A:200:ILE:HD11	0.48	1.85	6	1
1:A:140:TYR:CB	1:A:148:LEU:HD21	0.48	2.38	16	1
1:A:205:GLN:NE2	1:A:211:VAL:H	0.48	2.07	13	1
1:A:148:LEU:HD13	1:A:156:LEU:CD1	0.47	2.39	2	2
1:A:123:ASN:HD21	1:A:126:VAL:HG23	0.47	1.69	9	5
1:A:136:LEU:HD23	1:A:160:THR:OG1	0.47	2.09	6	4
1:A:148:LEU:O	1:A:191:VAL:HG13	0.47	2.10	17	1
1:A:148:LEU:HD13	1:A:156:LEU:HD22	0.47	1.87	11	1

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Clash(Å)	$\operatorname{Distance}(\mathring{\mathrm{A}})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:129:LYS:HG3	1:A:204:LEU:HD21	0.46	1.88	17	1
1:A:170:THR:HG22	1:A:171:PRO:HD2	0.46	1.87	3	17
1:A:157:LEU:HD13	1:A:171:PRO:CB	0.46	2.41	7	3
1:A:153:ILE:HD11	1:A:191:VAL:CG1	0.46	2.40	15	1
1:A:127:GLU:HA	1:A:130:LEU:HD12	0.45	1.87	6	2
1:A:166:MET:SD	1:A:209:ILE:HD11	0.45	2.52	20	4
1:A:179:TRP:CE3	1:A:199:LEU:HD21	0.44	2.48	4	1
1:A:140:TYR:CG	1:A:156:LEU:HD12	0.44	2.48	20	1
1:A:148:LEU:CD2	1:A:156:LEU:HD12	0.43	2.43	17	1
1:A:134:ARG:HG2	1:A:193:LEU:HD21	0.43	1.89	6	1
1:A:148:LEU:HD13	1:A:152:GLU:OE2	0.43	2.13	15	1
1:A:136:LEU:HD12	1:A:140:TYR:CE1	0.43	2.49	18	1
1:A:156:LEU:HD21	1:A:196:TYR:CE2	0.42	2.49	7	1
1:A:156:LEU:HD23	1:A:156:LEU:O	0.42	2.15	2	1
1:A:179:TRP:CB	1:A:199:LEU:HD12	0.41	2.46	16	1
1:A:132:VAL:HG13	1:A:163:GLU:HB3	0.40	1.92	14	1
1:A:204:LEU:HD22	1:A:209:ILE:CD1	0.40	2.46	18	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	84/102 (82%)	80±1 (95±2%)	4±2 (5±2%)	0±0 (0±1%)	38	78
All	All	1680/2040 (82%)	1594 (95%)	80 (5%)	6 (0%)	38	78

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	141	ASP	5
1	A	209	ILE	1



6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric C		Outliers	Percentiles		
1	A	74/90 (82%)	65±2 (88±3%)	9±2 (12±3%)	9	52	
All	All	1480/1800 (82%)	1308 (88%)	172 (12%)	9	52	

All 37 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	170	THR	20
1	A	172	THR	18
1	A	121	LYS	17
1	A	206	LYS	11
1	A	123	ASN	11
1	A	175	ASP	7
1	A	213	LYS	7
1	A	167	SER	6
1	A	136	LEU	6
1	A	152	GLU	5
1	A	177	LYS	5
1	A	131	ASP	5
1	A	142	LYS	5
1	A	214	GLN	4
1	A	173	LYS	4
1	A	163	GLU	4
1	A	138	LYS	3
1	A	134	ARG	3
1	A	198	ASP	3
1	A	199	LEU	3
1	A	212	GLU	2
1	A	143	ASP	2
1	A	158	LYS	2
1	A	210	ARG	2
1	A	135	ARG	2
1	A	203	SER	2
1	A	202	LYS	2
1	A	148	LEU	2
1	A	125	GLU	1

Continued on next page..



Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	139	ARG	1
1	A	151	ASP	1
1	A	195	GLU	1
1	A	156	LEU	1
1	A	149	GLN	1
1	A	174	GLU	1
1	A	192	SER	1
1	A	194	GLU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1228
Number of shifts mapped to atoms	1228
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction} \pm {\rm precision}, ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	100	-0.32 ± 0.08	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	95	0.31 ± 0.05	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	91	-0.35 ± 0.08	None needed ($< 0.5 \text{ ppm}$)
^{15}N	92	-0.13 ± 0.25	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1051 atoms were assigned a chemical shift out of a possible 1175. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$		
Backbone	412/419 (98%)	168/170 (99%)	164/168 (98%)	80/81 (99%)		
Sidechain	575/688 (84%)	377/441 (85%)	192/219 (88%)	6/28 (21%)		

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	64/68 (94%)	32/32 (100%)	31/35 (89%)	1/1 (100%)
Overall	1051/1175 (89%)	577/643 (90%)	387/422 (92%)	87/110 (79%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1228 atoms were assigned a chemical shift out of a possible 1388. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	481/510 (94%)	198/207 (96%)	191/204 (94%)	92/99 (93%)
Sidechain	673/800 (84%)	442/514 (86%)	223/255 (87%)	8/31 (26%)
Aromatic	74/78 (95%)	37/37 (100%)	36/40~(90%)	1/1 (100%)
Overall	1228/1388 (88%)	677/758 (89%)	450/499 (90%)	101/131 (77%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1740
Intra-residue ($ i-j =0$)	416
Sequential ($ i-j =1$)	347
Medium range ($ i-j >1$ and $ i-j <5$)	401
Long range (i-j ≥5)	576
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	17.1
Number of long range restraints per residue ¹	5.6

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.3	0.19
0.2-0.5 (Medium)	0.1	0.26
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dantuninta tema	C	% ¹	Vio	lated	3	Consis	tentl	$\overline{ m y~Violated^4}$
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	416	23.9	4	1.0	0.2	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	342	19.7	4	1.2	0.2	0	0.0	0.0
Sidechain-Sidechain	74	4.3	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	347	19.9	1	0.3	0.1	0	0.0	0.0
Backbone-Backbone	79	4.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	199	11.4	1	0.5	0.1	0	0.0	0.0
Sidechain-Sidechain	69	4.0	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	401	23.0	1	0.2	0.1	0	0.0	0.0
Backbone-Backbone	60	3.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	185	10.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	156	9.0	1	0.6	0.1	0	0.0	0.0
Long range ($ i-j \ge 5$)	576	33.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	7	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	178	10.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	391	22.5	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1740	100.0	6	0.3	0.3	0	0.0	0.0
Backbone-Backbone	146	8.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	904	52.0	5	0.6	0.3	0	0.0	0.0
Sidechain-Sidechain	690	39.7	1	0.1	0.1	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID	Number of violations						Magn (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (A)	$SD^*(A)$	Median (A)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	1	0	0	0	0	1	0.15	0.15	0.0	0.15
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0
11	1	0	0	0	0	1	0.19	0.19	0.0	0.19

Continued on next page...




Continued from previous page...

Madal ID	Number of violations						Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	0	1	0	0	0	1	0.11	0.11	0.0	0.11
13	1	0	0	0	0	1	0.17	0.17	0.0	0.17
14	1	0	0	0	0	1	0.26	0.26	0.0	0.26
15	1	0	0	0	0	1	0.14	0.14	0.0	0.14
16	1	0	0	0	0	1	0.17	0.17	0.0	0.17
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	1	0	0	1	0.12	0.12	0.0	0.12
19	0	0	0	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0	0	0	0.0	0.0	0.0	0.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1734(IR:412, SQ:346, MR:400, LR:576, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
2	1	1	0	0	4	1	5.0
2	0	0	0	0	2	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	${f Models}^1$	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG2	2	0.22	0.04	0.22
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG3	2	0.22	0.04	0.22
(1,1009)	1:A:174:GLU:H	1:A:174:GLU:HB2	2	0.17	0.0	0.17


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table : All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG2	14	0.26
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG3	14	0.26
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG2	11	0.19
(1,1723)	1:A:212:GLU:HA	1:A:212:GLU:HG3	11	0.19
(1,1009)	1:A:174:GLU:H	1:A:174:GLU:HB2	13	0.17
(1,1009)	1:A:174:GLU:H	1:A:174:GLU:HB2	16	0.17
(1,1093)	1:A:119:LYS:HA	1:A:119:LYS:HD2	5	0.15
(1,1093)	1:A:119:LYS:HA	1:A:119:LYS:HD3	5	0.15
(1,1152)	1:A:125:GLU:HA	1:A:125:GLU:HG2	15	0.14
(1,1152)	1:A:125:GLU:HA	1:A:125:GLU:HG3	15	0.14
(1,217)	1:A:136:LEU:HD21	1:A:140:TYR:HE1	18	0.12
(1,217)	1:A:136:LEU:HD21	1:A:140:TYR:HE2	18	0.12
(1,217)	1:A:136:LEU:HD22	1:A:140:TYR:HE1	18	0.12
(1,217)	1:A:136:LEU:HD22	1:A:140:TYR:HE2	18	0.12
(1,217)	1:A:136:LEU:HD23	1:A:140:TYR:HE1	18	0.12
(1,217)	1:A:136:LEU:HD23	1:A:140:TYR:HE2	18	0.12

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)	
(1,1250)	1:A:134:ARG:HB2	1:A:135:ARG:H	12	0.11	
(1,1250)	1:A:134:ARG:HB3	1:A:135:ARG:H	12	0.11	



10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

