

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 04:23 AM EDT

PDB ID	:	6NES
Title	:	FAD-dependent monooxygenase TropB from T. stipitatus
Authors	:	Rodriguez Benitez, A.; Tweedy, S.E.; Baker Dockrey, S.A.; Lukowski, A.L.;
		Wymore, T.; Khare, D.; Brooks, C.L.; Palfey, B.A.; Smith, J.L.; Narayan,
		A.R.H.
Deposited on		
Resolution	:	1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.75 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 14287 atoms, of which 6831 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called FAD-dependent monooxygenase tropB.

Mol	Chain	Residues		Atoms						AltConf	Trace
1	А	431	Total 6750	-	Н 3341		O 610	S 23	0	5	0
1	В	430	Total 6762	-	Н 3350		0 612	S 22	0	7	0

• Molecule 2 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	А	1	Total 22	C 6	Н 14	O 2	0	0
2	А	1	Total 22	C 6	Н 14	O 2	0	0
2	В	1	Total 22	С 6	H 14	O 2	0	0
2	В	1	Total 22	С 6	Н 14	O 2	0	0


Continued on next page...



Continued from previous page...


Mol	Chain	Residues	A	Ator	ns		ZeroOcc	AltConf
2	В	1	Total 22	С 6	H 14	0 2	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C H O 14 3 8 3	0	0

• Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).





6NES

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	Δ	1	Total	С	Η	Ν	Ο	Р	0	0
4	A	1	84	27	31	9	15	2	0	
4	р	1	Total	С	Η	Ν	0	Р	0	0
4	В	1	84	27	31	9	15	2	0	

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Cl 1 1	0	0
5	В	1	Total Cl 1 1	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	218	Total O 218 218	0	0
6	В	263	Total O 263 263	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 2 2 21	Depositor	
Cell constants	70.81Å 184.49Å 164.13Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	46.43 - 1.75	Depositor	
% Data completeness	99.6 (46.43-1.75)	Depositor	
(in resolution range)	55.0 (40.40 1.10)	-	
R_{merge}	0.10	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$0.56 (at 1.75 \text{\AA})$	Xtriage	
Refinement program	PHENIX 1.14_3260	Depositor	
R, R_{free}	0.170 , 0.195	Depositor	
Wilson B-factor ($Å^2$)	30.0	Xtriage	
Anisotropy	0.316	Xtriage	
L-test for twinning ²	$ L > = 0.49, < L^2 > = 0.33$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	14287	wwPDB-VP	
Average B, all atoms $(Å^2)$	48.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mol	Turne	Chain	Res	Link	B	ond leng	gths	Bond angles				
	Type	Unam	nes	nes	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	HEZ	А	602	-	7,7,7	0.37	0	6,6,6	0.70	0		
2	HEZ	В	502	-	$7,\!7,\!7$	0.34	0	$6,\!6,\!6$	0.70	0		
3	GOL	А	603	-	$5,\!5,\!5$	1.34	1 (20%)	$5,\!5,\!5$	0.57	0		
2	HEZ	А	601	-	7,7,7	0.34	0	6,6,6	1.12	1 (16%)		
4	FAD	В	504	-	$53,\!58,\!58$	3.25	19 (35%)	68,89,89	2.04	15 (22%)		
4	FAD	А	604	-	$53,\!58,\!58$	3.24	18 (33%)	68,89,89	1.81	16 (23%)		
2	HEZ	В	501	-	7,7,7	0.41	0	6,6,6	0.72	0		
2	HEZ	В	503	-	7,7,7	0.33	0	6,6,6	0.69	0		

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEZ	А	602	-	-	3/5/5/5	-
2	HEZ	В	502	-	-	2/5/5/5	-
3	GOL	А	603	-	-	0/4/4/4	-
2	HEZ	А	601	-	-	0/5/5/5	-
4	FAD	В	504	-	-	1/30/50/50	0/6/6/6
4	FAD	А	604	-	-	1/30/50/50	0/6/6/6
2	HEZ	В	501	-	-	1/5/5/5	-
2	HEZ	В	503	-	-	2/5/5/5	-

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
4	В	504	FAD	C2B-C1B	-13.52	1.33	1.53
4	А	604	FAD	C2B-C1B	-13.32	1.33	1.53
4	В	504	FAD	O4B-C1B	12.35	1.58	1.41
4	А	604	FAD	O4B-C1B	11.89	1.57	1.41
4	В	504	FAD	C4X-N5	6.55	1.43	1.30

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
4	В	504	FAD	C5A-C6A-N6A	7.32	131.47	120.35

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	В	504	FAD	N3A-C2A-N1A	-5.58	119.96	128.68
4	А	604	FAD	N3A-C2A-N1A	-5.17	120.61	128.68
4	В	504	FAD	N6A-C6A-N1A	-4.83	108.54	118.57
4	В	504	FAD	C3B-C2B-C1B	4.69	108.04	100.98

Continued from previous page...

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	602	HEZ	C1-C2-C3-C4
2	В	502	HEZ	C1-C2-C3-C4
2	В	501	HEZ	C4-C5-C6-O6
2	А	602	HEZ	O1-C1-C2-C3
2	В	503	HEZ	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

