

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 - 02:49 AM EDT

PDB ID : 6NGA

Title: Structure of human neuronal nitric oxide synthase R354A/G357D mutant

heme domain in complex with 6-(3-(3-(dimethylamino)propyl)-2,6-difluo

rophenethyl)-4-methylpyridin-2-amine

Authors : Li, H.; Poulos, T.L.

Deposited on : 2018-12-21

Resolution : 1.98 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Nitric oxide synthase, brain.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace
1	A	413	Total 3374	C 2161	- 1	O 616	S 23	0	3	0
1	В	408	Total 3333	C 2136		O 608	S 21	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	ALA	ARG	engineered mutation	UNP P29475
A	357	ASP	GLY	engineered mutation	UNP P29475
В	354	ALA	ARG	engineered mutation	UNP P29475
В	357	ASP	GLY	engineered mutation	UNP P29475


• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total					0	0
		_	43	34	1	4	4		
2	D	1	Total	С	Fe	N	O	0	0
2	Б	1	43	34	1	4	4	0	U


• Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 17	9	5	3	0	0
3	В	1	Total 17		N 5		0	0

• Molecule 4 is 6-(2-{3-[3-(dimethylamino)propyl]-2,6-difluorophenyl}ethyl)-4-methylpyridin-2-amine (three-letter code: KL7) (formula: $C_{19}H_{25}F_2N_3$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 24			0	0
4	В	1	Total 24		F 2	0	0

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Zn 1 1	0	0

• Molecule 6 is water.

N	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	6	A	300	Total O 300 300	0	0
	6	В	209	Total O 209 209	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	52.19Å 122.77Å 164.67Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.82 - 1.98	Depositor
% Data completeness	99.9 (37.82-1.98)	Depositor
(in resolution range)	,	-
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) > 1$	1.42 (at 1.98Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, R_{free}	0.181 , 0.214	Depositor
Wilson B-factor ($Å^2$)	36.9	Xtriage
Anisotropy	0.588	Xtriage
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7385	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	$_{ m ths}$	Bond angles		
MIOI	туре	Chain	nes	LillK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	H4B	A	802	-	16,18,18	0.88	1 (6%)	11,26,26	2.49	5 (45%)
2	HEM	A	801	1	41,50,50	1.46	5 (12%)	45,82,82	1.82	6 (13%)
2	HEM	В	802	1	41,50,50	1.54	6 (14%)	45,82,82	1.81	9 (20%)
4	KL7	В	804	-	25,25,25	0.44	0	34,34,34	2.40	13 (38%)
3	H4B	В	803	-	16,18,18	1.01	1 (6%)	11,26,26	2.75	6 (54%)
4	KL7	A	803	-	25,25,25	0.52	0	34,34,34	2.37	11 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
2	HEM	A	801	1	-	2/12/54/54	-
2	HEM	В	802	1	-	0/12/54/54	-
4	KL7	В	804	-	-	1/11/11/11	0/2/2/2
3	H4B	В	803	-	-	0/8/17/17	0/2/2/2
4	KL7	A	803	-	-	5/11/11/11	0/2/2/2

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	A	801	HEM	C3C-C2C	-3.92	1.34	1.40
2	В	802	HEM	C3C-CAC	3.85	1.55	1.47
2	В	802	HEM	C3C-C2C	-3.81	1.35	1.40
2	A	801	HEM	C3C-CAC	3.26	1.54	1.47
2	A	801	HEM	CAB-C3B	2.87	1.55	1.47

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	A	801	HEM	CBA-CAA-C2A	-8.87	97.48	112.62
4	В	804	KL7	C11-C16-C15	-7.84	120.21	124.98
4	A	803	KL7	C11-C16-C15	-6.82	120.83	124.98
2	В	802	HEM	CBA-CAA-C2A	-5.84	102.65	112.62
3	В	803	H4B	C8A-C4A-C4	5.52	119.47	114.57



There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	KL7	C15-C17-C18-C19
4	A	803	KL7	C18-C19-N20-C22
4	A	803	KL7	C18-C19-N20-C21
4	В	804	KL7	C17-C18-C19-N20
4	A	803	KL7	C17-C18-C19-N20

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

