

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 09:12 AM EDT

PDB ID	:	6NHA
Title	:	Crystal structure of SYNT001, a human FcRn blocking monoclonal antibody
Authors	:	Blumberg, R.S.; Cheung, J.; Mahmood, A.; Gandhi, A.K.
Deposited on		
Resolution	:	2.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)		
Validation Pipeline (wwPDB-VP)	:	2.35.1
EDS Percentile statistics Ideal geometry (proteins)	: : :	FAILED 20191225.v01 (using entries in the PDB archive December 25th 2019) Engh & Huber (2001) Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.38 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called IgG receptor FcRn large subunit p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	256	Total 2014	C 1293	N 343	O 370	S 8	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	102	ALA	ASN	engineered mutation	UNP P55899
А	274	GLU	-	expression tag	UNP P55899
А	275	ASN	-	expression tag	UNP P55899
А	276	LEU	-	expression tag	UNP P55899
А	277	TYR	-	expression tag	UNP P55899
А	278	PHE	-	expression tag	UNP P55899
A	279	GLN	-	expression tag	UNP P55899

• Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	99	Total 828	C 528	N 140	0 157	${ m S} { m 3}$	0	0	0

• Molecule 3 is a protein called SYNT001-Fab light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	212	Total	С	N	0	S	0	0	0
	Ъ		1633	1023	274	331	5		Ū	U U

• Molecule 4 is a protein called SYNT001-Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	Н	211	Total 1583	C 999	N 265	0 314	${f S}{5}$	0	0	0



• Molecule 5 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula: $C_8H_{17}NO_3S$).



Mol	Chain	Residues	Ato	oms			ZeroOcc	AltConf
5	А	1	Total C 13 8			S 1	0	0
5	А	1	Total C 13 8	N 1		S 1	0	0
5	Н	1	Total C 13 8		0 3	S 1	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	33	Total O 33 33	0	0
6	В	10	Total O 10 10	0	0
6	L	69	Total O 69 69	0	0
6	Н	54	$\begin{array}{cc} \text{Total} & \text{O} \\ 54 & 54 \end{array}$	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	64.83Å 77.66Å 215.60Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.77 - 2.38	Depositor
% Data completeness	93.2 (49.77-2.38)	Depositor
(in resolution range)	55.2 (45.11-2.56)	-
R_{merge}	0.11	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.45 (at 2.39 \text{\AA})$	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.190 , 0.245	Depositor
Wilson B-factor $(Å^2)$	53.0	Xtriage
Anisotropy	0.261	Xtriage
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6263	wwPDB-VP
Average B, all atoms $(Å^2)$	60.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mol	Isl True Chain Das Link		Bo	ond leng	\mathbf{ths}	Bond angles				
	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NHE	Н	301	-	13,13,13	1.35	3 (23%)	$16,\!17,\!17$	1.90	6 (37%)
5	NHE	А	301	-	13,13,13	1.31	1 (7%)	$16,\!17,\!17$	1.84	4 (25%)
5	NHE	А	302	-	13,13,13	1.30	3 (23%)	16,17,17	2.06	4 (25%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NHE	Н	301	-	-	2/7/15/15	0/1/1/1
5	NHE	А	301	-	-	4/7/15/15	0/1/1/1
5	NHE	А	302	-	-	4/7/15/15	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
5	А	301	NHE	C2-S	3.31	1.82	1.77
5	А	302	NHE	C2-S	3.06	1.81	1.77
5	Н	301	NHE	C2-S	3.00	1.81	1.77
5	Н	301	NHE	O1-S	2.36	1.52	1.45
5	Н	301	NHE	O2-S	2.20	1.51	1.45

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	302	NHE	01-S-C2	4.98	112.92	106.92
5	А	302	NHE	02-S-01	-4.31	99.05	113.95
5	Н	301	NHE	02-S-01	-3.50	101.83	113.95
5	А	301	NHE	O3-S-C2	3.31	111.11	105.77
5	А	301	NHE	01-S-C2	3.23	110.80	106.92

There are no chirality outliers.

5 of 10 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	А	301	NHE	C1-C2-S-O1
5	А	301	NHE	C1-C2-S-O2
5	А	301	NHE	C1-C2-S-O3
5	А	302	NHE	N-C1-C2-S
5	Н	301	NHE	N-C1-C2-S

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

