



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2023 – 12:51 PM EDT

PDB ID : 6NLH
Title : Structure of human triose phosphate isomerase R189A
Authors : Richards, K.R.; Roland, B.P.; Palladino, M.J.; VanDemark, A.P.
Deposited on : 2019-01-08
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 29986 atoms, of which 14602 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	244	3671	1161	1838	314	351	7	0	0	0
1	E	238	3589	1137	1795	307	343	7	0	0	0
1	B	245	3695	1167	1851	318	352	7	0	0	0
1	C	239	3612	1143	1807	311	344	7	0	0	0
1	D	245	3695	1167	1851	318	352	7	0	0	0
1	F	239	3612	1143	1807	311	344	7	0	0	0
1	G	245	3695	1167	1851	318	352	7	0	0	0
1	H	239	3603	1141	1802	308	345	7	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	189	ALA	ARG	engineered mutation	UNP P60174
E	189	ALA	ARG	engineered mutation	UNP P60174
B	189	ALA	ARG	engineered mutation	UNP P60174
C	189	ALA	ARG	engineered mutation	UNP P60174
D	189	ALA	ARG	engineered mutation	UNP P60174
F	189	ALA	ARG	engineered mutation	UNP P60174
G	189	ALA	ARG	engineered mutation	UNP P60174
H	189	ALA	ARG	engineered mutation	UNP P60174

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Na 3 3	0	0
2	E	1	Total Na 1 1	0	0
2	B	4	Total Na 4 4	0	0
2	C	3	Total Na 3 3	0	0
2	D	3	Total Na 3 3	0	0
2	F	2	Total Na 2 2	0	0
2	G	2	Total Na 2 2	0	0
2	H	2	Total Na 2 2	0	0

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total Br 1 1	0	0
3	C	1	Total Br 1 1	0	0
3	F	1	Total Br 1 1	0	0
3	H	1	Total Br 1 1	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	F	1	Total O P 5 4 1	0	0
4	H	1	Total O P 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	118	Total O 118 118	0	0
5	E	79	Total O 79 79	0	0
5	B	114	Total O 114 114	0	0
5	C	86	Total O 86 86	0	0
5	D	97	Total O 97 97	0	0
5	F	85	Total O 85 85	0	0
5	G	106	Total O 106 106	0	0
5	H	85	Total O 85 85	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.14Å 73.66Å 92.84Å 90.03° 90.03° 90.00°	Depositor
Resolution (Å)	19.66 – 2.20	Depositor
% Data completeness (in resolution range)	95.9 (19.66-2.20)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.54 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.174 , 0.216	Depositor
Wilson B-factor (Å ²)	21.8	Xtrriage
Anisotropy	0.614	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.478 for h,-k,-l 0.477 for -h,k,-l 0.478 for -h,-k,l	Xtrriage
Total number of atoms	29986	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 24 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	H	304	-	4,4,4	0.84	0	6,6,6	0.49	0
4	PO4	C	305	-	4,4,4	0.67	0	6,6,6	0.47	0
4	PO4	E	303	-	4,4,4	0.87	0	6,6,6	0.40	0
4	PO4	F	304	-	4,4,4	0.71	0	6,6,6	0.61	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.