

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 2, 2023 – 04:28 PM EDT

PDB ID	:	6NLY
Title	:	Fragment of human mitochondrial Alanyl-tRNA Synthetase C-Ala domain
Authors	:	Kuhle, B.; Schimmel, P.
Deposited on		
Resolution	:	2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	FAILED
:	1.13
:	FAILED
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	Engh & Huber $(2001)$
:	Parkinson et al. (1996)
:	2.35.1
	:::::::::::::::::::::::::::::::::::::::

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 2.31 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1 B	178	Total	С	Ν	0	$\mathbf{S}$	0	0	0
	D		1339	836	244	249	10			
1	Λ	181	Total	С	Ν	0	S	0	0	0
			1357	847	247	253	10			
1	1 C	C 176	Total	С	Ν	0	S	0	0	0
			1324	827	240	247	10	0	0	0
1 D	182	Total	С	Ν	0	S	0	0	0	
		102	1384	863	258	253	10		0	U

• Molecule 1 is a protein called Alanine–tRNA ligase, mitochondrial.

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	986	LEU	-	expression tag	UNP Q5JTZ9
В	987	GLU	-	expression tag	UNP Q5JTZ9
В	988	HIS	-	expression tag	UNP Q5JTZ9
В	989	HIS	-	expression tag	UNP Q5JTZ9
В	990	HIS	-	expression tag	UNP Q5JTZ9
В	991	HIS	-	expression tag	UNP Q5JTZ9
В	992	HIS	-	expression tag	UNP Q5JTZ9
В	993	HIS	-	expression tag	UNP Q5JTZ9
А	986	LEU	-	expression tag	UNP Q5JTZ9
А	987	GLU	-	expression tag	UNP Q5JTZ9
А	988	HIS	-	expression tag	UNP Q5JTZ9
А	989	HIS	-	expression tag	UNP Q5JTZ9
A	990	HIS	-	expression tag	UNP Q5JTZ9
А	991	HIS	-	expression tag	UNP Q5JTZ9
A	992	HIS	-	expression tag	UNP Q5JTZ9
А	993	HIS	-	expression tag	UNP Q5JTZ9
С	986	LEU	-	expression tag	UNP Q5JTZ9
С	987	GLU	-	expression tag	UNP Q5JTZ9
С	988	HIS	-	expression tag	UNP Q5JTZ9
С	989	HIS	-	expression tag	UNP Q5JTZ9
С	990	HIS	-	expression tag	UNP Q5JTZ9

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	991	HIS	-	expression tag	UNP Q5JTZ9
С	992	HIS	-	expression tag	UNP Q5JTZ9
С	993	HIS	-	expression tag	UNP Q5JTZ9
D	986	LEU	-	expression tag	UNP Q5JTZ9
D	987	GLU	-	expression tag	UNP Q5JTZ9
D	988	HIS	-	expression tag	UNP Q5JTZ9
D	989	HIS	-	expression tag	UNP Q5JTZ9
D	990	HIS	-	expression tag	UNP Q5JTZ9
D	991	HIS	-	expression tag	UNP Q5JTZ9
D	992	HIS	-	expression tag	UNP Q5JTZ9
D	993	HIS	-	expression tag	UNP Q5JTZ9

Continued from previous page...

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	5	Total O 5 5	0	0
2	А	6	Total O 6 6	0	0
2	С	4	Total O 4 4	0	0
2	D	6	Total O 6 6	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	103.22Å 317.23Å 54.71Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	38.02 - 2.31	Depositor
% Data completeness	99.1 (38.02-2.31)	Depositor
(in resolution range)		Depositor
R <sub>merge</sub>	0.04	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.27 (at 2.31 \text{\AA})$	Xtriage
Refinement program	PHENIX	Depositor
$R, R_{free}$	0.209 , $0.257$	Depositor
Wilson B-factor $(Å^2)$	50.2	Xtriage
Anisotropy	0.848	Xtriage
L-test for twinning <sup>2</sup>	$ L  > = 0.49, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5425	wwPDB-VP
Average B, all atoms $(Å^2)$	77.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 21.11 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.5256e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 4.6 Ligand geometry (i)

There are no ligands in this entry.

#### 4.7 Other polymers (i)

There are no such residues in this entry.



## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

## 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

## 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

