

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 2, 2023 – 05:38 PM EDT

PDB ID : 6NMY

Title: A Cytokine-receptor complex

Authors: Dhagat, U.; Kan, W.L.; Hercus, T.R.; Broughton, S.E.; Nero, T.L.; Lopez,

A.F.; Parker, M.W.

Deposited on : 2019-01-13

Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 12975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Interleukin-3 receptor subunit alpha.

Mol	Chain	Residues		$\mathbf{At}$	oms		ZeroOcc	AltConf	Trace	
1	F	270	Total 2189	C 1381	N 392	O 400	S 16	0	0	0
1	M	262	Total 2128	C 1341	N 383	O 388	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	ain Residue Modelled Actual		Comment	Reference	
F	212	GLN	ASN	engineered mutation	UNP P26951
M	212	GLN	ASN	engineered mutation	UNP P26951

• Molecule 2 is a protein called Cytokine receptor common subunit beta.

Mol	Chain	Residues		At	oms		ZeroOcc	AltConf	Trace	
9	Λ	215	Total	Total C N C	О	S	0	0	0	
	A	210	1722	1079	304	329	10	0	0	
9	С	215	Total	С	N	О	S	0	0	0
$\begin{vmatrix} 2 \end{vmatrix}$	C	C 215	1722	1079	304	329	10	0		

• Molecule 3 is a protein called Interleukin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	3 I 1	108	Total	С	N	О	S	0	0	0
3		100	874	554	155	161	4	0	U	
2	Т	108	Total	С	N	О	S	0	0	0
3	J	J 108	874	554	155	161	4	0	U	U

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	8	GLY	-	expression tag	UNP P08700
I	9	ALA	-	expression tag	UNP P08700



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Chain	Residue	Modelled	Actual	Comment	Reference
I	10	MET	-	expression tag	UNP P08700
I	11	GLY	-	expression tag	UNP P08700
I	13	TYR	TRP	engineered mutation	UNP P08700
J	8	GLY	-	expression tag	UNP P08700
J	9	ALA	-	expression tag	UNP P08700
J	10	MET	-	expression tag	UNP P08700
J	11	GLY	-	expression tag	UNP P08700
J	13	TYR	TRP	engineered mutation	UNP P08700

• Molecule 4 is a protein called Cytokine receptor common subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	196	Total	С	N	О	S	0	0	0
4	Б	190	1580	987	283	302	8	0		
4	D	107	Total	С	N	О	S	0	0	0
4	D	D 197	1589	992	284	305	8	0	U	

There are 2 discrepancies between the modelled and reference sequences:

$\mathbf{C}$	hain	Residue	Modelled	Actual	Comment	Reference	
	В	346	GLN	ASN	engineered mutation	UNP P32927	
	D	346	GLN	ASN	engineered mutation	UNP P32927	

• Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	4	Total	С	N	О	0	0	0
)	E	4	49	28	2	19	0	Ü	U
	C	4	Total	С	N	О	0	0	0
5	G	G 4	49	28	2	19	0	U	U

• Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	Н	3	Total 39	C 22			0	0	0
6	K	3	Total 39	C 22			0	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	N	3	Total 39	C 22	N 2	O 15	0	0	0

• Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.

M	Iol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
,	7	L	2	Total 28			0	0	0
,	7	О	2	Total 28	C 16		0	0	0

• Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	С	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	F	2	Total O 2 2	0	0
9	M	2	Total O 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total O 1 1	0	0
9	С	1	Total O 1 1	0	0
9	I	1	Total O 1 1	0	0
9	J	2	Total O 2 2	0	0
9	В	2	Total O 2 2	0	0
9	D	1	Total O 1 1	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



## 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source		
Space group	P 21 21 21	Depositor		
Cell constants	111.65Å 157.25Å 168.31Å	Depositor		
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor		
Resolution (Å)	47.45 - 3.30	Depositor		
% Data completeness	99.8 (47.45-3.30)	Depositor		
(in resolution range)	` '			
$R_{merge}$	0.17	Depositor		
$R_{sym}$	(Not available)	Depositor		
$< I/\sigma(I) > 1$	1.51  (at  3.33Å)	Xtriage		
Refinement program	PHENIX	Depositor		
$R, R_{free}$	0.236 , $0.278$	Depositor		
Wilson B-factor $(A^2)$	95.0	Xtriage		
Anisotropy	0.224	Xtriage		
L-test for twinning <sup>2</sup>	$ < L >=0.46, < L^2>=0.29$	Xtriage		
Estimated twinning fraction	No twinning to report.	Xtriage		
Total number of atoms	12975	wwPDB-VP		
Average B, all atoms $(\mathring{A}^2)$	95.0	wwPDB-VP		

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

## 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 4.5 Carbohydrates (i)

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Trino	Chain	Dag	Link	Во	nd leng	ths	В	ond ang	les
IVIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	Е	1	5,1	14,14,15	0.29	0	17,19,21	0.55	0
5	NAG	Е	2	5	14,14,15	0.48	0	17,19,21	0.59	0
5	BMA	E	3	5	11,11,12	1.09	1 (9%)	15,15,17	0.99	1 (6%)
5	FUC	Е	4	5	10,10,11	0.79	0	14,14,16	0.85	0
5	NAG	G	1	5,1	14,14,15	1.02	1 (7%)	17,19,21	1.39	3 (17%)
5	NAG	G	2	5	14,14,15	0.31	0	17,19,21	0.38	0
5	BMA	G	3	5	11,11,12	0.70	0	15,15,17	0.72	0
5	FUC	G	4	5	10,10,11	1.78	2 (20%)	14,14,16	1.74	3 (21%)
6	NAG	Н	1	1,6	14,14,15	0.73	0	17,19,21	1.12	2 (11%)
6	NAG	Н	2	6	14,14,15	0.44	0	17,19,21	1.28	1 (5%)
6	BMA	Н	3	6	11,11,12	0.79	0	15,15,17	0.87	0
6	NAG	K	1	1,6	14,14,15	0.36	0	17,19,21	0.83	1 (5%)
6	NAG	K	2	6	14,14,15	0.31	0	17,19,21	0.67	0
6	BMA	K	3	6	11,11,12	0.80	0	15,15,17	0.83	0
7	NAG	L	1	1,7	14,14,15	0.39	0	17,19,21	0.57	0
7	NAG	L	2	7	14,14,15	0.63	0	17,19,21	0.43	0
6	NAG	N	1	2,6	14,14,15	0.63	1 (7%)	17,19,21	0.91	1 (5%)
6	NAG	N	2	6	14,14,15	1.64	2 (14%)	17,19,21	1.12	3 (17%)
6	BMA	N	3	6	11,11,12	0.90	0	15,15,17	1.26	1 (6%)
7	NAG	О	1	2,7	14,14,15	0.35	0	17,19,21	0.70	0
7	NAG	О	2	7	14,14,15	0.24	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	Е	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	Е	2	5	-	4/6/23/26	0/1/1/1
5	BMA	Е	3	5	-	1/2/19/22	0/1/1/1
5	FUC	Е	4	5	-	-	0/1/1/1
5	NAG	G	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	G	2	5	-	0/6/23/26	0/1/1/1
5	BMA	G	3	5	-	2/2/19/22	0/1/1/1
5	FUC	G	4	5	-	-	0/1/1/1
6	NAG	Н	1	1,6	-	4/6/23/26	0/1/1/1
6	NAG	Н	2	6	-	2/6/23/26	0/1/1/1
6	BMA	Н	3	6	-	0/2/19/22	0/1/1/1
6	NAG	K	1	1,6	-	1/6/23/26	0/1/1/1



 $Continued\ from\ previous\ page...$ 

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	K	2	6	-	3/6/23/26	0/1/1/1
6	BMA	K	3	6	-	0/2/19/22	0/1/1/1
7	NAG	L	1	1,7	-	2/6/23/26	0/1/1/1
7	NAG	L	2	7	-	0/6/23/26	0/1/1/1
6	NAG	N	1	2,6	-	1/6/23/26	0/1/1/1
6	NAG	N	2	6	-	2/6/23/26	0/1/1/1
6	BMA	N	3	6	-	2/2/19/22	0/1/1/1
7	NAG	О	1	2,7	-	1/6/23/26	0/1/1/1
7	NAG	О	2	7	-	0/6/23/26	0/1/1/1

### All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
6	N	2	NAG	O5-C1	-5.34	1.35	1.43
5	G	4	FUC	C1-C2	3.90	1.61	1.52
5	G	4	FUC	O5-C1	3.59	1.49	1.43
5	G	1	NAG	O5-C1	-3.53	1.38	1.43
5	Е	3	BMA	C1-C2	3.12	1.59	1.52
6	N	2	NAG	C1-C2	2.33	1.55	1.52
6	N	1	NAG	C1-C2	2.07	1.55	1.52

#### All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	Н	2	NAG	C1-O5-C5	4.44	118.21	112.19
5	G	4	FUC	C1-O5-C5	4.17	122.23	112.78
5	G	1	NAG	C1-O5-C5	3.77	117.31	112.19
5	G	4	FUC	C1-C2-C3	3.34	113.77	109.67
6	N	3	BMA	C1-C2-C3	-3.06	105.91	109.67
6	Н	1	NAG	O4-C4-C5	2.94	116.60	109.30
6	Н	1	NAG	C1-O5-C5	2.83	116.03	112.19
6	N	1	NAG	C1-O5-C5	2.81	116.00	112.19
6	N	2	NAG	C4-C3-C2	2.78	115.09	111.02
6	N	2	NAG	O5-C5-C4	-2.52	104.71	110.83
5	G	1	NAG	O4-C4-C3	-2.42	104.76	110.35
5	Е	3	BMA	O2-C2-C3	-2.38	105.38	110.14
5	G	4	FUC	O5-C5-C4	2.32	113.68	109.52
6	K	1	NAG	C1-O5-C5	2.19	115.16	112.19
5	G	1	NAG	C3-C4-C5	2.17	114.11	110.24
6	N	2	NAG	O3-C3-C2	-2.02	105.28	109.47



There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	Е	2	NAG	O5-C5-C6-O6
7	L	1	NAG	O5-C5-C6-O6
6	N	2	NAG	O5-C5-C6-O6
6	N	2	NAG	C4-C5-C6-O6
6	Н	1	NAG	O5-C5-C6-O6
6	N	3	BMA	C4-C5-C6-O6
6	Н	2	NAG	O5-C5-C6-O6
7	L	1	NAG	C4-C5-C6-O6
6	N	3	BMA	O5-C5-C6-O6
6	Н	1	NAG	C4-C5-C6-O6
5	Е	2	NAG	C4-C5-C6-O6
5	G	3	BMA	C4-C5-C6-O6
6	Н	2	NAG	C4-C5-C6-O6
6	Н	1	NAG	C1-C2-N2-C7
5	G	3	BMA	O5-C5-C6-O6
5	Е	3	BMA	O5-C5-C6-O6
6	K	2	NAG	O5-C5-C6-O6
5	Е	2	NAG	C1-C2-N2-C7
6	K	1	NAG	O5-C5-C6-O6
6	N	1	NAG	C3-C2-N2-C7
7	О	1	NAG	C3-C2-N2-C7
6	K	2	NAG	C1-C2-N2-C7
5	Е	2	NAG	C3-C2-N2-C7
6	Н	1	NAG	C3-C2-N2-C7
6	K	2	NAG	C3-C2-N2-C7


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





























## 4.6 Ligand geometry (i)

#### 1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Trmo	Chain	Dog	Tiple	Bo	ond leng	$ ag{ths}$	Bond angles			
		Type		nes	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
	8	NAG	С	301	2	14,14,15	0.46	0	17,19,21	0.65	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	С	301	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
8	С	301	NAG	C1-O5-C5	2.30	115.30	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

#### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

## 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

