



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2023 – 12:50 PM EDT

PDB ID : 2NRX
Title : Crystal structure of the C-terminal half of UvrC, in the presence of sulfate molecules
Authors : Karakas, E.; Truglio, J.J.; Kisker, C.
Deposited on : 2006-11-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	 % 78% 19% ..
1	B	220	 6% 73% 19% .. 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	905	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UvrABC system protein C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	217	1763	1119	323	318	3	0	0	0
1	B	208	1683	1071	304	305	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	558	GLY	-	cloning artifact	UNP Q9WYA3
B	558	GLY	-	cloning artifact	UNP Q9WYA3

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	176	Total O 176 176	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	158	Total 158	O 158	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UvrABC system protein C



- Molecule 1: UvrABC system protein C



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.36Å 81.02Å 99.62Å 90.00° 98.02° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 37.47 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.2 (40.00-1.90) 97.2 (37.47-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 1.89Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.240 0.184 , 0.239	Depositor DCC
R_{free} test set	2145 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.030 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3826	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.81	1/1795 (0.1%)	0.81	1/2414 (0.0%)
1	B	0.72	0/1713	0.78	2/2307 (0.1%)
All	All	0.77	1/3508 (0.0%)	0.79	3/4721 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	449	CYS	CB-SG	-6.36	1.71	1.82

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	495	HIS	N-CA-C	6.98	129.83	111.00
1	B	344	LYS	CD-CE-NZ	6.00	125.50	111.70
1	A	380	LEU	CA-CB-CG	-5.42	102.84	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	495	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1763	0	1816	54	0
1	B	1683	0	1721	38	0
2	A	20	0	0	3	1
2	B	20	0	0	0	0
3	A	6	0	8	4	0
4	A	176	0	0	14	1
4	B	158	0	0	7	0
All	All	3826	0	3545	89	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:GLU:H	1:A:345:GLU:CD	1.70	0.94
1:B:514:ILE:HG22	1:B:519:LYS:HG3	1.52	0.92
1:B:376:THR:HG23	1:B:400:GLN:HE21	1.36	0.91
1:A:460:THR:HG21	1:A:467:GLU:HB3	1.55	0.88
1:A:516:PRO:O	1:A:520:LYS:HG2	1.73	0.88
1:B:370:HIS:ND1	1:B:376:THR:HG22	1.91	0.85
1:B:342:LEU:N	4:B:311:HOH:O	2.13	0.82
1:A:460:THR:CG2	1:A:467:GLU:HB3	2.10	0.81
1:B:514:ILE:CG2	1:B:519:LYS:HG3	2.11	0.81
1:B:362:ARG:H	1:B:424:ASN:HD22	1.28	0.81
1:B:376:THR:CG2	1:B:400:GLN:HE21	1.95	0.79
1:B:525:HIS:HE1	1:B:539:GLU:OE2	1.70	0.75
1:B:525:HIS:HD2	4:B:152:HOH:O	1.70	0.75
1:A:469:HIS:HB3	4:A:78:HOH:O	1.88	0.74
1:A:422:LEU:HD11	1:A:447:LYS:HB3	1.69	0.73
1:B:495:HIS:N	1:B:496:ARG:HB2	2.07	0.70
1:A:466:ARG:HH12	1:A:468:ILE:HD11	1.57	0.70
1:A:478:ARG:HG2	4:A:322:HOH:O	1.93	0.69
1:A:474:HIS:HD2	1:A:476:VAL:H	1.39	0.69
1:A:466:ARG:NH1	1:A:468:ILE:HD11	2.08	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:422:LEU:HD11	1:B:447:LYS:HB3	1.76	0.68
1:A:348:GLU:HA	1:A:351:MET:HE3	1.77	0.67
1:A:525:HIS:HD2	4:A:136:HOH:O	1.77	0.67
1:A:348:GLU:HA	1:A:351:MET:CE	2.26	0.66
1:A:525:HIS:HE1	1:A:539:GLU:OE2	1.79	0.66
1:A:460:THR:OG1	1:A:469:HIS:HD2	1.79	0.65
1:B:376:THR:HG21	4:B:42:HOH:O	1.98	0.64
1:A:473:ASP:OD1	1:A:478:ARG:NH2	2.31	0.63
1:A:478:ARG:CG	4:A:322:HOH:O	2.47	0.63
1:B:514:ILE:HD11	1:B:553:VAL:HG22	1.81	0.61
1:A:355:ASN:ND2	4:A:313:HOH:O	2.26	0.61
1:B:514:ILE:HD11	1:B:553:VAL:CG2	2.30	0.61
1:A:357:LYS:HG3	1:B:421:PRO:HB2	1.84	0.60
1:B:495:HIS:CA	1:B:496:ARG:HB2	2.31	0.59
1:B:524:GLU:OE1	4:B:320:HOH:O	2.16	0.57
1:A:482:GLN:NE2	4:A:41:HOH:O	2.29	0.57
1:A:548:GLU:HG2	4:A:242:HOH:O	2.05	0.56
1:B:495:HIS:H	1:B:496:ARG:HB2	1.70	0.56
1:A:343:ARG:HB3	1:A:345:GLU:OE1	2.07	0.55
1:A:460:THR:OG1	1:A:469:HIS:CD2	2.58	0.54
1:B:348:GLU:HA	1:B:351:MET:HE3	1.88	0.54
1:A:362:ARG:NH2	3:A:905:GOL:H12	2.23	0.53
1:A:345:GLU:OE1	1:A:345:GLU:N	2.38	0.53
1:A:349:GLU:HG3	4:A:238:HOH:O	2.09	0.52
1:B:368:ILE:HG13	1:B:409:ILE:HG12	1.91	0.52
1:A:361:TYR:OH	1:B:344:LYS:HE3	2.10	0.52
1:A:434:GLN:HG3	2:A:903:SO4:O4	2.12	0.48
1:A:488:HIS:HB3	4:A:223:HOH:O	2.13	0.48
1:A:422:LEU:HD13	1:A:449:CYS:HB2	1.95	0.48
1:B:398:ILE:HD12	1:B:411:THR:HG21	1.96	0.48
1:A:420:HIS:CE1	3:A:905:GOL:C3	2.96	0.48
1:B:348:GLU:HA	1:B:351:MET:CE	2.43	0.48
1:B:547:THR:O	1:B:551:ARG:HG3	2.14	0.47
1:B:342:LEU:HD22	1:B:479:LEU:HD12	1.95	0.47
1:A:474:HIS:CD2	1:A:476:VAL:H	2.27	0.47
1:A:460:THR:HG23	1:A:467:GLU:HB3	1.94	0.47
1:B:353:LEU:HD22	1:B:476:VAL:HG11	1.97	0.47
1:A:385:ASP:OD2	1:B:344:LYS:HE2	2.15	0.46
1:A:345:GLU:CD	1:A:345:GLU:N	2.51	0.46
1:A:362:ARG:HH22	3:A:905:GOL:H12	1.81	0.46
1:B:402:HIS:N	4:B:318:HOH:O	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:GLU:HG2	4:A:260:HOH:O	2.14	0.45
1:A:458:GLU:O	1:A:460:THR:N	2.50	0.45
1:A:474:HIS:HE1	4:A:241:HOH:O	1.99	0.45
1:A:514:ILE:HD13	1:A:549:ILE:HG21	1.99	0.45
1:B:372:GLN:HG3	4:B:208:HOH:O	2.17	0.44
1:B:351:MET:HE1	1:B:359:PHE:HD1	1.81	0.44
1:A:460:THR:HG21	1:A:467:GLU:CB	2.38	0.44
1:B:370:HIS:HA	1:B:376:THR:HA	2.00	0.43
1:A:496:ARG:NH1	4:A:321:HOH:O	2.39	0.43
1:B:511:VAL:HG22	1:B:514:ILE:HG13	2.01	0.43
1:A:468:ILE:HA	4:A:297:HOH:O	2.18	0.42
1:B:443:LYS:HG3	4:B:155:HOH:O	2.19	0.42
1:A:409:ILE:HD11	1:A:434:GLN:OE1	2.20	0.42
1:A:514:ILE:HD11	1:A:518:ARG:HB3	2.00	0.42
1:B:362:ARG:H	1:B:424:ASN:ND2	2.07	0.42
1:A:429:ASP:CG	1:A:488:HIS:HE2	2.22	0.42
1:B:390:LYS:HA	1:B:393:TYR:CD1	2.55	0.41
1:A:395:ARG:NH2	2:A:902:SO4:O1	2.50	0.41
1:A:516:PRO:HD2	2:A:901:SO4:O3	2.21	0.41
1:B:432:ILE:HA	1:B:462:VAL:HG11	2.01	0.41
1:B:511:VAL:HB	1:B:556:ILE:HD13	2.02	0.41
1:A:381:VAL:HG22	1:A:487:THR:HG22	2.02	0.41
1:A:409:ILE:HD11	1:A:434:GLN:CD	2.41	0.41
1:A:351:MET:HE3	1:A:351:MET:HB3	1.85	0.41
1:B:414:LYS:O	1:B:418:SER:HB3	2.21	0.40
1:A:342:LEU:HD22	1:A:479:LEU:HD12	2.03	0.40
1:A:420:HIS:CE1	3:A:905:GOL:H31	2.56	0.40
1:A:474:HIS:CE1	4:A:241:HOH:O	2.74	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:904:SO4:O2	4:A:297:HOH:O[1_655]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/220 (98%)	210 (98%)	5 (2%)	0	100	100
1	B	202/220 (92%)	197 (98%)	5 (2%)	0	100	100
All	All	417/440 (95%)	407 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/195 (99%)	188 (97%)	5 (3%)	46	39
1	B	185/195 (95%)	175 (95%)	10 (5%)	22	13
All	All	378/390 (97%)	363 (96%)	15 (4%)	31	22

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	345	GLU
1	A	351	MET
1	A	380	LEU
1	A	466	ARG
1	A	498	ARG
1	B	344	LYS
1	B	351	MET
1	B	376	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	380	LEU
1	B	448	ASP
1	B	460	THR
1	B	466	ARG
1	B	505	ARG
1	B	511	VAL
1	B	537	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	372	GLN
1	A	402	HIS
1	A	469	HIS
1	A	474	HIS
1	A	525	HIS
1	B	400	GLN
1	B	402	HIS
1	B	424	ASN
1	B	436	ASN
1	B	525	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	903	-	4,4,4	0.19	0	6,6,6	0.25	0
2	SO4	B	904	-	4,4,4	0.35	0	6,6,6	0.75	0
3	GOL	A	905	-	5,5,5	0.58	0	5,5,5	0.73	0
2	SO4	B	901	-	4,4,4	0.20	0	6,6,6	0.33	0
2	SO4	A	904	-	4,4,4	0.29	0	6,6,6	0.81	0
2	SO4	A	901	-	4,4,4	0.15	0	6,6,6	0.27	0
2	SO4	A	902	-	4,4,4	0.17	0	6,6,6	0.56	0
2	SO4	B	902	-	4,4,4	0.13	0	6,6,6	0.22	0
2	SO4	A	903	-	4,4,4	0.06	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	905	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	905	GOL	O1-C1-C2-C3
3	A	905	GOL	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	905	GOL	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	904	SO4	0	1
2	A	901	SO4	1	0
2	A	902	SO4	1	0
2	A	903	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/220 (98%)	0.07	3 (1%) 75 77	15, 22, 32, 45	0
1	B	208/220 (94%)	0.28	14 (6%) 17 20	19, 28, 37, 41	0
All	All	425/440 (96%)	0.17	17 (4%) 38 41	15, 25, 35, 45	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	529	LEU	3.5
1	B	373	GLY	3.1
1	A	498	ARG	3.0
1	B	372	GLN	2.8
1	A	459	GLU	2.7
1	B	496	ARG	2.7
1	B	371	LEU	2.6
1	B	466	ARG	2.6
1	B	530	GLU	2.5
1	B	370	HIS	2.4
1	B	401	ASP	2.4
1	B	402	HIS	2.2
1	B	398	ILE	2.2
1	A	466	ARG	2.2
1	B	406	TYR	2.2
1	B	375	TYR	2.1
1	B	403	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	902	5/5	0.79	0.35	69,70,72,72	0
2	SO4	A	901	5/5	0.87	0.17	63,65,66,66	0
2	SO4	B	902	5/5	0.87	0.11	58,59,60,61	0
3	GOL	A	905	6/6	0.87	0.17	31,39,40,42	0
2	SO4	B	903	5/5	0.94	0.13	50,50,51,51	0
2	SO4	B	901	5/5	0.94	0.19	42,43,45,46	0
2	SO4	A	903	5/5	0.96	0.11	32,37,38,38	0
2	SO4	A	904	5/5	0.98	0.09	20,20,30,30	0
2	SO4	B	904	5/5	0.99	0.09	21,22,27,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.