



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 07:22 PM EDT

PDB ID : 4NRW  
Title : MvNei1-G86D  
Authors : Prakash, A.; Doubleie, S.  
Deposited on : 2013-11-27  
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	 15% 82% 15%
1	B	294	 86% 11%
2	C	13	 92% 8%
2	E	13	 92% 8%
3	D	13	 15% 69% 31%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	13	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a red segment on the left labeled '15%', a large green segment labeled '77%', a yellow segment labeled '15%', and a small orange segment on the right labeled '8%'.</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called formamidopyrimidine-DNA glycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	Total	C	N	O	S	0	0	0
			2336	1516	381	434	5			
1	B	285	Total	C	N	O	S	0	0	0
			2346	1525	383	433	5			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	ASP	GLY	engineered mutation	UNP Q5UQ00
A	288	LEU	-	expression tag	UNP Q5UQ00
A	289	GLU	-	expression tag	UNP Q5UQ00
A	290	HIS	-	expression tag	UNP Q5UQ00
A	291	HIS	-	expression tag	UNP Q5UQ00
A	292	HIS	-	expression tag	UNP Q5UQ00
A	293	HIS	-	expression tag	UNP Q5UQ00
A	294	HIS	-	expression tag	UNP Q5UQ00
A	295	HIS	-	expression tag	UNP Q5UQ00
B	86	ASP	GLY	engineered mutation	UNP Q5UQ00
B	288	LEU	-	expression tag	UNP Q5UQ00
B	289	GLU	-	expression tag	UNP Q5UQ00
B	290	HIS	-	expression tag	UNP Q5UQ00
B	291	HIS	-	expression tag	UNP Q5UQ00
B	292	HIS	-	expression tag	UNP Q5UQ00
B	293	HIS	-	expression tag	UNP Q5UQ00
B	294	HIS	-	expression tag	UNP Q5UQ00
B	295	HIS	-	expression tag	UNP Q5UQ00

- Molecule 2 is a DNA chain called 5'-D(\*GP\*TP\*AP\*GP\*AP\*CP\*CP\*TP\*GP\*GP\*AP\*C P\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	13	Total	C	N	O	P	0	0	0
			267	127	53	75	12			
2	E	13	Total	C	N	O	P	0	0	0
			267	127	53	75	12			

- Molecule 3 is a DNA chain called 5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(3DR)P\*GP\*TP\*CP\*TP\*AP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	13	Total	C	N	O	P	0	0	0
			249	120	41	76	12			
3	F	13	Total	C	N	O	P	0	0	0
			249	120	41	76	12			



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		
4	B	5	Total	O	0	0
			5	5		





- Molecule 3: 5'-D(\*CP\*GP\*TP\*CP\*CP\*AP\*(3DR)P\*GP\*TP\*CP\*TP\*AP\*C)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.40Å 121.68Å 79.99Å 90.00° 95.68° 90.00°	Depositor
Resolution (Å)	14.97 – 2.85 14.97 – 2.85	Depositor EDS
% Data completeness (in resolution range)	98.9 (14.97-2.85) 99.0 (14.97-2.85)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 2.86Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.260 , 0.315 0.256 , 0.306	Depositor DCC
$R_{free}$ test set	1759 reflections (10.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.6	Xtrriage
Anisotropy	0.935	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.83	EDS
Total number of atoms	5722	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/2398	0.39	1/3238 (0.0%)
1	B	0.22	0/2408	0.37	0/3247
2	C	0.44	0/300	0.95	0/462
2	E	0.45	0/300	0.95	0/462
3	D	0.45	0/264	1.03	0/402
3	F	0.44	0/264	1.04	0/402
All	All	0.28	0/5934	0.57	1/8213 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	ASP	CB-CG-OD2	5.16	122.95	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2336	0	2289	27	0
1	B	2346	0	2320	18	0
2	C	267	0	147	1	0
2	E	267	0	147	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	249	0	145	2	0
3	F	249	0	145	2	0
4	A	3	0	0	0	0
4	B	5	0	0	0	0
All	All	5722	0	5193	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:14:DC:H2''	3:F:15:DG:H5'	1.67	0.76
1:A:98:ALA:HB3	1:A:110:PHE:HB3	1.79	0.65
1:B:39:GLU:OE1	1:B:120:LYS:HE3	1.98	0.63
1:A:100:LEU:HB2	1:A:108:ALA:HB3	1.84	0.60
1:B:100:LEU:HB2	1:B:108:ALA:HB3	1.82	0.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/294 (96%)	271 (96%)	12 (4%)	0	100	100
1	B	283/294 (96%)	265 (94%)	18 (6%)	0	100	100
All	All	566/588 (96%)	536 (95%)	30 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/264 (95%)	250 (100%)	0	100	100
1	B	252/264 (96%)	252 (100%)	0	100	100
All	All	502/528 (95%)	502 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	3DR	F	20	3	8,11,12	2.32	4 (50%)	9,14,17	1.56	3 (33%)
3	3DR	D	20	3	8,11,12	2.36	4 (50%)	9,14,17	1.57	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3DR	F	20	3	-	1/3/15/16	0/1/1/1
3	3DR	D	20	3	-	2/3/15/16	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	20	3DR	C2'-C3'	-4.60	1.44	1.52
3	F	20	3DR	C2'-C3'	-4.52	1.44	1.52
3	F	20	3DR	C3'-C4'	-2.64	1.45	1.53
3	D	20	3DR	C3'-C4'	-2.61	1.45	1.53
3	D	20	3DR	O5'-C5'	-2.40	1.38	1.44

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	20	3DR	O4'-C4'-C3'	2.87	107.95	103.73
3	F	20	3DR	O4'-C4'-C3'	2.79	107.84	103.73
3	D	20	3DR	C2'-C3'-C4'	2.53	108.00	102.75
3	F	20	3DR	C2'-C3'-C4'	2.50	107.93	102.75
3	F	20	3DR	C1'-O4'-C4'	-2.09	105.11	108.48

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	20	3DR	O4'-C4'-C5'-O5'
3	F	20	3DR	O4'-C4'-C5'-O5'
3	D	20	3DR	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	20	3DR	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/294 (96%)	0.16	3 (1%) 80 78	14, 25, 37, 47	11 (3%)
1	B	285/294 (96%)	0.21	3 (1%) 80 78	14, 27, 38, 44	10 (3%)
2	C	13/13 (100%)	0.40	0 100 100	23, 34, 48, 53	0
2	E	13/13 (100%)	0.62	0 100 100	25, 47, 55, 60	0
3	D	12/13 (92%)	0.68	2 (16%) 1 1	30, 43, 67, 71	0
3	F	12/13 (92%)	0.91	2 (16%) 1 1	33, 48, 84, 94	0
All	All	620/640 (96%)	0.22	10 (1%) 72 68	14, 26, 43, 94	21 (3%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	72	LYS	2.7
3	F	26	DC	2.6
3	D	14	DC	2.6
1	B	246	LYS	2.6
1	B	260	ASN	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	3DR	D	20	11/12	0.95	0.16	12,15,22,22	0
3	3DR	F	20	11/12	0.96	0.15	17,18,21,22	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.