

Full wwPDB X-ray Structure Validation Report (i)

Oct 2, 2023 – 02:57 PM EDT

PDB ID	:	6NU5
Title	:	Pyruvate Kinase M2 Mutant - S437Y in Complex with L-cysteine
Authors	:	Srivastava, D.; Nandi, S.; Dey, M.
Deposited on		
Resolution	:	1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 9057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	522	Total	С	Ν	0	S	47	19	0
	A	322	4007	2538	700	742	27	47	12	0
1	D	518	Total	С	Ν	0	S	0	17	0
1	D	510	3997	2533	696	738	30	0		0

• Molecule 1 is a protein called Pyruvate kinase PKM.

Chain	Residue	Modelled	Actual	Comment	Reference
А	-19	MET	-	initiating methionine	UNP P14618
А	-18	GLY	-	expression tag	UNP P14618
А	-17	SER	-	expression tag	UNP P14618
А	-16	SER	-	expression tag	UNP P14618
А	-15	HIS	-	expression tag	UNP P14618
А	-14	HIS	-	expression tag	UNP P14618
А	-13	HIS	-	expression tag	UNP P14618
А	-12	HIS	-	expression tag	UNP P14618
А	-11	HIS	-	expression tag	UNP P14618
А	-10	HIS	-	expression tag	UNP P14618
А	-9	SER	-	expression tag	UNP P14618
А	-8	SER	-	expression tag	UNP P14618
А	-7	GLY	-	expression tag	UNP P14618
А	-6	LEU	-	expression tag	UNP P14618
A	-5	VAL	-	expression tag	UNP P14618
A	-4	PRO	-	expression tag	UNP P14618
А	-3	ARG	-	expression tag	UNP P14618
А	-2	GLY	-	expression tag	UNP P14618
А	-1	SER	-	expression tag	UNP P14618
А	0	HIS	-	expression tag	UNP P14618
А	437	TYR	SER	engineered mutation	UNP P14618
В	-19	MET	-	initiating methionine	UNP P14618
В	-18	GLY	-	expression tag	UNP P14618
В	-17	SER	-	expression tag	UNP P14618
В	-16	SER	-	expression tag	UNP P14618

There are 42 discrepancies between the modelled and reference sequences:


Continued on next page...



Chain	Residue	Modelled	Actual Comment		Reference
В	-15	HIS	-	expression tag	UNP P14618
В	-14	HIS	-	expression tag	UNP P14618
В	-13	HIS	-	expression tag	UNP P14618
В	-12	HIS	-	expression tag	UNP P14618
В	-11	HIS	-	expression tag	UNP P14618
В	-10	HIS	-	expression tag	UNP P14618
В	-9	SER	-	expression tag	UNP P14618
В	-8	SER	-	expression tag	UNP P14618
В	-7	GLY	-	expression tag	UNP P14618
В	-6	LEU	-	expression tag	UNP P14618
В	-5	VAL	-	expression tag	UNP P14618
В	-4	PRO	-	expression tag	UNP P14618
В	-3	ARG	-	expression tag	UNP P14618
В	-2	GLY	-	expression tag	UNP P14618
В	-1	SER	_	expression tag	UNP P14618
В	0	HIS	-	expression tag	UNP P14618
В	437	TYR	SER	engineered mutation	UNP P14618


Continued from previous page...

• Molecule 2 is CYSTEINE (three-letter code: CYS) (formula: $C_3H_7NO_2S$).




Mo	Chain	Residues		Ato	oms		ZeroOcc	AltConf
2	А	1	Total 7				0	0
2	В	1	Total 7		N 1		0	0





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 2 4 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 2 4 \end{array}$	0	0


• Molecule 4 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{S} \\ 3 & 1 & 1 & 1 \end{array}$	0	0
4	В	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{S} \\ 3 & 1 & 1 & 1 \end{array}$	0	0



• Molecule 5 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYL AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total C N O 19 11 2 6	0	0
5	В	1	Total C N O 19 11 2 6	0	0

• Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mg 1 1	0	0
6	В	1	Total Mg 1 1	0	0

• Molecule 7 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	Total K 1 1	0	0
7	В	1	Total K 1 1	0	0

• Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
8	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	478	Total O 478 478	0	0
9	В	493	Total O 493 493	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	110.00Å 94.13Å 109.26Å	Depositor	
a, b, c, α , β , γ	90.00° 95.61° 90.00°	Depositor	
Resolution (Å)	41.98 - 1.60	Depositor	
% Data completeness	99.5 (41.98-1.60)	Depositor	
(in resolution range)		-	
R _{merge}	0.05	Depositor	
R _{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$2.55 (at 1.60 \text{\AA})$	Xtriage	
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor	
R, R_{free}	0.166 , 0.182	Depositor	
Wilson B-factor $(Å^2)$	14.6	Xtriage	
Anisotropy	0.450	Xtriage	
L-test for twinning ²	$ < L >=0.49, < L^2>=0.33$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	9057	wwPDB-VP	
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	Mol Type Chai		ain Res	Link	Bo	Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	SCN	А	603	-	1,2,2	0.45	0	$0,\!1,\!1$	-	-	
5	B3P	В	604	-	18,18,18	0.42	0	$21,\!23,\!23$	1.30	2(9%)	
3	OXL	В	602	6	$5,\!5,\!5$	1.29	0	$6,\!6,\!6$	1.61	2 (33%)	
2	CYS	В	601	-	$5,\!6,\!6$	0.98	0	5,7,7	1.29	0	
5	B3P	А	604	-	18,18,18	0.47	0	21,23,23	1.37	3 (14%)	
8	EDO	В	605	-	3,3,3	0.30	0	2,2,2	1.14	0	
2	CYS	А	601	-	$5,\!6,\!6$	0.91	0	5,7,7	1.50	1 (20%)	
3	OXL	А	602	6	$5,\!5,\!5$	1.28	0	$6,\!6,\!6$	1.56	2 (33%)	
4	SCN	В	603	-	1,2,2	0.38	0	$0,\!1,\!1$	-	-	
8	EDO	В	606	-	$3,\!3,\!3$	0.31	0	$2,\!2,\!2$	0.68	0	

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	B3P	В	604	-	-	4/28/28/28	-
5	B3P	А	604	-	-	6/28/28/28	-
2	CYS	В	601	-	-	0/6/6/6	-
3	OXL	В	602	6	-	0/4/4/4	-
8	EDO	В	605	-	-	1/1/1/1	-
2	CYS	А	601	-	-	0/6/6/6	-
3	OXL	А	602	6	-	0/4/4/4	-
8	EDO	В	606	-	-	0/1/1/1	-

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	604	B3P	C2-N2-C8	-3.43	111.21	116.08
5	В	604	B3P	C2-N2-C8	-3.43	111.22	116.08
5	А	604	B3P	C3-N1-C4	-3.03	111.78	116.08
5	В	604	B3P	C3-N1-C4	-2.79	112.11	116.08
3	А	602	OXL	O3-C1-C2	2.66	121.05	113.16
3	В	602	OXL	O4-C2-C1	2.65	121.05	113.16

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	601	CYS	OXT-C-O	-2.53	118.34	124.09
3	В	602	OXL	O3-C1-C2	2.19	119.68	113.16
3	А	602	OXL	O4-C2-C1	2.08	119.33	113.16
5	А	604	B3P	O3-C11-C8	-2.05	107.48	111.63

Continued from previous page...

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
5	А	604	B3P	C5-C4-N1-C3
5	А	604	B3P	C6-C4-N1-C3
5	А	604	B3P	C7-C4-N1-C3
5	А	604	B3P	O2-C10-C8-C11
5	А	604	B3P	O2-C10-C8-C9
8	В	605	EDO	O1-C1-C2-O2
5	В	604	B3P	C6-C4-N1-C3
5	В	604	B3P	C9-C8-N2-C2
5	В	604	B3P	C10-C8-N2-C2
5	А	604	B3P	O2-C10-C8-N2
5	В	604	B3P	C7-C4-N1-C3


All (11) torsion outliers are listed below:

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

