



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 16, 2021 – 02:06 pm BST

PDB ID : 7O1K
Title : Structure of Mycobacterium tuberculosis beta-oxidation trifunctional enzyme alpha-E141A, beta-C92A mutant
Authors : Dalwani, S.; Wierenga, R.K.; Venkatesan, R.
Deposited on : 2021-03-29
Resolution : 2.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 17030 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxyacyl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	730	5411	3424	934	1033	20	0	0	0
1	B	724	5362	3396	925	1020	21	0	3	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	initiating methionine	UNP O53872
A	-14	GLY	-	expression tag	UNP O53872
A	-13	SER	-	expression tag	UNP O53872
A	-12	SER	-	expression tag	UNP O53872
A	-11	HIS	-	expression tag	UNP O53872
A	-10	HIS	-	expression tag	UNP O53872
A	-9	HIS	-	expression tag	UNP O53872
A	-8	HIS	-	expression tag	UNP O53872
A	-7	HIS	-	expression tag	UNP O53872
A	-6	HIS	-	expression tag	UNP O53872
A	-5	SER	-	expression tag	UNP O53872
A	-4	GLN	-	expression tag	UNP O53872
A	-3	ASP	-	expression tag	UNP O53872
A	-2	PRO	-	expression tag	UNP O53872
A	-1	ASN	-	expression tag	UNP O53872
A	0	SER	-	expression tag	UNP O53872
A	141	ALA	GLU	engineered mutation	UNP O53872
B	-15	MET	-	initiating methionine	UNP O53872
B	-14	GLY	-	expression tag	UNP O53872
B	-13	SER	-	expression tag	UNP O53872
B	-12	SER	-	expression tag	UNP O53872
B	-11	HIS	-	expression tag	UNP O53872
B	-10	HIS	-	expression tag	UNP O53872
B	-9	HIS	-	expression tag	UNP O53872
B	-8	HIS	-	expression tag	UNP O53872

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	HIS	-	expression tag	UNP O53872
B	-6	HIS	-	expression tag	UNP O53872
B	-5	SER	-	expression tag	UNP O53872
B	-4	GLN	-	expression tag	UNP O53872
B	-3	ASP	-	expression tag	UNP O53872
B	-2	PRO	-	expression tag	UNP O53872
B	-1	ASN	-	expression tag	UNP O53872
B	0	SER	-	expression tag	UNP O53872
B	141	ALA	GLU	engineered mutation	UNP O53872


- Molecule 2 is a protein called Putative acyltransferase Rv0859.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	402	2967	1853	528	572	14	0	2	0
2	D	402	2964	1852	525	573	14	0	3	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	92	ALA	CYS	engineered mutation	UNP O53871
D	92	ALA	CYS	engineered mutation	UNP O53871

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	36	Total	O	0	0
			36	36		
5	B	46	Total	O	0	0
			46	46		
5	C	30	Total	O	0	0
			30	30		
5	D	26	Total	O	0	0
			26	26		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: 3-hydroxyacyl-CoA dehydrogenase




- Molecule 1: 3-hydroxyacyl-CoA dehydrogenase



- Molecule 2: Putative acyltransferase Rv0859



- Molecule 2: Putative acyltransferase Rv0859





4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	248.94Å 134.96Å 118.93Å 90.00° 110.41° 90.00°	Depositor
Resolution (Å)	48.38 – 2.86	Depositor
% Data completeness (in resolution range)	97.9 (48.38-2.86)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.185 , 0.225	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	17030	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/5515	0.43	0/7471
1	B	0.27	0/5471	0.43	0/7408
2	C	0.27	0/3017	0.46	0/4086
2	D	0.26	0/3017	0.45	0/4088
All	All	0.26	0/17020	0.44	0/23053

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5411	0	5420	24	0
1	B	5362	0	5387	28	0
2	C	2967	0	2989	23	0
2	D	2964	0	2986	26	0
3	A	18	0	24	0	0
3	B	12	0	16	0	0
3	C	12	0	16	0	0
3	D	6	0	8	1	0
4	A	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	40	0	0	1	0
4	C	45	0	0	0	0
4	D	25	0	0	0	0
5	A	36	0	0	1	0
5	B	46	0	0	1	0
5	C	30	0	0	0	0
5	D	26	0	0	0	0
All	All	17030	0	16846	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:296:PRO:HD3	2:D:81:VAL:HG21	1.75	0.69
1:B:250:ARG:NH1	2:C:142:TYR:O	2.26	0.68
2:D:252:HIS:HE1	2:D:332:SER:H	1.42	0.67
2:D:338:GLN:NE2	2:D:345:ASP:OD1	2.29	0.66
2:C:338:GLN:NE2	2:C:345:ASP:OD1	2.27	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/736 (99%)	704 (97%)	22 (3%)	0	100	100
1	B	721/736 (98%)	701 (97%)	18 (2%)	2 (0%)	41	68
2	C	402/403 (100%)	386 (96%)	16 (4%)	0	100	100
2	D	403/403 (100%)	389 (96%)	14 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2252/2278 (99%)	2180 (97%)	70 (3%)	2 (0%)	100	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	556[A]	ASN
1	B	556[B]	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	552/565 (98%)	551 (100%)	1 (0%)	93	98
1	B	548/565 (97%)	545 (100%)	3 (0%)	88	96
2	C	307/309 (99%)	307 (100%)	0	100	100
2	D	308/309 (100%)	302 (98%)	6 (2%)	57	81
All	All	1715/1748 (98%)	1705 (99%)	10 (1%)	86	95

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	235	ASP
2	D	236	ASP
2	D	299	MET
1	B	568	ARG
2	D	81	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

36 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	C	509	-	4,4,4	0.13	0	6,6,6	0.08	0
4	SO4	B	803	-	4,4,4	0.14	0	6,6,6	0.07	0
3	GOL	A	802	-	5,5,5	0.91	0	5,5,5	1.00	0
4	SO4	B	809	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	B	808	-	4,4,4	0.14	0	6,6,6	0.08	0
3	GOL	B	801	-	5,5,5	0.90	0	5,5,5	0.99	0
4	SO4	C	507	-	4,4,4	0.15	0	6,6,6	0.08	0
4	SO4	B	810	-	4,4,4	0.13	0	6,6,6	0.08	0
4	SO4	C	511	-	4,4,4	0.14	0	6,6,6	0.08	0
4	SO4	A	805	-	4,4,4	0.13	0	6,6,6	0.09	0
4	SO4	C	505	-	4,4,4	0.13	0	6,6,6	0.10	0
4	SO4	D	503	-	4,4,4	0.14	0	6,6,6	0.10	0
4	SO4	A	804	-	4,4,4	0.14	0	6,6,6	0.10	0
3	GOL	D	501	-	5,5,5	0.98	0	5,5,5	0.97	0
4	SO4	C	510	-	4,4,4	0.13	0	6,6,6	0.09	0
4	SO4	C	508	-	4,4,4	0.14	0	6,6,6	0.05	0
3	GOL	C	502	-	5,5,5	1.04	0	5,5,5	0.96	0
4	SO4	C	506	-	4,4,4	0.12	0	6,6,6	0.15	0
4	SO4	D	505	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	B	806	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	B	807	-	4,4,4	0.15	0	6,6,6	0.04	0
4	SO4	C	504	-	4,4,4	0.13	0	6,6,6	0.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	809	-	4,4,4	0.15	0	6,6,6	0.06	0
3	GOL	B	802	-	5,5,5	0.90	0	5,5,5	0.99	0
4	SO4	B	805	-	4,4,4	0.14	0	6,6,6	0.07	0
3	GOL	A	803	-	5,5,5	0.95	0	5,5,5	0.90	0
4	SO4	A	808	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	D	504	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	B	804	-	4,4,4	0.14	0	6,6,6	0.06	0
3	GOL	C	501	-	5,5,5	0.94	0	5,5,5	0.91	0
4	SO4	D	502	-	4,4,4	0.14	0	6,6,6	0.06	0
3	GOL	A	801	-	5,5,5	0.95	0	5,5,5	0.98	0
4	SO4	C	503	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	D	506	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	A	807	-	4,4,4	0.13	0	6,6,6	0.09	0
4	SO4	A	806	-	4,4,4	0.15	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	501	-	-	0/4/4/4	-
3	GOL	A	801	-	-	2/4/4/4	-
3	GOL	B	802	-	-	0/4/4/4	-
3	GOL	A	803	-	-	1/4/4/4	-
3	GOL	A	802	-	-	0/4/4/4	-
3	GOL	D	501	-	-	0/4/4/4	-
3	GOL	B	801	-	-	1/4/4/4	-
3	GOL	C	502	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	GOL	O1-C1-C2-O2
3	B	801	GOL	O2-C2-C3-O3
3	A	801	GOL	O1-C1-C2-C3
3	A	803	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	809	SO4	1	0
3	D	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.