

Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 – 04:19 pm BST

PDB ID : 3O2C

Title : Crystal structure of a rod form of c-phycocyanin from Themosynechococcus

vulcanus at 1.5 angstroms

Authors : David, L.; Marx, A.; Adir, N.

Deposited on : 2010-07-22

Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4: 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	162	86%	13%	•				
2	В	172	91%	9%	•				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MEN	В	72	_	X	_	_



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3093 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called C-phycocyanin alpha subunit.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	162	Total	С	N	О	S	0	0	0
1	A	102	1226	772	205	242	7	0	U	U

• Molecule 2 is a protein called C-phycocyanin beta subunit.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	172	Total 1279	C 792	N 229	O 250	S 8	0	1	0

• Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: C₃₃H₄₀N₄O₆).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 43				0	0
3	В	1	Total 43	C 33			0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	В	1	Total 43	C 33	N 4	O 6	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	245	Total O 245 245	0	0
4	В	214	Total O 214 214	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: C-phycocyanin alpha subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	187.47Å 187.47Å 60.18Å	Danagitan
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.00 - 1.50	Depositor
Resolution (A)	26.00 - 1.50	EDS
% Data completeness	98.0 (26.00-1.50)	Depositor
(in resolution range)	98.0 (26.00-1.50)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.02	Depositor
$< I/\sigma(I) > 1$	12.46 (at 1.50Å)	Xtriage
Refinement program	CNS 1.1	Depositor
D D.	0.239 , 0.263	Depositor
R, R_{free}	0.233 , 0.259	DCC
R_{free} test set	3194 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	12.8	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38 , 45.8	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3093	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, MEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.77	$2/1248 \ (0.2\%)$	0.60	0/1692	
2	В	0.71	0/1286	0.66	$2/1740 \ (0.1\%)$	
All	All	0.74	2/2534~(0.1%)	0.63	2/3432 (0.1%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
1	A	74	TYR	CE2-CZ	-5.58	1.31	1.38
1	A	32	LYS	CE-NZ	-5.51	1.35	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	В	79	ARG	NE-CZ-NH2	8.71	124.65	120.30
2	В	79	ARG	NE-CZ-NH1	-8.58	116.01	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1226	0	1204	20	0
2	В	1279	0	1282	16	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	43	0	37	4	0
3	В	86	0	76	11	0
4	A	245	0	0	2	0
4	В	214	0	0	2	0
All	All	3093	0	2599	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom-2	Interatomic	Clash	
Atom-1	Atom-2	${f distance} \; ({f \AA})$	overlap (Å)	
2:B:155:CYS:SG	3:B:555:CYC:HAC1	1.27	1.68	
1:A:65:TYR:HD2	1:A:69:MET:SD	1.95	0.89	
2:B:79:ARG:NH2	3:B:584:CYC:O2D	2.11	0.83	
2:B:155:CYS:SG	3:B:555:CYC:HAC2	2.18	0.82	
1:A:65:TYR:CD2	1:A:69:MET:SD	2.73	0.81	
3:B:555:CYC:O2D	3:B:555:CYC:HHA	1.82	0.78	
1:A:38:MET:O	1:A:42:ARG:HG3	1.95	0.67	
3:B:584:CYC:HC	3:B:584:CYC:HMD2	1.59	0.67	
2:B:72:MEN:HE22	3:B:584:CYC:HBD2	1.76	0.66	
1:A:33:ARG:HH12	1:A:147:GLN:HE21	1.42	0.66	
1:A:10:ALA:O	1:A:14:THR:HG23	1.96	0.65	
3:A:484:CYC:HMD2	3:A:484:CYC:HC	1.65	0.62	
1:A:21:ASN:HB2	4:A:359:HOH:O	2.01	0.61	
2:B:7:LYS:O	2:B:11:GLN:HG3	2.04	0.57	
2:B:115:LEU:HD13	3:B:584:CYC:HMB3	1.86	0.56	
1:A:73:GLN:NE2	1:A:74:TYR:CE2	2.74	0.55	
2:B:116:ARG:O	2:B:120:GLN:HG3	2.09	0.53	
3:A:484:CYC:NB	3:A:484:CYC:HMA1	2.25	0.52	
1:A:73:GLN:HA	3:A:484:CYC:HBD2	1.92	0.51	
2:B:29:ASN:HA	2:B:32:LYS:HE2	1.93	0.51	
1:A:42:ARG:NH1	2:B:24:PHE:HB3	2.27	0.49	
3:B:555:CYC:O2D	3:B:555:CYC:CHA	2.59	0.47	
2:B:57:ARG:HD2	4:B:213:HOH:O	2.14	0.47	
2:B:150:ILE:HG21	3:B:555:CYC:HMC3	1.96	0.47	
1:A:28:ASP:OD2	1:A:32:LYS:HE2	2.15	0.47	
2:B:28:SER:O	2:B:32:LYS:HG3	2.15	0.47	
2:B:72:MEN:CE2	3:B:584:CYC:HBD2	2.43	0.47	
1:A:33:ARG:HH12	1:A:147:GLN:NE2	2.12	0.46	
1:A:65:TYR:HB2	1:A:69:MET:SD	2.56	0.46	



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance} \; ({f \AA})$	$overlap(ext{\AA})$
2:B:1:MET:HB3	4:B:412:HOH:O	2.18	0.43
3:A:484:CYC:HB	3:A:484:CYC:HMA1	1.83	0.43
1:A:33:ARG:NH1	1:A:147:GLN:HE21	2.14	0.42
2:B:155:CYS:SG	3:B:555:CYC:C3C	2.99	0.42
1:A:57:GLN:O	1:A:61:GLN:HG3	2.19	0.42
1:A:11:ALA:O	1:A:15:GLN:HG3	2.20	0.42
1:A:77:THR:O	1:A:78:PRO:C	2.58	0.41
1:A:15:GLN:HB2	1:A:17:ARG:HG2	2.02	0.41
1:A:171:ASN:ND2	4:A:441:HOH:O	2.53	0.41
1:A:30:ARG:O	1:A:30:ARG:HD2	2.21	0.41
1:A:77:THR:O	1:A:80:GLY:N	2.54	0.41
2:B:15:ARG:HE	2:B:15:ARG:HB2	1.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Favoured Allowed		Outliers Percenti	
1	A	160/162~(99%)	157 (98%)	3 (2%)	0	100	100
2	В	170/172 (99%)	167 (98%)	3 (2%)	0	100	100
All	All	330/334 (99%)	324 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed.	and	the	total	number	of	residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers		Percentiles		
1	A	123/123 (100%)	123 (100%)	0	100	100		
2	В	128/127 (101%)	128 (100%)	0	100	100		
All	All	$251/250 \; (100\%)$	251 (100%)	0	100	100		

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	47	ASN
1	A	57	GLN
1	A	147	GLN
1	A	171	ASN
2	В	35	ASN
2	В	54	ASN
2	В	78	ASN
2	В	145	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Iol Type Chain Res		Chain Res		Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MEN	В	72	2	7,8,9	4.62	4 (57%)	6,9,11	11.46	5 (83%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	В	72	2	-	6/7/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\mathring{\mathrm{A}})$
2	В	72	MEN	OD1-CG	9.73	1.43	1.23
2	В	72	MEN	CG-ND2	5.60	1.60	1.34
2	В	72	MEN	CE2-ND2	3.85	1.52	1.45
2	В	72	MEN	CB-CA	2.76	1.59	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	В	72	MEN	OD1-CG-CB	-24.73	85.27	121.50
2	В	72	MEN	CB-CG-ND2	12.03	131.66	115.48
2	В	72	MEN	OD1-CG-ND2	3.97	143.02	122.36
2	В	72	MEN	CE2-ND2-CG	-3.01	98.99	121.93
2	В	72	MEN	CB-CA-C	-2.55	106.69	111.47

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	72	MEN	C-CA-CB-CG
2	В	72	MEN	OD1-CG-ND2-CE2
2	В	72	MEN	N-CA-CB-CG
2	В	72	MEN	CA-CB-CG-OD1
2	В	72	MEN	CA-CB-CG-ND2
2	В	72	MEN	CB-CG-ND2-CE2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

\mathbf{Mol}	Chain	${f Res}$	Type	Clashes	Symm-Clashes
2	В	72	MEN	2	0



5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	Bond lengths			Bond angles		
10101					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	CYC	A	484	1	36,46,46	2.25	12 (33%)	44,67,67	3.41	14 (31%)	
3	CYC	В	555	2	36,46,46	2.21	11 (30%)	44,67,67	3.70	16 (36%)	
3	CYC	В	584	2	36,46,46	2.36	13 (36%)	44,67,67	3.59	16 (36%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	A	484	1	-	7/21/74/74	0/4/4/4
3	CYC	В	555	2	-	5/21/74/74	0/4/4/4
3	CYC	В	584	2	-	6/21/74/74	0/4/4/4

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
3	A	484	CYC	OB-C4B	7.14	1.37	1.23
3	В	584	CYC	OB-C4B	7.05	1.37	1.23
3	В	555	CYC	OB-C4B	6.71	1.36	1.23
3	В	555	CYC	C1C-NC	-5.14	1.30	1.37
3	В	555	CYC	OC-C1C	4.76	1.32	1.23
3	В	584	CYC	OC-C1C	4.70	1.32	1.23
3	В	584	CYC	C1C-NC	-4.68	1.31	1.37
3	A	484	CYC	C1C-NC	-4.54	1.31	1.37



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	${ m Observed}(m \AA)$	Ideal(A)
3	Α	484	CYC	OC-C1C	4.52	1.32	1.23
3	В	584	CYC	CHB-C4A	3.98	1.49	1.40
3	В	584	CYC	C2A-C3A	3.90	1.45	1.36
3	В	555	CYC	CHB-C4A	3.48	1.48	1.40
3	A	484	CYC	C3D-C2D	3.39	1.47	1.37
3	В	584	CYC	C3D-C2D	3.38	1.47	1.37
3	A	484	CYC	CHB-C4A	3.37	1.48	1.40
3	В	555	CYC	C2A-C3A	3.34	1.43	1.36
3	A	484	CYC	C2A-C3A	3.32	1.43	1.36
3	В	555	CYC	C3D-C2D	3.23	1.47	1.37
3	В	555	CYC	C4B-C3B	-2.81	1.42	1.48
3	В	584	CYC	C1B-NB	-2.65	1.33	1.37
3	В	584	CYC	CHA-C1A	2.64	1.37	1.35
3	В	584	CYC	C3B-C2B	2.59	1.42	1.36
3	В	584	CYC	C4B-C3B	-2.58	1.43	1.48
3	В	584	CYC	CMB-C2B	-2.56	1.45	1.50
3	A	484	CYC	C2C-C1C	2.48	1.54	1.52
3	В	555	CYC	C3B-C2B	2.44	1.41	1.36
3	Α	484	CYC	C2C-C3C	-2.37	1.47	1.54
3	A	484	CYC	C4B-C3B	-2.37	1.43	1.48
3	В	584	CYC	C4B-NB	-2.31	1.33	1.38
3	В	555	CYC	CMB-C2B	-2.27	1.46	1.50
3	A	484	CYC	C4B-NB	-2.25	1.33	1.38
3	В	555	CYC	C1B-NB	-2.18	1.34	1.37
3	A	484	CYC	CMB-C2B	-2.18	1.46	1.50
3	В	555	CYC	C4B-NB	-2.13	1.33	1.38
3	A	484	CYC	C3B-C2B	2.11	1.41	1.36
3	В	584	CYC	C1D-CHD	2.02	1.49	1.41

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	В	555	CYC	C3B-C4B-NB	12.61	116.97	106.78
3	В	584	CYC	C3B-C4B-NB	12.51	116.89	106.78
3	A	484	CYC	C3B-C4B-NB	12.12	116.57	106.78
3	В	584	CYC	OC-C1C-C2C	-11.10	117.35	126.17
3	В	555	CYC	OC-C1C-C2C	-10.28	118.00	126.17
3	A	484	CYC	OC-C1C-C2C	-10.06	118.17	126.17
3	В	584	CYC	C2C-C1C-NC	8.70	115.77	108.27
3	В	555	CYC	C2C-C1C-NC	8.41	115.53	108.27
3	В	555	CYC	OB-C4B-C3B	-8.16	119.18	128.04
3	В	584	CYC	OB-C4B-C3B	-7.82	119.55	128.04



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	A	484	CYC	C2C-C1C-NC	7.82	115.01	108.27
3	A	484	CYC	OB-C4B-C3B	-7.56	119.84	128.04
3	В	555	CYC	CBC-CAC-C3C	5.94	126.69	113.47
3	A	484	CYC	CAB-C3B-C4B	5.24	129.65	121.38
3	В	584	CYC	CAB-C3B-C4B	5.13	129.47	121.38
3	В	555	CYC	C1B-NB-C4B	-4.99	104.31	110.67
3	A	484	CYC	C1B-NB-C4B	-4.96	104.36	110.67
3	В	555	CYC	CAB-C3B-C4B	4.59	128.62	121.38
3	В	555	CYC	CAD-CBD-CGD	-4.56	105.03	112.67
3	В	584	CYC	C1B-NB-C4B	-4.48	104.96	110.67
3	В	555	CYC	C4D-CHA-C1A	3.90	133.46	128.81
3	A	484	CYC	C1A-C2A-C3A	-3.27	103.17	106.78
3	В	555	CYC	C1A-C2A-C3A	-3.23	103.21	106.78
3	В	584	CYC	C4D-CHA-C1A	3.18	132.60	128.81
3	В	584	CYC	C1A-C2A-C3A	-3.16	103.28	106.78
3	A	484	CYC	CAA-CBA-CGA	-3.01	107.62	112.67
3	В	584	CYC	CBD-CAD-C3D	2.81	117.67	112.49
3	В	555	CYC	CHB-C4A-C3A	2.78	132.05	124.90
3	В	584	CYC	C1A-NA-C4A	2.75	111.70	106.51
3	В	555	CYC	CHD-C4C-NC	-2.69	122.01	125.20
3	В	584	CYC	CHD-C4C-NC	-2.58	122.13	125.20
3	В	555	CYC	C1A-NA-C4A	2.56	111.33	106.51
3	В	555	CYC	C2B-C1B-NB	2.53	110.69	106.99
3	В	584	CYC	CHB-C4A-C3A	2.49	131.31	124.90
3	В	584	CYC	C4A-C3A-C2A	2.49	109.37	106.51
3	A	484	CYC	C1A-NA-C4A	2.47	111.17	106.51
3	A	484	CYC	CAC-C3C-C2C	2.45	120.39	114.26
3	A	484	CYC	C2B-C1B-NB	2.45	110.58	106.99
3	A	484	CYC	CHB-C4A-C3A	2.37	130.99	124.90
3	A	484	CYC	CHD-C4C-NC	-2.36	122.39	125.20
3	В	584	CYC	CAA-CBA-CGA	-2.33	108.77	112.67
3	В	584	CYC	C2B-C1B-NB	2.32	110.38	106.99
3	В	555	CYC	CAB-C3B-C2B	2.29	131.45	127.53
3	В	555	CYC	C4A-C3A-C2A	2.29	109.14	106.51
3	A	484	CYC	C4A-C3A-C2A	2.29	109.13	106.51
3	В	584	CYC	C3A-C4A-NA	-2.05	106.15	110.53

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	484	CYC	NA-C4A-CHB-C1B



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	484	CYC	C3A-C4A-CHB-C1B
3	A	484	CYC	C4C-C3C-CAC-CBC
3	A	484	CYC	ND-C1D-CHD-C4C
3	A	484	CYC	C2D-C1D-CHD-C4C
3	В	555	CYC	NA-C4A-CHB-C1B
3	В	555	CYC	C3A-C4A-CHB-C1B
3	В	555	CYC	C2C-C3C-CAC-CBC
3	В	555	CYC	C4C-C3C-CAC-CBC
3	В	555	CYC	C3D-CAD-CBD-CGD
3	В	584	CYC	NA-C4A-CHB-C1B
3	В	584	CYC	C3A-C4A-CHB-C1B
3	В	584	CYC	ND-C1D-CHD-C4C
3	В	584	CYC	C2D-C1D-CHD-C4C
3	В	584	CYC	C2B-C3B-CAB-CBB
3	В	584	CYC	C4B-C3B-CAB-CBB
3	A	484	CYC	C2B-C3B-CAB-CBB
3	A	484	CYC	C2C-C3C-CAC-CBC


There are no ring outliers.


3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	484	CYC	4	0
3	В	555	CYC	6	0
3	В	584	CYC	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$162/162 \; (100\%)$	0.30	6 (3%) 41 46	10, 15, 24, 33	0
2	В	171/172 (99%)	0.60	11 (6%) 19 20	12, 20, 32, 44	0
All	All	333/334 (99%)	0.45	17 (5%) 28 30	10, 17, 30, 44	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	174	ALA	4.7
2	В	114	GLY	4.1
1	A	77	THR	3.4
2	В	21	ASN	3.3
2	В	113[A]	ASN	3.1
2	В	148	ASN	2.9
2	В	32	LYS	2.7
1	A	69	MET	2.7
2	В	173	VAL	2.5
1	A	70	GLN	2.5
2	В	29	ASN	2.4
2	В	22	ALA	2.3
1	A	15	GLN	2.3
1	A	14	THR	2.1
2	В	2	LEU	2.1
2	В	15	ARG	2.0
1	A	65	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	MEN	В	72	9/10	0.92	0.12	19,19,31,31	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
3	CYC	В	555	43/43	0.79	0.18	16,20,36,44	0
3	CYC	В	584	43/43	0.81	0.17	13,25,33,35	0
3	CYC	A	484	43/43	0.95	0.11	8,12,16,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers (i)

There are no such residues in this entry.

