



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 12:32 AM EDT

PDB ID : 3O31  
Title : E81Q mutant of MtNAS in complex with a reaction intermediate  
Authors : Dreyfus, C.; Pignol, D.; Arnoux, P.  
Deposited on : 2010-07-23  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	
1	B	296	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ThermoNicotianamine Synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2111	1347	360	393	11	0	0	0
1	B	264	2111	1347	360	393	11	0	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	81	GLN	GLU	ENGINEERED MUTATION	UNP O26771
A	266	LYS	-	EXPRESSION TAG	UNP O26771
A	267	GLY	-	EXPRESSION TAG	UNP O26771
A	268	GLU	-	EXPRESSION TAG	UNP O26771
A	269	LEU	-	EXPRESSION TAG	UNP O26771
A	270	ASN	-	EXPRESSION TAG	UNP O26771
A	271	SER	-	EXPRESSION TAG	UNP O26771
A	272	LYS	-	EXPRESSION TAG	UNP O26771
A	273	LEU	-	EXPRESSION TAG	UNP O26771
A	274	GLU	-	EXPRESSION TAG	UNP O26771
A	275	GLY	-	EXPRESSION TAG	UNP O26771
A	276	LYS	-	EXPRESSION TAG	UNP O26771
A	277	PRO	-	EXPRESSION TAG	UNP O26771
A	278	ILE	-	EXPRESSION TAG	UNP O26771
A	279	PRO	-	EXPRESSION TAG	UNP O26771
A	280	ASN	-	EXPRESSION TAG	UNP O26771
A	281	PRO	-	EXPRESSION TAG	UNP O26771
A	282	LEU	-	EXPRESSION TAG	UNP O26771
A	283	LEU	-	EXPRESSION TAG	UNP O26771
A	284	GLY	-	EXPRESSION TAG	UNP O26771
A	285	LEU	-	EXPRESSION TAG	UNP O26771
A	286	ASP	-	EXPRESSION TAG	UNP O26771
A	287	SER	-	EXPRESSION TAG	UNP O26771
A	288	THR	-	EXPRESSION TAG	UNP O26771
A	289	ARG	-	EXPRESSION TAG	UNP O26771

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	THR	-	EXPRESSION TAG	UNP O26771
A	291	GLY	-	EXPRESSION TAG	UNP O26771
A	292	HIS	-	EXPRESSION TAG	UNP O26771
A	293	HIS	-	EXPRESSION TAG	UNP O26771
A	294	HIS	-	EXPRESSION TAG	UNP O26771
A	295	HIS	-	EXPRESSION TAG	UNP O26771
A	296	HIS	-	EXPRESSION TAG	UNP O26771
A	297	HIS	-	EXPRESSION TAG	UNP O26771
B	81	GLN	GLU	ENGINEERED MUTATION	UNP O26771
B	266	LYS	-	EXPRESSION TAG	UNP O26771
B	267	GLY	-	EXPRESSION TAG	UNP O26771
B	268	GLU	-	EXPRESSION TAG	UNP O26771
B	269	LEU	-	EXPRESSION TAG	UNP O26771
B	270	ASN	-	EXPRESSION TAG	UNP O26771
B	271	SER	-	EXPRESSION TAG	UNP O26771
B	272	LYS	-	EXPRESSION TAG	UNP O26771
B	273	LEU	-	EXPRESSION TAG	UNP O26771
B	274	GLU	-	EXPRESSION TAG	UNP O26771
B	275	GLY	-	EXPRESSION TAG	UNP O26771
B	276	LYS	-	EXPRESSION TAG	UNP O26771
B	277	PRO	-	EXPRESSION TAG	UNP O26771
B	278	ILE	-	EXPRESSION TAG	UNP O26771
B	279	PRO	-	EXPRESSION TAG	UNP O26771
B	280	ASN	-	EXPRESSION TAG	UNP O26771
B	281	PRO	-	EXPRESSION TAG	UNP O26771
B	282	LEU	-	EXPRESSION TAG	UNP O26771
B	283	LEU	-	EXPRESSION TAG	UNP O26771
B	284	GLY	-	EXPRESSION TAG	UNP O26771
B	285	LEU	-	EXPRESSION TAG	UNP O26771
B	286	ASP	-	EXPRESSION TAG	UNP O26771
B	287	SER	-	EXPRESSION TAG	UNP O26771
B	288	THR	-	EXPRESSION TAG	UNP O26771
B	289	ARG	-	EXPRESSION TAG	UNP O26771
B	290	THR	-	EXPRESSION TAG	UNP O26771
B	291	GLY	-	EXPRESSION TAG	UNP O26771
B	292	HIS	-	EXPRESSION TAG	UNP O26771
B	293	HIS	-	EXPRESSION TAG	UNP O26771
B	294	HIS	-	EXPRESSION TAG	UNP O26771
B	295	HIS	-	EXPRESSION TAG	UNP O26771
B	296	HIS	-	EXPRESSION TAG	UNP O26771
B	297	HIS	-	EXPRESSION TAG	UNP O26771

- Molecule 2 is N-[(3S)-3-amino-3-carboxypropyl]-L-glutamic acid (three-letter code: 3O3)





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.37Å 67.90Å 147.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.39 – 1.70 32.39 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.3 (32.39-1.70) 97.3 (32.39-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.188 , 0.233 0.187 , 0.233	Depositor DCC
$R_{free}$ test set	3493 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.5	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 59.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5223	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 3O3, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/2153	0.66	0/2912
1	B	0.62	0/2153	0.71	2/2912 (0.1%)
All	All	0.59	0/4306	0.69	2/5824 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	217	ARG	NE-CZ-NH2	-7.76	116.42	120.30
1	B	217	ARG	NE-CZ-NH1	7.70	124.15	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2111	0	2108	30	0
1	B	2111	0	2108	43	0
2	A	17	0	13	0	0
2	B	17	0	13	0	0
3	A	4	0	0	1	0
3	B	2	0	0	0	0
4	A	461	0	0	18	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	500	0	0	20	0
All	All	5223	0	4242	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:902:HOH:O	1:B:56:ILE:HG21	1.36	1.23
1:A:147:MET:SD	4:A:826:HOH:O	2.03	1.17
1:B:63:ASN:HB2	4:B:488:HOH:O	1.44	1.17
1:B:112:LYS:HE2	4:B:395:HOH:O	1.60	0.98
1:B:27:MET:HB2	1:B:79:LYS:HD2	1.47	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/296 (88%)	260 (99%)	2 (1%)	0	100	100
1	B	262/296 (88%)	257 (98%)	5 (2%)	0	100	100
All	All	524/592 (88%)	517 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/257 (89%)	225 (98%)	4 (2%)	60	46
1	B	229/257 (89%)	221 (96%)	8 (4%)	36	17
All	All	458/514 (89%)	446 (97%)	12 (3%)	46	28

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	121	ARG
1	B	136	LEU
1	B	197	LEU
1	B	161	LEU
1	A	182	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	113	ASN
1	B	175	ASN
1	B	207	ASN
1	A	207	ASN
1	B	64	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	3O3	A	1	-	15,16,16	0.93	0	14,20,20	1.42	2 (14%)
2	3O3	B	1	-	15,16,16	1.05	1 (6%)	14,20,20	1.09	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3O3	A	1	-	-	5/19/19/19	-
2	3O3	B	1	-	-	6/19/19/19	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	3O3	O5-C11	-2.05	1.24	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	3O3	C7-N3-C8	2.74	118.67	113.76
2	A	1	3O3	C10-C9-C8	2.41	117.60	113.16
2	B	1	3O3	C10-C9-C8	2.04	116.93	113.16

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	3O3	C11-C8-N3-C7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	1	3O3	C11-C8-N3-C7
2	B	1	3O3	C9-C8-N3-C7
2	A	1	3O3	O6-C11-C8-C9
2	B	1	3O3	O5-C11-C8-C9

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	264/296 (89%)	-0.07	4 (1%) 73 77	7, 13, 21, 29	0
1	B	264/296 (89%)	-0.11	3 (1%) 80 83	7, 12, 21, 26	0
All	All	528/592 (89%)	-0.09	7 (1%) 77 81	7, 12, 21, 29	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	265	PRO	5.0
1	B	265	PRO	3.6
1	B	119	ARG	2.8
1	A	121	ARG	2.5
1	A	119	ARG	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	3O3	A	1	17/17	0.94	0.09	9,12,16,17	0
2	3O3	B	1	17/17	0.95	0.10	9,11,15,16	0
3	BR	A	300	1/1	0.99	0.06	25,25,25,25	0
3	BR	A	301	1/1	0.99	0.07	25,25,25,25	0
3	BR	B	298	1/1	0.99	0.02	16,16,16,16	0
3	BR	A	299	1/1	1.00	0.01	13,13,13,13	0
3	BR	A	298	1/1	1.00	0.02	16,16,16,16	0
3	BR	B	299	1/1	1.00	0.04	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.