



## wwPDB EM Validation Summary Report ⓘ

Oct 18, 2021 – 10:17 am BST

PDB ID : 7O37  
EMDB ID : EMD-12702  
Title : Murine supercomplex CIII2CIV in the assembled locked conformation  
Authors : Vercellino, I.; Sazanov, L.A.  
Deposited on : 2021-04-01  
Resolution : 3.20 Å (reported)  
Based on initial models : 3L75, 5Z62, 1NTZ, 5IY5

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev97  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	89% 11%
1	L	446	87% 13%
2	B	439	80% 15% .
2	M	439	82% 14% .
3	C	381	86% 14%
3	N	381	89% 10%
4	D	241	87% 13%
4	O	241	91% 9%


Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	196	51% 84% 16%
5	P	196	43% 81% 19%
6	F	110	81% 12% 7%
6	Q	110	77% 15% 8%
7	G	81	80% 11% 9%
7	R	81	77% 12% 11%
8	H	76	7% 83% 7% 11%
8	S	76	9% 79% 11% 11%
9	J	63	73% 22% 5%
9	U	63	75% 21% 5%
10	K	56	21% 88% 5% 7%
10	V	56	20% 79% 16% 5%
11	T	78	9% 79% 21%
12	I	113	19% 78% 19% ..
13	a	514	99% .
14	b	227	5% 98% .
15	c	261	99% .
16	d	147	6% 97% ..
17	e	109	95% 5%
18	f	99	5% 95% ..
19	g	85	24% 86% . 12%
20	h	85	14% 93% 7%
21	i	75	96% .
22	k	56	5% 86% . 12%
23	l	47	98% .

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
24	m	46	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
34	HEA	a	603	X	-	-	-
34	HEA	a	604	X	-	-	-

## 2 Entry composition [i](#)

There are 37 unique types of molecules in this entry. The entry contains 48610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome b-c1 complex subunit 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	446	Total	C	N	O	S	0	0
			3466	2167	611	671	17		
1	L	445	Total	C	N	O	S	0	0
			3460	2163	610	670	17		

- Molecule 2 is a protein called Cytochrome b-c1 complex subunit 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	420	Total	C	N	O	S	0	0
			3154	1980	555	610	9		
2	M	420	Total	C	N	O	S	0	0
			3154	1980	555	610	9		

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	380	Total	C	N	O	S	0	0
			3045	2052	473	499	21		
3	N	380	Total	C	N	O	S	0	0
			3046	2052	473	499	22		

- Molecule 4 is a protein called Cytochrome c1, heme protein, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	241	Total	C	N	O	S	0	0
			1919	1224	329	352	14		
4	O	240	Total	C	N	O	S	0	0
			1909	1218	327	350	14		

- Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	196	Total	C	N	O	S	0	0
			1512	952	263	290	7		
5	P	196	Total	C	N	O	S	0	0
			1512	952	263	290	7		

- Molecule 6 is a protein called Cytochrome b-c1 complex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	102	Total	C	N	O	S	0	0
			900	575	160	162	3		
6	Q	101	Total	C	N	O	S	0	0
			894	572	159	160	3		

- Molecule 7 is a protein called Cytochrome b-c1 complex subunit 8.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	G	74	Total	C	N	O	0	0
			624	404	117	103		
7	R	72	Total	C	N	O	0	0
			609	396	112	101		

- Molecule 8 is a protein called Cytochrome b-c1 complex subunit 6, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	68	Total	C	N	O	S	0	0
			563	343	103	112	5		
8	S	68	Total	C	N	O	S	0	0
			563	343	103	112	5		

- Molecule 9 is a protein called Cytochrome b-c1 complex subunit 9.

Mol	Chain	Residues	Atoms				AltConf	Trace
9	J	60	Total	C	N	O	0	0
			495	323	86	86		
9	U	60	Total	C	N	O	0	0
			495	323	86	86		

- Molecule 10 is a protein called Cytochrome b-c1 complex subunit 10.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	K	52	Total	C	N	O	S	0	0
			429	286	75	66	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
10	V	53	Total	C	N	O	S	0	0
			438	292	77	67	2		

- Molecule 11 is a protein called Cytochrome b-c1 complex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	T	78	Total	C	N	O	S	0	0
			554	352	103	97	2		

- Molecule 12 is a protein called Cox7a2l protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	I	111	Total	C	N	O	S	0	0
			828	539	138	146	5		

- Molecule 13 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	a	514	Total	C	N	O	S	0	0
			4021	2691	623	675	32		

- Molecule 14 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	b	227	Total	C	N	O	S	0	0
			1817	1180	282	336	19		

- Molecule 15 is a protein called Cytochrome c oxidase subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	c	259	Total	C	N	O	S	0	0
			2111	1414	338	349	10		

- Molecule 16 is a protein called Cytochrome c oxidase subunit 4 isoform 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	d	144	Total	C	N	O	S	0	0
			1195	770	199	219	7		

- Molecule 17 is a protein called Cytochrome c oxidase subunit 5A, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	e	104	Total	C	N	O	S	0	0
			842	538	141	161	2		

- Molecule 18 is a protein called Cytochrome c oxidase subunit 5B, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	f	95	Total	C	N	O	S	0	0
			727	452	127	140	8		

- Molecule 19 is a protein called Cytochrome c oxidase subunit 6A2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	g	75	Total	C	N	O	S	0	0
			605	392	114	96	3		

- Molecule 20 is a protein called Cytochrome c oxidase subunit 6B1.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	h	79	Total	C	N	O	S	0	0
			654	416	116	117	5		

- Molecule 21 is a protein called Cytochrome c oxidase subunit 6C.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	i	72	Total	C	N	O	S	0	0
			572	372	103	94	3		

- Molecule 22 is a protein called Cytochrome c oxidase subunit 7B, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	k	49	Total	C	N	O	S	0	0
			383	248	65	68	2		

- Molecule 23 is a protein called Cytochrome c oxidase subunit 7C, mitochondrial.


Mol	Chain	Residues	Atoms					AltConf	Trace
23	l	46	Total	C	N	O	S	0	0
			380	253	64	61	2		

- Molecule 24 is a protein called Cytochrome c oxidase subunit 8B, mitochondrial.



Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	m	43	311	203	51	56	1	0	0

- Molecule 25 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula:  $C_{41}H_{82}NO_8P$ ).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
25	A	1	23	13	1	8	1	0
25	C	1	35	25	1	8	1	0
25	D	1	32	22	1	8	1	0
25	F	1	37	27	1	8	1	0
25	G	1	42	32	1	8	1	0
25	L	1	23	13	1	8	1	0
25	N	1	37	27	1	8	1	0
25	O	1	23	13	1	8	1	0
25	R	1	51	41	1	8	1	0
25	a	1	62	42	2	16	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
25	a	1	Total 62	C 42	N 2	O 16	P 2	0
25	b	1	Total 57	C 37	N 2	O 16	P 2	0
25	b	1	Total 57	C 37	N 2	O 16	P 2	0
25	c	1	Total 45	C 35	N 1	O 8	P 1	0
25	g	1	Total 25	C 15	N 1	O 8	P 1	0
25	k	1	Total 27	C 17	N 1	O 8	P 1	0

- Molecule 26 is CARDIOLIPIN (three-letter code: CDL) (formula:  $C_{81}H_{156}O_{17}P_2$ ).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
26	A	1	Total 46	C 27	O 17	P 2	0
26	D	1	Total 56	C 37	O 17	P 2	0
26	G	1	Total 42	C 23	O 17	P 2	0
26	L	1	Total 46	C 27	O 17	P 2	0
26	O	1	Total 57	C 38	O 17	P 2	0


Continued on next page...






Mol	Chain	Residues	Atoms				AltConf	
28	D	1	Total	C	Fe	N	O	0
			43	34	1	4	4	
28	O	1	Total	C	Fe	N	O	0
			43	34	1	4	4	

- Molecule 29 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).




Mol	Chain	Residues	Atoms			AltConf
29	E	1	Total	Fe	S	0
			4	2	2	
29	P	1	Total	Fe	S	0
			4	2	2	

- Molecule 30 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PC1) (formula:  $C_{44}H_{88}NO_8P$ ).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
30	E	1	Total	C	N	O	P	0
			35	25	1	8	1	
30	G	1	Total	C	N	O	P	0
			78	58	2	16	2	
30	G	1	Total	C	N	O	P	0
			78	58	2	16	2	
30	U	1	Total	C	N	O	P	0
			24	14	1	8	1	
30	a	1	Total	C	N	O	P	0
			43	33	1	8	1	

- Molecule 31 is TRISTEAROYLGLYCEROL (three-letter code: TGL) (formula:  $C_{57}H_{110}O_6$ ).



Mol	Chain	Residues	Atoms			AltConf
31	G	1	Total	C	O	0
			44	38	6	


- Molecule 32 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		AltConf
32	a	1	Total	Cu	0
			1	1	

- Molecule 33 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		AltConf
33	a	1	Total	Na	0
			1	1	

- Molecule 34 is HEME-A (three-letter code: HEA) (formula: C<sub>49</sub>H<sub>56</sub>FeN<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				AltConf	
			Total	C	Fe	N		O
34	a	1	120	98	2	8	12	0
34	a	1	120	98	2	8	12	0

- Molecule 35 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
35	b	1	1	1	0

- Molecule 36 is DINUCLEAR COPPER ION (three-letter code: CUA) (formula: Cu<sub>2</sub>).



Mol	Chain	Residues	Atoms		AltConf
36	b	1	Total	Cu	0
			2	2	

- Molecule 37 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
37	f	1	Total	Zn	0
			1	1	




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome b-c1 complex subunit 1, mitochondrial





- Molecule 1: Cytochrome b-c1 complex subunit 1, mitochondrial





- Molecule 2: Cytochrome b-c1 complex subunit 2, mitochondrial





- Molecule 2: Cytochrome b-c1 complex subunit 2, mitochondrial





• Molecule 3: Cytochrome b





• Molecule 3: Cytochrome b




• Molecule 4: Cytochrome c1, heme protein, mitochondrial




• Molecule 4: Cytochrome c1, heme protein, mitochondrial




• Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial




• Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial




• Molecule 6: Cytochrome b-c1 complex subunit 7




• Molecule 6: Cytochrome b-c1 complex subunit 7



• Molecule 7: Cytochrome b-c1 complex subunit 8





- Molecule 7: Cytochrome b-c1 complex subunit 8

Chain R:  77% 12% 11%





- Molecule 8: Cytochrome b-c1 complex subunit 6, mitochondrial

Chain H:  7% 83% 7% 11%



- Molecule 8: Cytochrome b-c1 complex subunit 6, mitochondrial

Chain S:  9% 79% 11% 11%



- Molecule 9: Cytochrome b-c1 complex subunit 9

Chain J:  73% 22% 5%





- Molecule 9: Cytochrome b-c1 complex subunit 9

Chain U:  75% 21% 5%




- Molecule 10: Cytochrome b-c1 complex subunit 10

Chain K:  21% 88% 5% 7%





- Molecule 10: Cytochrome b-c1 complex subunit 10



Chain V:  20% 79% 16% 5%




• Molecule 11: Cytochrome b-c1 complex subunit 9




• Molecule 12: Cox7a2l protein




• Molecule 13: Cytochrome c oxidase subunit 1



• Molecule 14: Cytochrome c oxidase subunit 2




• Molecule 15: Cytochrome c oxidase subunit 3




• Molecule 16: Cytochrome c oxidase subunit 4 isoform 1, mitochondrial






- Molecule 17: Cytochrome c oxidase subunit 5A, mitochondrial

Chain e: 95% 5%




- Molecule 18: Cytochrome c oxidase subunit 5B, mitochondrial

Chain f: 5% 95% ..




- Molecule 19: Cytochrome c oxidase subunit 6A2, mitochondrial

Chain g: 24% 86% 12%



- Molecule 20: Cytochrome c oxidase subunit 6B1

Chain h: 14% 93% 7%



- Molecule 21: Cytochrome c oxidase subunit 6C

Chain i: 96%



- Molecule 22: Cytochrome c oxidase subunit 7B, mitochondrial

Chain k: 5% 86% 12%





- Molecule 23: Cytochrome c oxidase subunit 7C, mitochondrial

Chain l:  98%



- Molecule 24: Cytochrome c oxidase subunit 8B, mitochondrial

Chain m:  7% 93% 7%



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided, Not provided, Not provided, C1, Not provided, Not provided, Not provided	Depositor
Number of particles used	14851	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	90.66	Depositor
Minimum defocus (nm)	200	Depositor
Maximum defocus (nm)	2700	Depositor
Magnification	75000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.332	Depositor
Minimum map value	-0.028	Depositor
Average map value	0.007	Depositor
Map value standard deviation	0.021	Depositor
Recommended contour level	0.05	Depositor
Map size ( $\text{\AA}$ )	165.98401, 217.056, 176.62401	wwPDB
Map dimensions	166, 204, 156	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.064, 1.064, 1.064	Depositor



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEA, HEM, MG, FES, PC1, NA, 3PE, ZN, CDL, TGL, CU, HEC, CUA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.30	0/3536	0.58	1/4803 (0.0%)
1	L	0.31	0/3530	0.60	2/4793 (0.0%)
2	B	0.28	0/3205	0.56	3/4332 (0.1%)
2	M	0.29	0/3205	0.53	0/4332
3	C	0.32	0/3147	0.56	1/4299 (0.0%)
3	N	0.32	0/3147	0.57	2/4297 (0.0%)
4	D	0.31	0/1978	0.57	1/2685 (0.0%)
4	O	0.29	0/1968	0.53	0/2674
5	E	0.28	0/1545	0.56	0/2091
5	P	0.27	0/1545	0.58	0/2091
6	F	0.28	0/922	0.55	0/1234
6	Q	0.29	0/916	0.57	0/1226
7	G	0.36	0/642	0.64	0/867
7	R	0.33	0/627	0.67	0/848
8	H	0.42	1/570 (0.2%)	0.71	2/763 (0.3%)
8	S	0.36	0/570	0.70	1/763 (0.1%)
9	J	0.29	0/509	0.57	0/687
9	U	0.30	0/509	0.64	0/687
10	K	0.27	0/445	0.62	0/608
10	V	0.29	0/454	0.69	1/619 (0.2%)
11	T	0.32	0/565	0.68	0/772
12	I	0.32	0/849	0.70	3/1153 (0.3%)
13	a	0.37	0/4162	0.63	3/5686 (0.1%)
14	b	0.37	0/1863	0.71	5/2542 (0.2%)
15	c	0.35	0/2195	0.59	0/3000
16	d	0.34	0/1229	0.61	1/1659 (0.1%)
17	e	0.34	0/860	0.67	0/1167
18	f	0.36	0/744	0.69	0/1009
19	g	0.28	0/632	0.61	1/866 (0.1%)
20	h	0.34	0/674	0.67	0/910
21	i	0.35	0/584	0.67	0/778
22	k	0.34	0/396	0.66	1/541 (0.2%)

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
23	l	0.37	0/393	0.55	0/527
24	m	0.35	0/318	0.60	0/433
All	All	0.32	1/48434 (0.0%)	0.60	28/65742 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
18	f	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	H	66	CYS	CB-SG	-5.48	1.72	1.81

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	316	ASP	CB-CG-OD1	11.46	128.62	118.30
14	b	68	LEU	CA-CB-CG	7.75	133.13	115.30
3	C	58	ASP	CB-CG-OD1	7.07	124.66	118.30
14	b	45	MET	CA-CB-CG	7.01	125.21	113.30
13	a	51	ASP	CB-CG-OD1	6.78	124.41	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
18	f	61	CYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3466	0	3377	31	0
1	L	3460	0	3367	31	0
2	B	3154	0	3158	40	0
2	M	3154	0	3158	36	0
3	C	3045	0	3107	36	0
3	N	3046	0	3112	25	0
4	D	1919	0	1867	23	0
4	O	1909	0	1854	17	0
5	E	1512	0	1495	21	0
5	P	1512	0	1495	25	0
6	F	900	0	887	7	0
6	Q	894	0	882	9	0
7	G	624	0	633	11	0
7	R	609	0	614	8	0
8	H	563	0	541	2	0
8	S	563	0	541	5	0
9	J	495	0	489	9	0
9	U	495	0	489	8	0
10	K	429	0	430	3	0
10	V	438	0	443	4	0
11	T	554	0	590	15	0
12	I	828	0	807	12	0
13	a	4021	0	3997	0	0
14	b	1817	0	1822	0	0
15	c	2111	0	2047	0	0
16	d	1195	0	1161	0	0
17	e	842	0	838	0	0
18	f	727	0	703	0	0
19	g	605	0	570	0	0
20	h	654	0	622	0	0
21	i	572	0	596	0	0
22	k	383	0	367	0	0
23	l	380	0	378	0	0
24	m	311	0	329	0	0
25	A	23	0	20	1	0
25	C	35	0	44	0	0
25	D	32	0	38	1	0
25	F	37	0	48	1	0
25	G	42	0	61	0	0
25	L	23	0	20	1	0
25	N	37	0	48	0	0
25	O	23	0	20	0	0
25	R	51	0	82	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
25	a	62	0	72	0	0
25	b	57	0	62	0	0
25	c	45	0	67	0	0
25	g	25	0	24	0	0
25	k	27	0	28	0	0
26	A	46	0	36	1	0
26	D	56	0	56	0	0
26	G	42	0	28	2	0
26	L	46	0	36	1	0
26	O	57	0	58	0	0
26	R	41	0	26	0	0
27	C	86	0	60	5	0
27	N	86	0	60	3	0
28	D	43	0	30	2	0
28	O	43	0	30	2	0
29	E	4	0	0	0	0
29	P	4	0	0	0	0
30	E	35	0	44	0	0
30	G	78	0	104	3	0
30	U	24	0	22	0	0
30	a	43	0	60	0	0
31	G	44	0	63	0	0
32	a	1	0	0	0	0
33	a	1	0	0	0	0
34	a	120	0	108	0	0
35	b	1	0	0	0	0
36	b	2	0	0	0	0
37	f	1	0	0	0	0
All	All	48610	0	48221	336	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 336 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:P:156:TYR:HB2	5:P:165:TYR:HB2	1.69	0.74
3:C:100:ARG:HH12	27:C:402:HEM:HBD1	1.58	0.68
3:N:98:VAL:HG22	27:N:402:HEM:HBC2	1.79	0.64
8:H:19:ARG:HG2	8:H:63:ARG:HD3	1.81	0.63
5:P:99:ARG:HH21	5:P:101:ARG:HD2	1.64	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/446 (100%)	426 (96%)	18 (4%)	0	100	100
1	L	443/446 (99%)	428 (97%)	15 (3%)	0	100	100
2	B	418/439 (95%)	403 (96%)	15 (4%)	0	100	100
2	M	418/439 (95%)	402 (96%)	16 (4%)	0	100	100
3	C	378/381 (99%)	367 (97%)	11 (3%)	0	100	100
3	N	378/381 (99%)	368 (97%)	10 (3%)	0	100	100
4	D	239/241 (99%)	237 (99%)	2 (1%)	0	100	100
4	O	238/241 (99%)	234 (98%)	4 (2%)	0	100	100
5	E	194/196 (99%)	188 (97%)	6 (3%)	0	100	100
5	P	194/196 (99%)	187 (96%)	7 (4%)	0	100	100
6	F	100/110 (91%)	100 (100%)	0	0	100	100
6	Q	99/110 (90%)	98 (99%)	1 (1%)	0	100	100
7	G	72/81 (89%)	66 (92%)	6 (8%)	0	100	100
7	R	70/81 (86%)	67 (96%)	3 (4%)	0	100	100
8	H	66/76 (87%)	64 (97%)	2 (3%)	0	100	100
8	S	66/76 (87%)	64 (97%)	2 (3%)	0	100	100
9	J	58/63 (92%)	56 (97%)	2 (3%)	0	100	100
9	U	58/63 (92%)	56 (97%)	2 (3%)	0	100	100
10	K	50/56 (89%)	48 (96%)	2 (4%)	0	100	100
10	V	51/56 (91%)	48 (94%)	3 (6%)	0	100	100
11	T	76/78 (97%)	67 (88%)	9 (12%)	0	100	100
12	I	109/113 (96%)	99 (91%)	10 (9%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
13	a	512/514 (100%)	498 (97%)	14 (3%)	0	100	100
14	b	225/227 (99%)	209 (93%)	16 (7%)	0	100	100
15	c	257/261 (98%)	248 (96%)	9 (4%)	0	100	100
16	d	142/147 (97%)	133 (94%)	9 (6%)	0	100	100
17	e	102/109 (94%)	99 (97%)	3 (3%)	0	100	100
18	f	93/99 (94%)	85 (91%)	8 (9%)	0	100	100
19	g	73/85 (86%)	68 (93%)	5 (7%)	0	100	100
20	h	77/85 (91%)	74 (96%)	3 (4%)	0	100	100
21	i	70/75 (93%)	68 (97%)	2 (3%)	0	100	100
22	k	47/56 (84%)	45 (96%)	2 (4%)	0	100	100
23	l	44/47 (94%)	43 (98%)	1 (2%)	0	100	100
24	m	41/46 (89%)	39 (95%)	2 (5%)	0	100	100
All	All	5902/6120 (96%)	5682 (96%)	220 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/373 (100%)	370 (99%)	3 (1%)	81	93
1	L	372/373 (100%)	370 (100%)	2 (0%)	88	95
2	B	330/344 (96%)	329 (100%)	1 (0%)	92	96
2	M	330/344 (96%)	330 (100%)	0	100	100
3	C	332/333 (100%)	332 (100%)	0	100	100
3	N	332/333 (100%)	330 (99%)	2 (1%)	86	94
4	D	206/206 (100%)	206 (100%)	0	100	100
4	O	205/206 (100%)	205 (100%)	0	100	100
5	E	166/166 (100%)	166 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	P	166/166 (100%)	166 (100%)	0	100	100
6	F	94/98 (96%)	93 (99%)	1 (1%)	73	88
6	Q	93/98 (95%)	93 (100%)	0	100	100
7	G	67/73 (92%)	67 (100%)	0	100	100
7	R	66/73 (90%)	66 (100%)	0	100	100
8	H	65/72 (90%)	65 (100%)	0	100	100
8	S	65/72 (90%)	65 (100%)	0	100	100
9	J	51/54 (94%)	51 (100%)	0	100	100
9	U	51/54 (94%)	51 (100%)	0	100	100
10	K	42/46 (91%)	42 (100%)	0	100	100
10	V	43/46 (94%)	42 (98%)	1 (2%)	50	78
11	T	58/58 (100%)	58 (100%)	0	100	100
12	I	79/95 (83%)	78 (99%)	1 (1%)	69	87
13	a	425/425 (100%)	423 (100%)	2 (0%)	88	95
14	b	210/210 (100%)	210 (100%)	0	100	100
15	c	225/227 (99%)	225 (100%)	0	100	100
16	d	127/128 (99%)	127 (100%)	0	100	100
17	e	91/95 (96%)	91 (100%)	0	100	100
18	f	81/83 (98%)	81 (100%)	0	100	100
19	g	62/67 (92%)	61 (98%)	1 (2%)	62	84
20	h	70/75 (93%)	70 (100%)	0	100	100
21	i	54/56 (96%)	54 (100%)	0	100	100
22	k	39/46 (85%)	39 (100%)	0	100	100
23	l	39/40 (98%)	39 (100%)	0	100	100
24	m	33/34 (97%)	33 (100%)	0	100	100
All	All	5042/5169 (98%)	5028 (100%)	14 (0%)	92	96

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	N	267	HIS
3	N	331	ASN
19	g	54	ARG

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
13	a	265	LYS
13	a	331	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
7	R	23	GLN
13	a	331	ASN
23	l	43	GLN
2	B	343	GLN
1	A	339	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 43 ligands modelled in this entry, 4 are monoatomic - leaving 39 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
25	3PE	a	607	-	27,27,50	0.41	0	30,32,55	0.49	0
30	PC1	a	606	-	42,42,53	0.33	0	48,50,61	0.34	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
25	3PE	g	101	-	24,24,50	0.45	0	27,29,55	0.70	1 (3%)
27	HEM	N	402	3	27,50,50	1.01	2 (7%)	17,82,82	1.32	2 (11%)
25	3PE	O	302	-	22,22,50	0.44	0	25,27,55	0.44	0
31	TGL	G	104	-	43,43,62	0.21	0	46,46,65	0.25	0
34	HEA	a	603	13	44,67,67	1.38	7 (15%)	37,103,103	2.93	17 (45%)
26	CDL	A	502	-	45,45,99	0.43	0	51,57,111	0.40	0
25	3PE	b	304	-	27,27,50	0.39	0	30,32,55	0.41	0
25	3PE	A	501	-	22,22,50	0.46	0	25,27,55	0.72	1 (4%)
27	HEM	C	401	3	27,50,50	1.00	1 (3%)	17,82,82	1.26	2 (11%)
27	HEM	C	402	3	27,50,50	1.04	2 (7%)	17,82,82	1.53	4 (23%)
26	CDL	D	301	-	55,55,99	0.39	0	61,67,111	0.35	0
26	CDL	O	301	-	56,56,99	0.38	0	62,68,111	0.33	0
28	HEC	D	303	4	26,50,50	2.37	3 (11%)	18,82,82	1.82	5 (27%)
25	3PE	k	101	-	26,26,50	0.40	0	29,31,55	0.38	0
26	CDL	R	101	-	40,40,99	0.47	0	46,52,111	0.60	1 (2%)
28	HEC	O	303	4	26,50,50	2.42	4 (15%)	18,82,82	1.70	4 (22%)
34	HEA	a	604	13	44,67,67	1.42	7 (15%)	37,103,103	3.00	14 (37%)
29	FES	E	201	5	0,4,4	-	-	-	-	-
25	3PE	G	102	-	41,41,50	0.34	0	44,46,55	0.31	0
25	3PE	D	302	-	31,31,50	0.38	0	34,36,55	0.36	0
30	PC1	G	105	-	42,42,53	0.32	0	48,50,61	0.30	0
36	CUA	b	303	14	0,1,1	-	-	-	-	-
27	HEM	N	401	3	27,50,50	0.98	1 (3%)	17,82,82	1.25	0
30	PC1	U	101	-	23,23,53	0.45	0	29,31,61	0.68	1 (3%)
25	3PE	F	201	-	36,36,50	0.35	0	39,41,55	0.31	0
26	CDL	G	101	-	41,41,99	0.44	0	47,53,111	0.37	0
30	PC1	E	202	-	34,34,53	0.35	0	40,42,61	0.34	0
25	3PE	c	301	-	44,44,50	0.32	0	47,49,55	0.34	0
25	3PE	R	102	-	50,50,50	0.32	0	53,55,55	0.30	0
29	FES	P	201	5	0,4,4	-	-	-	-	-
25	3PE	b	302	-	28,28,50	0.40	0	31,33,55	0.35	0
25	3PE	C	403	-	34,34,50	0.36	0	37,39,55	0.35	0
26	CDL	L	502	-	45,45,99	0.44	0	51,57,111	0.57	1 (1%)
30	PC1	G	103	-	34,34,53	0.36	0	40,42,61	0.69	1 (2%)
25	3PE	N	403	-	36,36,50	0.35	0	39,41,55	0.32	0
25	3PE	a	605	-	33,33,50	0.38	0	36,38,55	0.61	1 (2%)
25	3PE	L	501	-	22,22,50	0.44	0	25,27,55	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
25	3PE	a	607	-	-	12/31/31/54	-
30	PC1	a	606	-	-	10/46/46/57	-
25	3PE	g	101	-	-	10/28/28/54	-
27	HEM	N	402	3	-	0/6/54/54	-
25	3PE	O	302	-	-	5/26/26/54	-
31	TGL	G	104	-	-	6/46/46/65	-
34	HEA	a	603	13	3/3/7/16	7/24/76/76	-
26	CDL	A	502	-	-	18/56/56/110	-
25	3PE	b	304	-	-	12/31/31/54	-
25	3PE	A	501	-	-	9/26/26/54	-
27	HEM	C	401	3	-	1/6/54/54	-
27	HEM	C	402	3	-	0/6/54/54	-
26	CDL	D	301	-	-	17/66/66/110	-
26	CDL	O	301	-	-	18/67/67/110	-
28	HEC	D	303	4	-	0/6/54/54	-
34	HEA	a	604	13	3/3/7/16	6/24/76/76	-
25	3PE	k	101	-	-	9/30/30/54	-
26	CDL	R	101	-	-	12/51/51/110	-
28	HEC	O	303	4	-	2/6/54/54	-
29	FES	E	201	5	-	-	0/1/1/1
25	3PE	G	102	-	-	11/45/45/54	-
25	3PE	D	302	-	-	11/35/35/54	-
30	PC1	G	105	-	-	13/46/46/57	-
27	HEM	N	401	3	-	1/6/54/54	-
30	PC1	U	101	-	-	10/27/27/57	-
25	3PE	F	201	-	-	16/40/40/54	-
26	CDL	G	101	-	-	14/52/52/110	-
30	PC1	E	202	-	-	11/38/38/57	-
25	3PE	c	301	-	-	10/48/48/54	-
25	3PE	R	102	-	-	14/54/54/54	-
30	PC1	G	103	-	-	11/37/37/57	-
25	3PE	b	302	-	-	9/32/32/54	-
25	3PE	C	403	-	-	3/38/38/54	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
26	CDL	L	502	-	-	21/56/56/110	-
29	FES	P	201	5	-	-	0/1/1/1
25	3PE	N	403	-	-	8/40/40/54	-
25	3PE	a	605	-	-	9/37/37/54	-
25	3PE	L	501	-	-	8/26/26/54	-

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
28	O	303	HEC	C3B-C2B	-6.59	1.33	1.40
28	D	303	HEC	C3B-C2B	-6.53	1.33	1.40
28	D	303	HEC	C3C-C2C	-6.45	1.34	1.40
28	O	303	HEC	C3C-C2C	-6.43	1.34	1.40
28	O	303	HEC	C3D-C2D	5.35	1.53	1.37

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
34	a	604	HEA	CMC-C2C-C3C	7.41	138.54	124.68
34	a	603	HEA	CMC-C2C-C3C	7.23	138.20	124.68
34	a	604	HEA	CMC-C2C-C1C	-6.56	118.38	128.46
34	a	603	HEA	CMC-C2C-C1C	-6.44	118.56	128.46
34	a	604	HEA	CMB-C2B-C1B	-6.27	118.82	128.46

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
34	a	603	HEA	NA
34	a	603	HEA	NB
34	a	603	HEA	ND
34	a	604	HEA	NA
34	a	604	HEA	NB

5 of 334 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
25	A	501	3PE	C1-O11-P-O12
25	A	501	3PE	O13-C11-C12-N
25	C	403	3PE	C11-O13-P-O14
25	D	302	3PE	C1-O11-P-O12
25	D	302	3PE	C1-O11-P-O13


There are no ring outliers.


16 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
27	N	402	HEM	1	0
26	A	502	CDL	1	0
25	A	501	3PE	1	0
27	C	401	HEM	2	0
27	C	402	HEM	3	0
28	D	303	HEC	2	0
28	O	303	HEC	2	0
25	D	302	3PE	1	0
30	G	105	PC1	2	0
27	N	401	HEM	2	0
25	F	201	3PE	1	0
26	G	101	CDL	2	0
25	R	102	3PE	2	0
26	L	502	CDL	1	0
30	G	103	PC1	1	0
25	L	501	3PE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12702. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 92



Y Index: 91




Z Index: 104


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is  $299 \text{ nm}^3$ ; this corresponds to an approximate mass of 270 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.


## 7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>



## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.18	3.90	3.29
Unmasked-calculated*	-	-	-

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12702 and PDB model 7O37. Per-residue inclusion information can be found in section 3 on page 17.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.